

Data collection

Siemens P4 diffractometer 1933 reflections with
 2 θ/ω scans $I > 2\sigma(I)$
 Absorption correction: $R_{\text{int}} = 0.013$
 empirical via ψ scans $\theta_{\text{max}} = 24.99^\circ$
 (SHELXL93; Sheldrick, 1993) $h = 0 \rightarrow 7$
 $T_{\text{min}} = 0.666$, $T_{\text{max}} = 0.760$ $k = 0 \rightarrow 13$
 2409 measured reflections $l = -22 \rightarrow 22$
 2183 independent reflections 3 standard reflections
 every 97 reflections
 intensity decay: 3.19%

Refinement

Refinement on F^2 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $R[F^2 > 2\sigma(F^2)] = 0.023$ $\Delta\rho_{\text{max}} = 0.253 \text{ e } \text{\AA}^{-3}$
 $wR(F^2) = 0.064$ $\Delta\rho_{\text{min}} = -0.278 \text{ e } \text{\AA}^{-3}$
 $S = 1.052$ Extinction correction: none
 2182 reflections Scattering factors from
 191 parameters *International Tables for*
 All H atoms refined *Crystallography* (Vol. C)
 $w = 1/[\sigma^2(F_o^2) + (0.0286P)^2$
 $+ 0.4364P]$
 where $P = (F_o^2 + 2F_c^2)/3$

Table 2. Selected geometric parameters (\AA , $^\circ$) for (II)

Pd—S2	2.2496 (6)	S2—C2	1.712 (2)
Pd—S1	2.2501 (6)	C1—C2	1.400 (3)
S1—C1	1.695 (2)		
S2 ⁱ —Pd—S1	92.08 (2)	C2—S2—Pd	105.93 (8)
S2—Pd—S1	87.92 (2)	C2—C1—S1	121.6 (2)
C1—S1—Pd	105.32 (8)	C1—C2—S2	119.2 (2)

Symmetry code: (i) $-x, -y, -z$.

For both compounds, data collection: XSCANS (Siemens, 1994a); cell refinement: XSCANS; data reduction: XSCANS; program(s) used to solve structures: SHELXS86 (Sheldrick, 1990); program(s) used to refine structures: SHELXL93 (Sheldrick, 1993); molecular graphics: XP (Siemens, 1994b); software used to prepare material for publication: SHELXL93.

This work has been generously supported by the National Science Council, Taiwan.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: TA1234). Services for accessing these data are described at the back of the journal.

References

- Alcàcer, L. & Novais, H. (1982). In *Extended Linear Chain Compounds*, Vol. 3. New York: Plenum.
- Broderick, W. E., Thompson, J. A. & Hoffman, B. M. (1991). *Inorg. Chem.* **30**, 2958–2960.
- Bünder, W. & Weiss, E. (1975). *J. Organomet. Chem.* **92**, 65–68.
- Clemenson, P. I. (1990). *Coord. Chem. Rev.* **106**, 171–203.
- Davison, A., Edelstein, N., Holm, R. H. & Maki, H. (1963). *Inorg. Chem.* **2**, 1227.
- Fang, Q., You, X.-Z., Cai, J.-H. & He, M.-Y. (1993). *Acta Cryst.* **C49**, 1347–1350.
- Miller, J. S. & Epstein, A. J. (1995). *Molecular Magnetism: From Molecular Assemblies to the Devices*, edited by E. Coronado, P. Delhaès, D. Gatteschi & J. S. Miller, pp. 379–414. Amsterdam: Kluwer Academic Publishers.
- Sartain, D. & Truter, M. R. (1967). *J. Chem. Soc. A*, pp. 1264–1274.

- Schläpfer, C. W. & Nakamoto, K. (1975). *Inorg. Chem.* **14**, 1338–1344.
- Schrauzer, G. N. & Mayweg, V. P. (1965). *J. Am. Chem. Soc.* **87**, 3585–3592.
- Sheldrick, G. M. (1990). *Acta Cryst.* **A46**, 467–473.
- Sheldrick, G. M. (1993). *SHELXL93. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.
- Siemens (1994a). *XSCANS. X-ray Single Crystal Analysis System*. Version 2.1. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Siemens (1994b). *XP. Molecular Graphics Program*. Version 5.03. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.

Acta Cryst. (1999). **C55**, 1072–1075

A linear trinuclear Cd^{II}–Cd^{II}–Cd^{II} complex with a μ -acetato bridge: bis{(μ -acetato)[μ -bis(salicylidene)-1,3-propanediaminato]-cadmium(II)}cadmium(II)

ORHAN ATAKOL,^a MECIT AKSU,^a FILIZ ERCAN,^b CENGİZ ARICI,^b M. NAWAZ TAHIR^b AND DİNÇER ÜLKÜ^b

^aDepartment of Chemistry, Ankara University, Tandogan 06532, Ankara, Turkey, and ^bDepartment of Engineering Physics, Hacettepe University, Beytepe 06100, Ankara, Turkey. E-mail: dulku@eti.cc.hun.edu.tr

(Received 23 September 1998; accepted 22 March 1999)


Abstract

The title complex, bis(μ -acetato)-1:2 $\kappa^2O:O'$;2:3 $\kappa^2O:O'$ -bis{ μ -2,2'-[1,3-propanediylbis(nitrilomethylidene)]diphenolato}-1 κ^4N,N',O,O' :2 κ^2O,O' ;3 κ^4N,N',O,O' :2 κ^2O,O' -tricadmium(II), [Cd{Cd(C₁₇H₁₆N₂O₂)(CH₃CO₂)₂}₂], consists of a doubly oxygen-bridged linear homotrimeric complex. The outer Cd^{II} ions, related by an inversion centre, have an irregular square-pyramidal coordination. The central Cd^{II} ion has an irregular octahedral coordination and lies on an inversion centre. The metal–metal bridging distance is 3.2582 (3) Å.

Comment

The structures and syntheses of several oxygen-bridged dimeric and trimeric linear homo- or heteronuclear metal complexes have been studied in our laboratories recently; examples include [Cu₂(C₁₁H₁₀CINO₂)₂] (Tahir *et al.*, 1996), [Cu₂(C₁₄H₁₁NO₂)₂] (Ülkü, Ercan, Atakol, Ercan & Gencer, 1997), [Ni₃{(SALPD)(CH₃CO₂)-[(CH₃)₂SO]}₂] (Ülkü, Ercan, Atakol & Dinçer, 1997), [Cd{Ni(SALPD)(CH₃CO₂)[(CH₃)₂NCHO]}₂] (Ülkü, Tahir *et al.*, 1997), [Cu₂(C₁₂H₁₃NO₂)₂] (Atakol *et al.*, 1997), [Cu₂(C₁₂H₁₃NO₂)₂] (Ülkü *et al.*, 1998), [Ni₂{Cu-

(SALPD)(NO₂)₂[(CH₃)₂NCHO]}₂·2[(CH₃)₂NCHO] (Tahir *et al.*, 1998), [Mn{Ni(SALPD)(CH₃CO₂)[(CH₃)₂NCHO]}₂] (Ercan & Atakol, 1998), [Cd{Cu(SALPD)(CH₃CO₂)₂}]·C₄H₈O₂ (Ercan *et al.*, 1998), [Cu₃{(SALPD)(CH₃CO₂)₂}]·C₄H₈O₂ (Atakol *et al.*, 1999), [Cu{Zn(C₂₃H₂₁I₂N₃O₂)}] (Ercan *et al.*, 1999) and [Ni{Zn(C₃₁H₃₄I₂N₄O₂)}] (Arıcı *et al.*, 1999), where H₂SALPD is *N,N'*-bis(salicylidene)-1,3-propanediamine {alternative name: 2,2'-[1,3-propanediylbis-(nitrilomethylidene)]diphenol}. Structural studies of some other oxygen-bridged homonuclear complexes have been reported previously by other workers (Kato *et al.*, 1964; Barclay & Hoskins, 1965; Butcher & Sinn, 1976; Kato & Muto, 1988). We report here a new linear homonuclear trimeric complex, [Cd₃{(SALPD)(CH₃CO₂)₂}]₂, (I).



The monoclinic unit cell of (I) contains two centrosymmetric trimer complexes, whose central Cd^{II} ion is located on the inversion centre at the origin. The

coordination around the central Cd^{II} ion is an irregular octahedron, involving four bridging O atoms from two SALPD²⁻ ligands in the equatorial plane and an O atom from each of the two bridging acetate groups in the apical positions. The inversion-related outer Cd^{II} ions have irregular square-pyramidal coordination polyhedra, with the four equatorial positions occupied by the two N and two O atoms of the SALPD²⁻ ligand. There is an O atom from each bridging acetate group in the axial positions of these irregular square-pyramidal polyhedra.

The Cd···Cd distance is 3.2582(3) Å; this is longer than the Cu···Cd distance [3.244(3) Å] in the similar structure described by Ercan *et al.* (1998). The SALPD²⁻ ligand in (I) is not planar. The dihedral angles between the best plane of the salicylidimine residues (N1, C1–C7, O1 and N2, C11–C17, O2) and the equatorial plane (O1, O2, N1, N2) are 143.4(1) and 40.7(1)°, respectively. The dihedral angle between the equatorial planes of the neighbouring polyhedra [O1, O2, N1, N2 and O1, O2, O1ⁱ, O2ⁱ; symmetry code: (i) $-x, -y, -z$] is 13.7(1)°. The outer Cd^{II} ion is 0.7407(1) Å out of its equatorial plane. The Cd1—O1, Cd1—O2 and Cd1—O4 bond lengths are 2.269(3), 2.297(3) and 2.353(3) Å, respectively, while the Cd—O distances were 2.280(3), 2.310(3) and 2.238(2) Å, respectively, in our previous investigation (Ercan *et al.*, 1998). The average Cd2—O bond length in (I) is 2.207(3) Å, and the Cd2—N1 and Cd2—N2 bond lengths are 2.258(4) and 2.236(4) Å, respectively. The chelate ring formed by atoms Cd2, N1, C8–C10 and N2 has a chair conformation. The distances




Fig. 1. PLATON (Spek, 1998) drawing of (I) with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level and H atoms are shown as small circles of arbitrary radii [symmetry code: (i) $-x, -y, -z$].

of the two *para*-positioned chair atoms, Cd2 and C9, from the plane of the other four atoms are $-1.035(6)$ and $0.697(6)$ Å, respectively. Although the type of metal ions and some of the coordinating groups of the metal complexes reported by our previous investigations are different from the case discussed here, the bond lengths, angles and dihedral angles of these structures are similar.

Experimental

N,N'-Bis(salicylidene)-1,3-propanediamine (0.282 g, 1 mmol) was dissolved in 75 ml hot methanol. Next, a solution of Cd(CH₃CO₂)₂ (0.335 g, 1.5 mmol) in 30 ml hot methanol was added slowly. The resulting mixture was set aside for 1 d and the yellow crystals which formed were filtered off and dried in air.

Crystal data

[Cd ₃ (C ₁₇ H ₁₆ N ₂ O ₂) ₂ ·(C ₂ H ₃ O ₂) ₂]	Mo K α radiation
$M_r = 1015.95$	$\lambda = 0.71073$ Å
Monoclinic	Cell parameters from 25 reflections
$P2_1/c$	$\theta = 12.04\text{--}20.91^\circ$
$a = 10.6433(12)$ Å	$\mu = 1.74$ mm ⁻¹
$b = 20.6568(13)$ Å	$T = 295$ K
$c = 8.5187(11)$ Å	Prismatic
$\beta = 91.762(2)^\circ$	$0.30 \times 0.20 \times 0.15$ mm
$V = 1872.0(3)$ Å ³	Yellow
$Z = 2$	
$D_x = 1.802$ Mg m ⁻³	
D_m not measured	

Data collection

Enraf-Nonius CAD-4 diffractometer	$R_{\text{int}} = 0.014$
$\omega/2\theta$ scans	$\theta_{\text{max}} = 25.08^\circ$
Absorption correction: empirical via ψ scans (Fair, 1990)	$h = -10 \rightarrow 0$
$T_{\text{min}} = 0.614$, $T_{\text{max}} = 0.770$	$k = -24 \rightarrow 0$
3669 measured reflections	$l = -12 \rightarrow 12$
3195 independent reflections	3 standard reflections
2772 reflections with $I > 1\sigma(I)$	frequency: 120 min
	intensity decay: -0.78%

Refinement

Refinement on F	$(\Delta/\sigma)_{\text{max}} < 0.001$
$R = 0.028$	$\Delta\rho_{\text{max}} = 0.48$ e Å ⁻³
$wR = 0.037$	$\Delta\rho_{\text{min}} = -0.20$ e Å ⁻³
$S = 0.90$	Extinction correction: none
2772 reflections	Scattering factors from <i>International Tables for X-ray Crystallography</i> (Vol. IV)
241 parameters	
H atoms: see below	
$w = 1/[\sigma F^2 + (0.02F)^2 + 0.4]$	

Table 1. Selected geometric parameters (Å, °)

Cd1...Cd2	3.2582(3)	O1—C1	1.318(5)
Cd1—O1	2.269(3)	O2—C17	1.319(5)
Cd1—O2	2.297(3)	O3—C18	1.271(6)

Cd1—O4	2.353(4)	O4—C18	1.229(6)
Cd2—O1	2.223(3)	N1—C7	1.284(6)
Cd2—O2	2.225(3)	N1—C8	1.475(6)
Cd2—O3	2.173(3)	N2—C10	1.472(6)
Cd2—N1	2.258(4)	N2—C11	1.273(6)
Cd2—N2	2.236(4)		
Cd1—Cd2—O1	44.07(8)	Cd2—O1—Cd1	93.0(1)
Cd1—Cd2—O2	44.78(8)	Cd2—O1—C1	130.3(3)
Cd1—Cd2—O3	93.0(1)	Cd1—O1—C1	132.6(3)
Cd1—Cd2—N1	127.3(1)	Cd2—O2—Cd1	92.2(1)
Cd1—Cd2—N2	128.0(1)	Cd2—O2—C17	129.4(3)
Cd2—Cd1—O1	42.96(8)	Cd1—O2—C17	132.6(3)
Cd2—Cd1—O2	43.04(8)	Cd1—O4—C18	150.7(3)
Cd2—Cd1—O4	60.30(9)	Cd2—O3—C18	110.8(3)
O1—Cd2—O2	80.8(1)	Cd2—N1—C7	125.8(3)
O1—Cd2—O3	105.1(1)	Cd2—N1—C8	116.6(3)
O1—Cd2—N1	83.6(1)	C7—N1—C8	117.5(4)
O1—Cd2—N2	141.5(1)	Cd2—N2—C10	114.8(3)
O2—Cd2—O3	110.0(1)	Cd2—N2—C11	126.3(3)
O2—Cd2—N1	140.8(1)	C10—N2—C11	118.8(4)
O2—Cd2—N2	83.4(1)	O1—C1—C2	119.5(4)
O3—Cd2—N1	108.6(1)	O1—C1—C6	123.3(4)
O3—Cd2—N2	113.3(1)	C1—C6—C7	126.5(4)
N1—Cd2—N2	87.0(1)	N1—C7—C6	128.8(4)
O1—Cd1—O2	78.3(1)	N1—C8—C9	112.9(4)
O1—Cd1—O4	82.1(1)	N2—C10—C9	112.4(4)
O2—Cd1—O4	81.9(1)		

All non-H atoms were refined with anisotropic displacement parameters. H atoms H81, H82, H91, H92, H101, H102, H191, H192 and H193 were taken from difference maps, while the other H atoms were placed geometrically 0.95 Å from their parent C atoms, and then a riding model was used for all H atoms, with $U_{\text{iso}}(\text{H}) = 1.3U_{\text{eq}}(\text{C})$.

Data collection: CAD-4 EXPRESS (Enraf-Nonius, 1993). Data reduction: MolEN (Fair, 1990). Program(s) used to solve structure: MolEN. Program(s) used to refine structure: MolEN. Molecular graphics: PLATON (Spek, 1998). Software used to prepare material for publication: MolEN.

The authors wish to acknowledge the purchase of the CAD-4 diffractometer under Grant DPT/TBAG1 of the Scientific and Technical Research Council of Turkey.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: FR1176). Services for accessing these data are described at the back of the journal.

References

- Arci, C., Ercan, F., Atakol, O., Akay, A. & Ülkü, D. (1999). *Acta Cryst.* **C55**, 928–930.
- Atakol, O., Arci, C., Ercan, F. & Ülkü, D. (1999). *Acta Cryst.* **C55**, 511–513.
- Atakol, O., Ercan, F., Ülkü, D. & Yılmaz, N. (1997). *Anal. Sci.* **13**, 1051–1052.
- Barclay, G. A. & Hoskins, B. F. (1965). *J. Chem. Soc.* pp. 1979–1991.
- Butcher, R. J. & Sinn, E. (1976). *Inorg. Chem.* **15**, 1604–1608.
- Enraf-Nonius (1993). CAD-4 EXPRESS. Version 1.1. Enraf-Nonius, Delft, The Netherlands.
- Ercan, F., Arci, C., Akay, A., Atakol, O. & Ülkü, D. (1999). *Acta Cryst.* **C55**, 925–928.
- Ercan, F. & Atakol, O. (1998). *Acta Cryst.* **C54**, 1268–1270.
- Ercan, F., Ülkü, D., Atakol, O. & Dinçer, F. N. (1998). *Acta Cryst.* **C54**, 1787–1790.
- Fair, C. K. (1990). *MolEN. An Interactive Intelligent System for Crystal Structure Analysis*. Enraf-Nonius, Delft, The Netherlands.
- Kato, M., Jonassen, H. B. & Fanning, J. (1964). *Chem. Rev.* **64**, 99–128.

- Kato, M. & Muto, Y. (1988). *Coord. Chem. Rev.* **92**, 45–83.
 Spek, A. L. (1998). *PLATON. Molecular Geometry Program*. University of Utrecht, The Netherlands.
 Tahir, M. N., Ülkü, D., Atakol, O. & Akay, A. (1996). *Acta Cryst.* **C52**, 2676–2678.
 Tahir, M. N., Ülkü, D., Atakol, O. & Çakırcer, O. (1998). *Acta Cryst.* **C54**, 468–470.
 Ülkü, D., Ancia, C., Tahir, M. N., Atakol, O. & Tastekin, M. (1998). *Anal. Sci.* **14**, 455–456.
 Ülkü, D., Ercan, F., Atakol, O. & Dinçer, F. N. (1997). *Acta Cryst.* **C53**, 1056–1057.
 Ülkü, D., Ercan, F., Atakol, O., Ercan, I. & Gencer, A. (1997). *Acta Cryst.* **C53**, 179–181.
 Ülkü, D., Tahir, M. N., Atakol, O. & Nazir, H. (1997). *Acta Cryst.* **C53**, 872–874.

Acta Cryst. (1999). **C55**, 1075–1079

Three anionic polymeric networks: sodium(I)–ruthenium(III), sodium(I)– rhodium(III) and sodium(I)–aluminium(III) μ -oxalato complexes with tris(2,2′-bi- pyridine)ruthenium(II) cations

RENE PELLAUX,^a SILVIO DECURTINS^a AND HELMUT W. SCHMALLE^b

^aDepartement für Chemie und Biochemie, Universität Bern, Freiestrasse 3, CH-3012 Bern, Switzerland, and
^bInstitute of Inorganic Chemistry, University of Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland.
 E-mail: schmalle@aci.unizh.ch


(Received 26 May 1998; accepted 26 February 1999)

Abstract

The crystal structures of $[\text{Ru}(\text{bpy})_3]_n^{2+}[\text{Na}M(\text{ox})_3]_n^{2n-}$, with $M = \text{Ru}^{\text{III}}$ [tris(2,2′-bipyridine)ruthenium(II) tris(μ -oxalato)ruthenium(III)sodium(I), (1)], $M = \text{Rh}^{\text{III}}$ [tris(2,2′-bipyridine)ruthenium(II) tris(μ -oxalato)rhodium(III)sodium(I), (2)] and $M = \text{Al}^{\text{III}}$ [tris(2,2′-bipyridine)ruthenium(II) tris(μ -oxalato)aluminium(III)sodium(I), (3)], where bpy is 2,2′-bipyridine ($\text{C}_{10}\text{H}_8\text{N}_2$) and ox is oxalate (C_2O_4), represent examples of a specific polymeric three-dimensional network topology. These networks, formally built up from $[M(\text{ox})_3/2]$ subunits and $[\text{Ru}(\text{bpy})_3]^{2+}$ counter-ions which template the polymeric structure, are chiral. The absolute configurations of the measured crystals have been determined as the Λ form for (1), and as the Δ form for (2) and (3). The coordinated metal ions form trigonally distorted octahedra. The average distances are 2.060 (2) Å for $\text{Ru}^{\text{II}}-\text{N}$ in the three $[\text{Ru}(\text{bpy})_3]^{2+}$ counter-ions, 2.338 (3) Å for $\text{Na}^{\text{I}}-\text{O}$, 2.031 (2) Å for $\text{Ru}^{\text{III}}-\text{O}$, 2.005 (2) Å for $\text{Rh}^{\text{III}}-\text{O}$ and 1.890 (3) Å for $\text{Al}^{\text{III}}-\text{O}$.

Comment

The anionic oxalate molecule can act as a mono-, bi-, tri- or tetradentate ligand, but is also well known for its ability to form bridged polynuclear complexes (Scott *et al.*, 1973). In particular, several two-dimensional heterometallic layer compounds with network stoichiometries $[M^{\text{II}}M^{\text{III}}(\text{ox})_3]_n^{n-}$ (Tamaki *et al.*, 1992; Decurtins, Schmalte, Oswald *et al.*, 1994; Pellaux *et al.*, 1997), as well as three-dimensional homometallic network compounds with stoichiometries $[M_2^{\text{II}}(\text{ox})_3]_n^{2n-}$ (Decurtins, Schmalte, Schneuwly *et al.*, 1994; Decurtins *et al.*, 1996), have been reported. Both structure types have been realised for many different transition metals and they are of interest in the field of molecule-based magnets due to the ability of the oxalato ligand to act as mediator for magnetic exchange interactions. In addition, a three-dimensional heterometallic structure type with stoichiometries $[M^{\text{I}}M^{\text{III}}(\text{ox})_3]_n^{2n-}$, of which three examples are presented in this report, has been studied mainly because of its specific photophysical properties (Hauser *et al.*, 1996; von Arx *et al.*, 1996). For a general description of the $[M^{\text{I}}M^{\text{III}}(\text{ox})_3]_n^{2n-}$ structures, these anionic networks may be considered as built up from formal $[M(\text{ox})_3/2]$ subunits, each representing three-connected points (Wells, 1984). If all subunits have the same chiral configuration, a well defined three-dimensional three-connected decagon network is formed, where the $[\text{Ru}(\text{bpy})_3]^{2+}$ cations occupy the vacancies of the framework.



In Fig. 1, a stereoplot of the structure of (1) in the Λ configuration is presented. Strands of bridged metal ions in the oxalato network form left-handed helices along the 2_1 axes. According to the space-group symmetry, all metal ions occupy sites with D_3 point-group symmetry (Wyckoff letter *a*). Contrary to the Λ configuration found in the case of (1), the absolute configurations of the coordinated metal ions in the measured crystals of (2) and (3) represent the Δ form; compound (3) is shown in Fig. 2 as an example. The Na—O 3+3 coordination distances range between 2.315 (2) Å in (1) and 2.388 (3) Å in (3). The most pronounced differences within the Na—O distances are observed in (3). Furthermore, these bond lengths are significantly elongated compared with values in (1) and (2). In a similar way, the Na—O distances reported for $[\text{Ni}(\text{bpy})_3]$ -