

Global, regional, and national disability-adjusted life-years (DALYs) for 315 diseases and injuries and healthy life expectancy (HALE), 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015



GBD 2015 DALYs and HALE Collaborators*



Summary

Background Healthy life expectancy (HALE) and disability-adjusted life-years (DALYs) provide summary measures of health across geographies and time that can inform assessments of epidemiological patterns and health system performance, help to prioritise investments in research and development, and monitor progress toward the Sustainable Development Goals (SDGs). We aimed to provide updated HALE and DALYs for geographies worldwide and evaluate how disease burden changes with development.

Methods We used results from the Global Burden of Diseases, Injuries, and Risk Factors Study 2015 (GBD 2015) for all-cause mortality, cause-specific mortality, and non-fatal disease burden to derive HALE and DALYs by sex for 195 countries and territories from 1990 to 2015. We calculated DALYs by summing years of life lost (YLLs) and years of life lived with disability (YLDs) for each geography, age group, sex, and year. We estimated HALE using the Sullivan method, which draws from age-specific death rates and YLDs per capita. We then assessed how observed levels of DALYs and HALE differed from expected trends calculated with the Socio-demographic Index (SDI), a composite indicator constructed from measures of income per capita, average years of schooling, and total fertility rate.

Findings Total global DALYs remained largely unchanged from 1990 to 2015, with decreases in communicable, neonatal, maternal, and nutritional (Group 1) disease DALYs offset by increased DALYs due to non-communicable diseases (NCDs). Much of this epidemiological transition was caused by changes in population growth and ageing, but it was accelerated by widespread improvements in SDI that also correlated strongly with the increasing importance of NCDs. Both total DALYs and age-standardised DALY rates due to most Group 1 causes significantly decreased by 2015, and although total burden climbed for the majority of NCDs, age-standardised DALY rates due to NCDs declined. Nonetheless, age-standardised DALY rates due to several high-burden NCDs (including osteoarthritis, drug use disorders, depression, diabetes, congenital birth defects, and skin, oral, and sense organ diseases) either increased or remained unchanged, leading to increases in their relative ranking in many geographies. From 2005 to 2015, HALE at birth increased by an average of 2·9 years (95% uncertainty interval 2·9–3·0) for men and 3·5 years (3·4–3·7) for women, while HALE at age 65 years improved by 0·85 years (0·78–0·92) and 1·2 years (1·1–1·3), respectively. Rising SDI was associated with consistently higher HALE and a somewhat smaller proportion of life spent with functional health loss; however, rising SDI was related to increases in total disability. Many countries and territories in central America and eastern sub-Saharan Africa had increasingly lower rates of disease burden than expected given their SDI. At the same time, a subset of geographies recorded a growing gap between observed and expected levels of DALYs, a trend driven mainly by rising burden due to war, interpersonal violence, and various NCDs.

Interpretation Health is improving globally, but this means more populations are spending more time with functional health loss, an absolute expansion of morbidity. The proportion of life spent in ill health decreases somewhat with increasing SDI, a relative compression of morbidity, which supports continued efforts to elevate personal income, improve education, and limit fertility. Our analysis of DALYs and HALE and their relationship to SDI represents a robust framework on which to benchmark geography-specific health performance and SDG progress. Country-specific drivers of disease burden, particularly for causes with higher-than-expected DALYs, should inform financial and research investments, prevention efforts, health policies, and health system improvement initiatives for all countries along the development continuum.

Funding Bill & Melinda Gates Foundation.

Copyright © The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY license.

Lancet 2016; 388: 1603–58

This online publication has been corrected. The corrected version first appeared at thelancet.com on January 5, 2017

See [Editorial](#) page 1447

See [Comment](#) pages 1448 and 1450

*Collaborators listed at the end of the Article

Correspondence to:
Prof Christopher J L Murray,
2301 5th Avenue, Suite 600,
Seattle, WA 98121, USA
cjl@uw.edu

Introduction

Summary measures of population health are crucial inputs to guide health system investments and set priorities at global, regional, national, and subnational levels. The Millennium Development Goals (MDGs), which sought to reduce extreme poverty and improve health, expired in 2015, and were replaced by the 2030 Agenda for Sustainable Development, or Sustainable Development Goals (SDGs).¹ The shift from the MDGs to the SDGs reflects a broadening of the global development agenda,^{2,3} expanding to include targets for non-communicable diseases (NCDs) and indicators that consider the interplay of environmental, societal, and economic factors on health.⁴ Within this context, summary population health measures are advantageous because they can easily be used to show progress toward SDG 3—to “ensure healthy lives and promote well-being for all at all ages”—and provide a metric by which comparative progress on other SDGs can be monitored.⁵ Summary measures also provide insights into whether, as societies live longer, they spend more or less of their time with functional health loss, known as the expansion or compression of morbidity, respectively, which has profound implications for societies and the financing of health systems.

Two types of population health summary measures exist: health expectancies and health gaps.⁶ Healthy life expectancy (HALE), which originates from Sullivan,⁷ provides a single summary measure of population health by weighting years lived with a measure of functional

health loss experienced before death. Many health expectancy measures have been proposed, but HALE is the only one that captures a full range of functional health loss.^{8–10} Health gap measures capture differences between a population and some normative standard such as a maximum lifespan in full health. Disability-adjusted life-years (DALY) are a widely used gap measure,^{6,9–11} representing the sum of years of life lost (YLLs) due to premature mortality and years lived with disability (YLDs). YLLs quantify the gap between observed mortality and a normative life expectancy,¹² and YLDs capture the prevalence of conditions that lead to non-fatal health loss while accounting for the severity of those conditions. Health gap measures can be easily disaggregated to examine contributions of relative morbidity and mortality, individual diseases, injuries, and attributable risk factors.

The Global Burden of Diseases, Injuries, and Risk Factors (GBD) study is the most comprehensive source of comparable summary population health measures because of its inclusion of country-level results, uncertainty quantification, and its effort to maximise comparability across geography, time, and across different health conditions. Alternative summary health assessments are not as standardised or comprehensive, with studies reporting only incomplete time-series, no uncertainty measures, or only a subset of countries and causes.^{13–15} WHO published DALY estimates for 2 years (2000 and 2012), with 132 causes and 174 countries and

Research in context

Evidence before this study

Disability-adjusted life-years (DALYs), a summary measure of population health based on estimates of premature mortality and non-fatal health loss, originated from the initial Global Burden of Disease (GBD) study in 1993. DALYs, in combination with other summary measures such as healthy life expectancy (HALE), offer relatively simple yet powerful metrics against which progress and challenges in improving disease burden and extending healthy lifespans can be effectively monitored over time. Published in 2012, GBD 2010 provided updated estimates of DALYs due to 291 causes and HALE in 187 countries from 1990 to 2010. GBD 2013 extended this time series to 2013, with 188 countries, and 306 causes. Novel analyses for quantifying epidemiological transitions were introduced as part of GBD 2013, enabling a comparison of shifts in years of life lost (YLLs) and years lived with disability (YLDs) with increasing levels of development. WHO has produced estimates of DALYs and HALE largely based on GBD 2010 and GBD 2013; however, modifications were implemented for a subset of causes, disability weights, and countries, and a normative life table of 91·9 years at birth was used for calculating YLLs.

Added value of this study

For GBD 2015, we generated estimates of HALE and DALYs for 315 causes by geography, sex, and age group from 1990 to

2015 for 195 countries and territories. We constructed a summary metric referred to as the Socio-demographic Index (SDI) based on measures of income per capita, average years of schooling, and total fertility rate. We estimated SDI for each geography-year, and characterised the average relationship for each age, sex, and cause for DALYs and HALE with SDI. Using these relationships, we calculated expected levels of DALYs, life expectancy, and HALE for each geography over time. We compared observed patterns of both DALYs and HALE with those expected on the basis of SDI, allowing us to explore where health gains exceeded—or lagged behind—corresponding changes in development.

Implications

Since 1990, overall health has improved in most countries, with particularly large gains occurring in the past 10 years. Although improved health means longer lifespans, it also translates to more years of functional health lost. The fraction of overall life expectancy spent in poor health is generally constant or has slightly declined in some countries, a result driven by declines in DALYs due to communicable, maternal, nutritional, and neonatal causes and increases in others, mainly non-communicable diseases. Country-specific drivers of disease burden, particularly when observed DALYs are higher than expected on the basis of SDI, should inform country-specific inquiry and action.

without uncertainty intervals. These estimates were derived primarily from GBD 2010 results, but were modified in 60 countries and for 12 cause groups separately estimated by WHO and UN agencies.^{13,16,17} WHO applied the same approach for GBD 2013 results and used their own life tables to produce HALE estimates for 2015.¹⁴ The European Commission (EC) and the Organisation for Economic Co-operation and Development (OECD) also reported healthy life expectancy estimates for European countries from 2004 through 2014, but these were based on self-reported health status.^{18,19}

Here we present GBD 2015 findings for DALYs and HALE, building upon updated estimates of mortality, causes of death, and non-fatal health loss.^{12,20} Overall analytic approaches are similar to previous GBD studies,^{9,10} but include new mortality and morbidity data, refined methods, and expanded geographies.^{12,20} This report supersedes all previous GBD studies on DALYs and HALE through the estimation of a complete time-series for 1990 to 2015. To facilitate a more in-depth examination of the drivers of DALY and HALE trends, we assess how HALE, along with overall and cause-specific DALYs, change as geographies move through the development continuum. We use this analysis to benchmark overall progress and decompose observed disease burden compared with levels expected for specific causes on the basis of development alone, to highlight potential areas for policy investment or further research.

Methods

Overview

Detailed methods for estimating DALYs and HALE, including analytic approaches for mortality and non-fatal health loss estimation, are provided in related publications.^{12,20} Additional detail on GBD metrics and definitions are found elsewhere.²¹ Interactive tools are also available to explore GBD 2015 results and data sources. This analysis follows the Guidelines for Accurate and Transparent Health Estimates Reporting (GATHER), which includes recommendations on documentation of data sources, estimation methods, and statistical analysis.^{22,23}

In brief, the GBD geographic hierarchy involves 519 total geographies within 195 countries and territories, 21 regions, and seven super-regions. This study reports results for all countries and territories. The GBD cause hierarchy has four levels of classification and causes reported within each level that are mutually exclusive and collectively exhaustive. The full GBD cause list with corresponding International Classification of Diseases (ICD)-9 and ICD-10 codes are available in our publications on cause-specific mortality¹² and non-fatal health outcomes.²⁰

Estimation of mortality and non-fatal health loss

We estimated all-cause and cause-specific mortality with a multistep computation process, which included systematically addressing known data challenges, such as different coding schemes, different age-group reporting,

variation in certification, misclassification of HIV/AIDS deaths in some countries, misclassification of maternal HIV/AIDS deaths, and incorporation of population-based cancer registry data, before computation of cause-specific mortality with analytic tools such as Cause of Death Ensemble Modelling (CODEm). Each death could have only one underlying cause. Additional detail, including model specifications and data availability for each cause-specific model, can be found in the supplementary material of the GBD 2015 mortality and causes of death publication.¹² We calculated normative life tables based on the lowest death rates for each age group among geographies with total populations greater than 5 million. We computed cause-specific YLLs by multiplying cause-specific deaths by the life expectancy at the age of death (ie, 86·59 years at age 0 years; 23·79 years at age 65 years) from this normative life table, and then used the GBD world population age standard to calculate age-standardised mortality rates and YLL rates.¹²

Our most commonly used analytic approach to estimate non-fatal health loss was DisMod-MR 2.1, a Bayesian meta-regression tool that synthesises diverse data sources to produce internally consistent estimates of incidence, prevalence, remission, and excess mortality. The use of other methods to estimate non-fatal health loss was determined by cause-specific data availability and epidemiological characteristics.²⁴ Additional detail, including model specifications and data availability for each cause-specific model, can be found in the supplementary material of the GBD 2015 non-fatal publication.²⁰ Each non-fatal sequela was estimated separately. We then applied a microsimulation framework to assess the occurrence of comorbidity in each age group, sex, geography, and year separately. Disability from comorbid conditions was apportioned to each of the contributing causes. GBD disability weights were based on population surveys with more than 60 000 respondents, and previous studies show that disability weights do not significantly vary across geographies, income, or educational attainment.^{25,26} In this study, disability weights are invariant over geography and time, although the distribution of sequelae, and therefore the severity and cumulative disability per case of a condition, can differ by age, sex, geography, and year.

Estimation of DALYs, HALE, and corresponding uncertainty

DALYs are the sum of YLLs and YLDs as estimated in GBD 2015 for each cause, geography, age group, sex, and year.^{12,20} Using methods developed by Sullivan,⁷ we calculated HALE by age group within abridged multiple-decrement life tables and estimates of YLDs per capita for each geography–age–sex–year from 1990 to 2015.^{8,10,27}

For all results, we report 95% uncertainty intervals (UIs), which were derived from 1000 draws from the posterior distribution of each step in the estimation process. UIs are distinct from confidence intervals, because confidence

For the interactive visualisation of GBD 2015 results see
<http://vizhub.healthdata.org/gbd-compare/>

For the data citation tool see
<http://ghdx.healthdata.org/gbd-data-input-sources>

intervals only capture the uncertainty associated with sampling error, whereas uncertainty intervals provide a method for propagation of uncertainty from multiple sources including sampling, model estimation, and model specification. 95% UIs represent the ordinal 25th and 975th draw of the quantity of interest. For mortality and YLLs, UIs reflect uncertainty that arises from sample sizes of studies used as data sources, adjustments to sources of all-cause mortality, parameter uncertainty in model estimation, and specification uncertainty for all-cause and cause-specific models. For prevalence, incidence, and YLDs, UIs reflect the uncertainty that arises from sample sizes of studies used as data sources, data adjustments from non-reference definitions, parameter uncertainty in model estimation, and uncertainty in the disability weights. In the absence of any direction information about the correlation between uncertainty in YLLs and YLDs, we assumed uncertainty in age-specific YLDs is independent of age-specific YLLs in DALYs and death rates in HALE.

Epidemiological transition and relationship between DALYs, HALE, and SDI

We examined the relationship between DALYs, HALE, and the Socio-demographic Index (SDI).²⁸ SDI was constructed based on the geometric mean of three indicators: income per capita, average years of schooling among people aged 15 years or older, and the total fertility rate. SDI values were scaled to a range of 0 to 1, with 0 equalling the lowest income, lowest schooling, and highest fertility rate observed from 1980 to 2015, and 1 equalling the highest income, highest schooling, and lowest fertility rate assessed during that time. The average relationships between each summary health measure and SDI were estimated using spline regressions. These regressions were used to estimate expected values at each level of SDI. Additional detail on SDI computation and geography-specific SDI values are available in the appendix (pp 4–5 and pp 74–80).

See Online for appendix

Role of the funding source

The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Results

Global trends for DALYs and HALE

Worldwide total DALYs due to Group 1 causes fell from 1·2 billion (95% CI UI 1·2–1·2) in 1990, to 741·6 million (703·9–787·7) in 2015, whereas total DALYs due to NCDs increased from 1·1 billion (1·0–1·2), to 1·5 billion (1·3–1·7; figure 1; table 1). Total injury DALYs were relatively unchanged between 1990 and 2015. All-age DALY rates for NCDs changed little between 1990 and 2015, whereas they declined substantially for Group 1 causes. Taking into account population ageing, DALY rates for Group 1 causes

and NCDs both decreased (figure 1). For injuries, reductions in all-age DALY rates and age-standardised DALY rates were similar between 1990 and 2015.

HALE at birth increased to 60·9 years for men and 64·9 years for women in 2015, rising 2·9 years and 3·5 years since 2005, respectively (table 2). The gap between life expectancy and HALE, which represents years of functional health lost, widened between 2005 and 2015 from 7·7 years to 8·1 years for men, and from 9·4 years to 10·0 years for women. Life expectancy at age 65 years was 15·5 years for men and 18·5 years for women, while HALE was 11·9 years and 14·2 years for each sex, respectively (table 2).

Global causes of DALYs

In 2015, Group 1 causes accounted for 30·1% (95% UI 28·6–31·7%) of global DALYs, with NCDs leading to 59·7% (57·8–61·5) and injuries to 10·1% (9·5–10·7; table 1). Since 2005, DALYs due to many of the world's leading communicable causes substantially declined, yet burden increased for a subset of infectious diseases. Age-standardised DALY rates from HIV/AIDS and malaria each fell by more than 40% and lower respiratory infections and diarrhoeal diseases had decreases in total and age-standardised rates of DALYs of more than 20% (table 1). From 2005 to 2015, reductions in both total and age-standardised DALY rates due to tetanus and measles surpassed 50% and 70%, respectively. African trypanosomiasis, a disease targeted for elimination, saw both total DALYs and age-standardised rates fall by more than 70% since 2005. DALY rates substantially fell for all types of hepatitis, with age-standardised DALY rates from acute hepatitis A declining by more than 35% by 2015. However, both total DALYs and age-standardised DALY rates from dengue increased by more than 50%. Although the west African Ebola virus outbreak peaked in 2014,²⁹ Ebola virus disease still caused substantial DALYs in 2015 (table 1). Maternal disorders significantly declined from 2005 to 2015, with total DALYs and age-standardised rates each falling by more than 20%. Reductions in the global burden of neonatal disorders were somewhat less pronounced; for instance, the number of DALYs due to neonatal sepsis was largely unchanged (table 1).

In 2015, cardiovascular diseases, cancers, and mental and substance use disorders were among the leading causes of NCD burden (table 1). For many NCDs, including most cardiovascular causes and most cancers, total DALYs increased but age-standardised DALY rates declined. Nearly all neurological disorders increased in total DALYs, including Alzheimer's disease and other dementias, which rose by more than 30%, whereas age-standardised rates either moderately decreased (eg, Alzheimer's disease and other dementias) or were relatively unchanged (eg, Parkinson's disease). Total DALYs from low back and neck pain also increased, rising by more than 17%. Cirrhosis caused more DALYs in 2015 than in 2005, although age-standardised DALY rates significantly fell. A similar overall




Figure 1: Trends from 1990 to 2015, by GBD Level 1 cause, in global DALYs (A), crude DALY rates (B), and age-standardised DALY rates (C)
The difference in trends between (A) and (B) is caused by population growth and the difference between (B) and (C) is caused by changes in the percentage distribution of the population by age. Shaded areas show 95% uncertainty intervals. DALYs=disability-adjusted life-years.

trend was found for diabetes and chronic kidney disease, with all aetiologies apart from chronic kidney disease due to diabetes, showing significant declines in age-standardised DALY rates amid rising total DALYs. For other NCDs, namely those associated with skin diseases, sensory conditions, and oral disorders, total DALYs significantly increased from 2005 to 2015, and age-standardised DALY rates either somewhat increased or did not significantly change since 2005. Age-standardised DALY rates due to chronic obstructive pulmonary disease fell by more than 20% from 2005 to 2015, while those due to asthma decreased by almost 17%. Peptic ulcer disease, a leading cause of digestive disease burden, saw marked reductions in total DALYs and age-standardised DALY rates, with the latter decreasing by nearly 30% (table 1).

A number of NCDs significantly increased in terms of total burden and age-standardised DALY rates. Osteoarthritis was the most notable example, with total DALYs rising by 35% and age-standardised DALY rates by 4% between 1990 and 2015. Major depressive disorders and drug use disorders, particularly of opioids and cocaine, both increased in total DALYs and age-standardised rates; however, age-standardised DALY rates from alcohol use disorders dropped by 19%. Total DALYs and age-standardised DALY rates from chronic kidney disease due to diabetes also significantly increased by 2015. Male and female infertility accounted for a relatively small fraction of NCDs, but burden due to both causes increased significantly since 2005. Oral disorders and sense organ diseases also had increased total DALYs, whereas age-standardised DALY rates were relatively unchanged since 2005.

Unintentional injuries and transport injuries each saw age-standardised DALY rates significantly decrease (20% and 17%, respectively). Road injury burden significantly declined from 2005, with age-standardised DALY rates falling by 18% by 2015. Among unintentional injuries, drowning had the largest reduction in both total burden (26%) and age-standardised DALY rates (32%). Age-standardised DALY rates from self-harm and interpersonal violence both fell by more than 16% since 2005. DALYs due to forces of nature, war, and legal intervention increased from 2005 to 2015, although not significantly; this rise was primarily driven by escalated violence and war in the Middle East. Despite still causing major health loss in 2015, forces of nature caused far fewer DALYs than in 2005, mainly because there were no large-scale losses of life like that seen in the 2005 earthquake that killed more than 70 000 people in India and Pakistan.

Changes in leading causes of disease burden over time

In 1990, lower respiratory infections, preterm birth complications, and diarrhoeal diseases were the three leading causes of global DALYs; by 2015, only lower respiratory infections remained in the leading three (figure 2). Many Group 1 causes had significant declines for total burden, as well as all-age DALY rates and age-standardised DALY rates, for both time periods; these causes included tuberculosis, meningitis, diarrhoeal diseases, protein-energy malnutrition, preterm birth complications, tetanus, and measles. Such reductions across measures of DALYs contributed to downward shifts in relative ranks for most Group 1 causes over time. Malaria and HIV/AIDS both diverged from this trend,

	All-age DALYs (thousands)			Age-standardised rate (per 100 000)		
	2005	2015	Percentage change, 2005–15	2005	2015	Percentage change, 2005–15
All causes	2553306.8 (2373137.6 to 2756328.0)	2464895.4 (2259889.0 to 2696510.8)	-3.5 (-5.3 to -1.8)*	41561.7 (38640.8 to 44850.4)	34445.7 (31603.0 to 37654.3)	-17.1 (-18.7 to -15.7)*
Communicable, maternal, neonatal, and nutritional diseases	968014.5 (931809.4 to 1012327.9)	741595.9 (703928.7 to 787659.9)	-23.4 (-25.2 to -21.7)*	14297.6 (13766.0 to 14955.1)	10007.2 (9498.5 to 10629.3)	-30.0 (-31.6 to -28.5)*
HIV/AIDS and tuberculosis	148685.6 (141013.7 to 160635.2)	106991.8 (99680.8 to 116789.9)	-28.0 (-30.6 to -25.1)*	2296.0 (2172.7 to 2490.5)	1435.5 (1336.0 to 1569.2)	-37.5 (-39.6 to -35.0)*
Tuberculosis	49769.6 (43196.4 to 60348.2)	40302.2 (34065.8 to 49653.9)	-19.0 (-24.2 to -13.6)*	817.0 (708.5 to 990.9)	552.4 (467.2 to 681.3)	-32.4 (-36.7 to -27.9)*
HIV/AIDS	98916.1 (94515.6 to 103953.2)	66689.5 (63342.9 to 70788.3)	-32.6 (-35.2 to -29.5)*	1478.9 (1414.0 to 1553.2)	883.1 (839.2 to 937.1)	-40.3 (-42.6 to -37.5)*
HIV/AIDS—tuberculosis	19327.0 (15512.5 to 22015.1)	11621.9 (8955.5 to 13412.3)	-39.9 (-44.2 to -34.5)*	289.6 (232.8 to 329.8)	154.0 (118.7 to 177.7)	-46.8 (-50.7 to -42.1)*
HIV/AIDS resulting in other diseases	79589.0 (74869.9 to 85308.1)	55067.6 (51480.9 to 59294.3)	-30.8 (-34.2 to -26.9)*	1189.3 (1119.8 to 1277.1)	729.1 (681.9 to 784.8)	-38.7 (-41.7 to -35.3)*
Diarrhoea, lower respiratory, and other common infectious diseases	333534.0 (317202.5 to 351569.5)	242875.8 (230350.0 to 255919.1)	-27.2 (-30.2 to -24.2)*	5005.5 (4769.5 to 5269.7)	3326.2 (3158.6 to 3503.4)	-33.5 (-36.2 to -30.9)*
Diarrhoeal diseases	98394.2 (90909.6 to 106634.4)	71589.5 (66442.9 to 77205.8)	-27.2 (-33.2 to -20.8)*	1479.8 (1371.9 to 1598.4)	976.9 (908.1 to 1052.7)	-34.0 (-39.3 to -28.4)*
Intestinal infectious diseases	15062.4 (8707.5 to 24651.4)	12632.0 (7250.5 to 20762.5)	-16.1 (-22.9 to -9.9)*	216.0 (124.9 to 353.2)	170.8 (98.2 to 280.7)	-20.9 (-27.3 to -14.9)*
Typhoid fever	12543.9 (6953.1 to 21143.4)	10575.6 (5896.1 to 17598.0)	-15.7 (-22.7 to -9.3)*	179.8 (99.7 to 302.9)	143.0 (79.8 to 237.7)	-20.5 (-26.9 to -14.2)*
Paratyphoid fever	2393.2 (1119.8 to 4569.5)	2014.8 (960.5 to 3848.4)	-15.8 (-24.2 to -7.4)*	34.4 (16.1 to 65.6)	27.2 (13.0 to 52.0)	-20.8 (-28.7 to -12.6)*
Other intestinal infectious diseases	125.4 (39.3 to 243.9)	41.5 (16.3 to 81.3)	-66.9 (-78.0 to -47.0)*	1.8 (0.6 to 3.5)	0.6 (0.2 to 1.1)	-69.3 (-79.5 to -51.6)*
Lower respiratory infections	135293.2 (127083.7 to 143499.4)	103048.6 (96128.2 to 109078.8)	-23.8 (-28.2 to -19.4)*	2070.3 (1939.4 to 2189.6)	1428.5 (1330.2 to 1511.1)	-31.0 (-34.8 to -27.1)*
Upper respiratory infections	2661.2 (1550.4 to 4406.1)	2868.4 (1650.2 to 4785.2)	7.8 (5.2 to 9.7)*	40.1 (23.5 to 66.4)	38.7 (22.3 to 64.5)	-3.5 (-5.6 to -2.1)*
Otitis media	3444.4 (2276.8 to 5004.0)	3497.6 (2284.9 to 5088.5)	1.5 (-1.3 to 4.1)	50.9 (33.7 to 74.1)	47.3 (30.9 to 68.7)	-7.2 (-9.9 to -4.9)*
Meningitis	28394.6 (24170.2 to 32237.2)	25394.6 (21653.2 to 30649.0)	-10.6 (-20.1 to 3.8)	416.8 (355.2 to 471.9)	342.4 (292.3 to 412.8)	-17.8 (-26.4 to -4.6)*
Pneumococcal meningitis	7826.5 (6404.7 to 9672.3)	7773.4 (6353.4 to 9900.2)	-0.7 (-11.2 to 14.9)	115.3 (94.8 to 141.6)	104.9 (85.9 to 133.4)	-9.0 (-18.5 to 5.3)
Haemophilus influenzae type B meningitis	8507.6 (6656.5 to 10590.4)	5345.7 (4146.7 to 7103.2)	-37.2 (-45.8 to -25.0)*	123.3 (96.7 to 153.4)	71.9 (55.9 to 95.4)	-41.7 (-49.7 to -30.5)*
Meningococcal meningitis	5239.8 (4012.0 to 6704.3)	5065.6 (3879.5 to 6721.8)	-3.3 (-16.9 to 16.3)	76.6 (59.0 to 97.6)	68.2 (52.3 to 90.3)	-11.0 (-23.4 to 6.9)
Other meningitis	6820.7 (5665.6 to 8234.7)	7210.0 (5953.9 to 8958.4)	5.7 (-4.6 to 20.3)	101.7 (84.9 to 122.0)	97.4 (80.5 to 120.9)	-4.2 (-13.1 to 9.0)
Encephalitis	8850.1 (8064.3 to 9665.8)	8452.5 (7668.6 to 9411.6)	-4.5 (-11.6 to 2.9)	132.1 (120.8 to 144.2)	114.6 (104.0 to 127.6)	-13.2 (-19.7 to -6.8)*
Diphtheria	443.2 (229.3 to 905.0)	170.8 (83.4 to 388.7)	-61.5 (-86.1 to 3.5)	6.4 (3.3 to 13.1)	2.3 (1.1 to 5.2)	-64.2 (-87.1 to -3.7)*
Whooping cough	8559.1 (3269.5 to 19234.7)	5070.5 (1815.7 to 10791.8)	-40.8 (-77.3 to 62.2)	123.2 (47.0 to 276.7)	68.0 (24.3 to 144.7)	-44.8 (-78.8 to 51.2)
Tetanus	7213.2 (6251.3 to 9082.1)	3510.0 (3002.2 to 4502.9)	-51.3 (-58.0 to -43.8)*	105.1 (90.6 to 134.4)	47.1 (40.3 to 60.6)	-55.1 (-61.2 to -48.1)*
Measles	24602.4 (9272.8 to 51178.0)	6150.1 (2193.1 to 13467.4)	-75.0 (-84.4 to -59.0)*	355.1 (133.9 to 738.2)	82.7 (29.5 to 181.1)	-76.7 (-85.5 to -61.8)*
Varicella and herpes zoster	616.0 (521.8 to 727.1)	491.2 (386.1 to 625.3)	-20.3 (-33.2 to -4.3)*	9.7 (8.2 to 11.5)	6.8 (5.3 to 8.6)	-29.9 (-40.6 to -16.7)*

(Table 1 continues on next page)

	All-age DALYs (thousands)			Age-standardised rate (per 100 000)		
	2005	2015	Percentage change, 2005–15	2005	2015	Percentage change, 2005–15
(Continued from previous page)						
Neglected tropical diseases and malaria	116 583·6 (96 865·6 to 137 551·5)	79 212·2 (63 820·2 to 97 299·0)	-32·1 (-40·2 to -23·7)*	1710·8 (1422·4 to 2019·1)	1066·4 (859·1 to 1310·2)	-37·7 (-45·1 to -30·1)*
Malaria	90 438·1 (73 336·5 to 107 546·9)	55 769·6 (42 478·4 to 69 078·5)	-38·3 (-48·1 to -27·8)*	1315·8 (1067·0 to 1562·5)	749·3 (571·0 to 927·9)	-43·1 (-52·0 to -33·4)*
Chagas disease	243·6 (220·0 to 273·5)	236·1 (211·8 to 265·3)	-3·0 (-8·5 to 3·0)	4·4 (4·0 to 5·0)	3·4 (3·1 to 3·9)	-22·7 (-27·0 to -18·0)*
Leishmaniasis	1367·1 (878·4 to 1979·8)	1418·9 (1005·8 to 1913·8)	3·8 (-8·6 to 19·3)	20·1 (13·0 to 28·9)	18·9 (13·4 to 25·5)	-5·9 (-16·9 to 8·0)
Visceral leishmaniasis	1334·4 (844·9 to 1949·5)	1377·4 (965·4 to 1863·8)	3·2 (-9·2 to 19·1)	19·6 (12·4 to 28·5)	18·3 (12·9 to 24·8)	-6·3 (-17·4 to 7·9)
Cutaneous and mucocutaneous leishmaniasis	32·6 (15·2 to 63·3)	41·5 (19·4 to 81·0)	27·3 (23·6 to 30·6)*	0·5 (0·2 to 1·0)	0·6 (0·3 to 1·1)	11·4 (8·2 to 14·5)*
African trypanosomiasis	803·1 (425·6 to 1311·4)	202·4 (104·6 to 322·3)	-74·8 (-80·4 to -68·2)*	11·8 (6·3 to 19·3)	2·7 (1·4 to 4·3)	-77·2 (-82·3 to -71·3)*
Schistosomiasis	3417·7 (1883·7 to 6044·0)	2613·3 (1409·9 to 4695·8)	-23·5 (-30·6 to -6·7)*	51·1 (28·4 to 90·0)	35·1 (19·0 to 62·9)	-31·3 (-37·7 to -16·2)*
Cysticercosis	369·6 (260·6 to 489·8)	303·6 (211·4 to 410·9)	-17·9 (-22·7 to -13·5)*	5·9 (4·2 to 7·8)	4·1 (2·9 to 5·5)	-30·4 (-34·3 to -26·5)*
Cystic echinococcosis	174·1 (141·3 to 212·4)	172·6 (132·1 to 220·6)	-0·8 (-7·4 to 4·5)	2·8 (2·2 to 3·4)	2·4 (1·8 to 3·0)	-14·7 (-20·2 to -10·2)*
Lymphatic filariasis	2476·6 (1272·2 to 4008·8)	2075·0 (1120·5 to 3311·5)	-16·2 (-32·1 to -3·8)*	38·9 (20·0 to 63·1)	28·1 (15·2 to 44·9)	-27·7 (-41·5 to -17·0)*
Onchocerciasis	1441·6 (835·2 to 2288·7)	1135·7 (545·8 to 2005·7)	-21·2 (-38·6 to -4·8)*	22·5 (13·2 to 35·5)	15·5 (7·5 to 27·3)	-31·2 (-46·8 to -16·0)*
Trachoma	282·6 (194·3 to 396·1)	279·2 (192·5 to 396·2)	-1·2 (-5·6 to 3·0)	5·5 (3·8 to 7·7)	4·2 (2·9 to 5·9)	-23·9 (-27·6 to -20·3)*
Dengue	1132·8 (795·9 to 1616·2)	1892·2 (1266·7 to 2925·2)	67·0 (16·8 to 166·1)*	16·6 (11·7 to 23·8)	25·5 (17·1 to 39·5)	53·5 (7·3 to 144·1)*
Yellow fever	400·8 (83·3 to 1083·2)	329·8 (66·9 to 898·1)	-17·7 (-34·1 to 1·7)	5·8 (1·2 to 15·6)	4·4 (0·9 to 12·1)	-23·0 (-38·2 to -4·9)*
Rabies	1760·6 (1525·6 to 2019·8)	931·6 (779·2 to 1120·9)	-47·1 (-54·2 to -39·3)*	26·4 (23·0 to 30·2)	12·6 (10·6 to 15·2)	-52·1 (-58·6 to -45·2)*
Intestinal nematode infections	4404·5 (2719·6 to 6783·4)	3378·3 (2046·2 to 5294·5)	-23·3 (-27·4 to -18·2)*	65·8 (40·6 to 101·4)	45·6 (27·6 to 71·4)	-30·7 (-34·5 to -26·1)*
Ascariasis	1693·0 (1061·5 to 2610·7)	1075·4 (685·1 to 1660·9)	-36·5 (-43·1 to -29·5)*	25·2 (15·8 to 39·0)	14·5 (9·2 to 22·4)	-42·5 (-48·6 to -36·2)*
Trichuriasis	653·1 (357·9 to 1076·3)	544·1 (289·3 to 946·1)	-16·7 (-30·6 to 3·7)	9·8 (5·4 to 16·1)	7·3 (3·9 to 12·8)	-24·9 (-37·5 to -6·7)*
Hookworm disease	2058·4 (1275·6 to 3171·3)	1758·8 (1085·8 to 2755·1)	-14·6 (-19·9 to -9·4)*	30·8 (19·0 to 47·4)	23·7 (14·7 to 37·1)	-22·9 (-27·7 to -18·1)*
Food-borne trematodiases	1625·6 (793·1 to 3058·3)	1686·5 (855·0 to 3072·8)	3·7 (-0·5 to 10·1)	25·1 (12·3 to 47·0)	22·6 (11·5 to 41·0)	-10·0 (-13·5 to -5·2)*
Leprosy	30·8 (20·6 to 43·0)	31·0 (20·8 to 43·6)	0·6 (-1·5 to 2·9)	0·5 (0·4 to 0·8)	0·4 (0·3 to 0·6)	-19·2 (-20·8 to -17·4)*
Ebola virus disease	0·8 (0·7 to 1·0)	295·4 (238·3 to 353·8)	36 024·8 (36 010·3 to 36 054·6)*	0·0 (0·0 to 0·0)	3·9 (3·2 to 4·7)	31 654·0 (31 640·5 to 31 681·3)*
Other neglected tropical diseases	6213·6 (3655·0 to 11309·8)	6461·0 (3516·3 to 13769·5)	4·0 (-18·2 to 40·2)	91·8 (53·8 to 167·4)	87·6 (47·7 to 187·2)	-4·6 (-24·9 to 28·6)
Maternal disorders	20 797·1 (19 462·4 to 22 290·3)	16 282·0 (14 542·5 to 18 451·2)	-21·7 (-30·1 to -11·7)*	299·0 (279·8 to 320·7)	212·6 (189·9 to 240·9)	-28·9 (-36·5 to -19·8)*
Maternal haemorrhage	5614·4 (4902·4 to 6 402·8)	4645·8 (3769·9 to 5 636·8)	-17·3 (-28·9 to -4·6)*	81·1 (70·8 to 92·3)	60·7 (49·3 to 73·6)	-25·1 (-35·6 to -13·5)*
Maternal sepsis and other maternal infections	1443·1 (1195·4 to 1742·2)	1050·0 (792·7 to 1382·8)	-27·2 (-41·1 to -10·8)*	20·8 (17·3 to 25·1)	13·7 (10·4 to 18·0)	-34·1 (-46·8 to -19·0)*

(Table 1 continues on next page)

	All-age DALYs (thousands)			Age-standardised rate (per 100 000)		
	2005	2015	Percentage change, 2005–15	2005	2015	Percentage change, 2005–15
(Continued from previous page)						
Maternal hypertensive disorders	3939·7 (3391·5 to 4579·2)	2938·0 (2318·0 to 3698·5)	-25·4 (-35·9 to -13·2)*	56·1 (48·3 to 65·2)	38·3 (30·2 to 48·2)	-31·7 (-41·2 to -20·7)*
Maternal obstructed labour and uterine rupture	1905·3 (1592·5 to 2242·8)	1627·1 (1257·6 to 2025·1)	-14·6 (-25·4 to -2·4)*	27·7 (23·2 to 32·5)	21·2 (16·4 to 26·4)	-23·2 (-32·9 to -12·2)*
Maternal abortion, miscarriage and ectopic pregnancy	2366·6 (1992·0 to 2830·9)	1809·7 (1416·9 to 2283·6)	-23·5 (-34·2 to -12·1)*	34·0 (28·7 to 40·8)	23·6 (18·5 to 29·8)	-30·5 (-40·1 to -20·1)*
Indirect maternal deaths	2192·6 (1803·5 to 2621·7)	1744·2 (1299·7 to 2286·8)	-20·4 (-33·2 to -3·4)*	31·4 (25·9 to 37·5)	22·7 (17·0 to 29·8)	-27·6 (-39·0 to -11·9)*
Late maternal deaths	449·8 (295·7 to 652·2)	377·7 (244·8 to 562·2)	-16·0 (-26·8 to -2·6)*	6·5 (4·2 to 9·4)	4·9 (3·2 to 7·3)	-23·6 (-33·7 to -11·4)*
Maternal deaths aggravated by HIV/AIDS	149·9 (96·3 to 204·8)	122·1 (73·3 to 176·5)	-18·6 (-35·2 to 4·2)	2·2 (1·4 to 3·0)	1·6 (1·0 to 2·3)	-27·1 (-42·1 to -6·8)*
Other maternal disorders	2735·8 (2324·0 to 3245·0)	1967·5 (1578·3 to 2434·9)	-28·1 (-36·5 to -18·1)*	39·3 (33·4 to 46·6)	25·7 (20·6 to 31·7)	-34·7 (-42·4 to -26·1)*
Neonatal disorders	239 098·4 (232 872·7 to 246 048·9)	197 924·8 (191 388·9 to 204 751·8)	-17·2 (-19·2 to -15·4)*	3370·5 (3282·2 to 3468·1)	2638·3 (2551·0 to 2729·5)	-21·7 (-23·6 to -20·0)*
Neonatal preterm birth complications	99 050·0 (92 118·6 to 110 250·8)	74 833·6 (68 500·3 to 83 015·4)	-24·4 (-29·6 to -19·2)*	1396·6 (1298·8 to 1554·2)	997·5 (913·1 to 1106·5)	-28·6 (-33·4 to -23·6)*
Neonatal encephalopathy due to birth asphyxia and trauma	79 423·6 (72 215·4 to 87 185·6)	67 856·5 (61 497·7 to 75 824·7)	-14·6 (-21·7 to -6·7)*	1118·8 (1017·3 to 1228·6)	904·4 (819·6 to 1010·5)	-19·2 (-26·0 to -11·7)*
Neonatal sepsis and other neonatal infections	30 510·1 (21 830·7 to 40 329·6)	30 454·5 (21 591·8 to 39 747·9)	-0·2 (-16·2 to 20·3)	429·6 (307·4 to 567·7)	406·0 (287·8 to 529·8)	-5·5 (-20·6 to 13·9)
Haemolytic disease and other neonatal jaundice	6427·0 (4234·6 to 9708·3)	4504·0 (3202·7 to 6420·6)	-29·9 (-44·1 to -13·6)*	90·9 (60·0 to 137·0)	60·1 (42·7 to 85·6)	-33·9 (-47·2 to -18·6)*
Other neonatal disorders	23 687·6 (17 420·2 to 30 795·3)	20 276·2 (15 531·0 to 25 385·9)	-14·4 (-32·8 to 7·1)	334·6 (246·3 to 434·7)	270·4 (207·1 to 338·5)	-19·2 (-36·6 to 1·1)
Nutritional deficiencies	84 133·3 (65 215·4 to 108 480·6)	76 517·1 (58 781·9 to 100 939·6)	-9·1 (-15·2 to -3·5)*	1242·7 (964·9 to 1598·6)	1034·4 (795·9 to 1363·6)	-16·8 (-22·4 to -11·7)*
Protein-energy malnutrition	26 655·2 (21 449·8 to 32 723·9)	21 094·0 (16 844·2 to 26 299·5)	-20·9 (-34·5 to -4·8)*	395·1 (319·4 to 484·0)	286·5 (228·9 to 356·6)	-27·5 (-39·8 to -13·0)*
Iodine deficiency	2320·5 (1500·7 to 3422·1)	2476·4 (1592·8 to 3662·1)	6·7 (3·4 to 9·9)*	35·3 (22·8 to 52·1)	33·4 (21·5 to 49·4)	-5·2 (-8·4 to -2·4)*
Vitamin A deficiency	209·5 (130·8 to 309·2)	232·4 (143·4 to 346·3)	10·9 (7·5 to 14·6)*	3·1 (2·0 to 4·6)	3·1 (1·9 to 4·7)	0·2 (-2·7 to 3·3)
Iron-deficiency anaemia	52 951·3 (36 342·0 to 74 873·5)	51 217·1 (35 014·4 to 72 661·0)	-3·3 (-4·8 to -1·8)*	778·6 (534·8 to 1099·4)	690·8 (472·4 to 979·6)	-11·3 (-12·7 to -10·0)*
Other nutritional deficiencies	1996·8 (1378·4 to 2848·4)	1497·2 (1135·7 to 1983·7)	-25·0 (-43·3 to -9·4)*	30·5 (21·2 to 43·5)	20·5 (15·7 to 27·1)	-32·7 (-48·9 to -19·0)*
Other communicable, maternal, neonatal, and nutritional diseases	25 182·6 (20 002·2 to 31 397·3)	21 792·2 (17 474·0 to 26 922·3)	-13·5 (-18·6 to -7·8)*	373·1 (298·4 to 462·5)	293·8 (235·7 to 362·6)	-21·2 (-25·9 to -16·1)*
Sexually transmitted diseases excluding HIV	12 378·7 (7 831·6 to 18 396·6)	10 330·9 (6 583·0 to 15 315·1)	-16·5 (-24·2 to -8·6)*	178·7 (113·6 to 264·7)	138·1 (87·9 to 205·0)	-22·7 (-29·6 to -15·5)*
Syphilis	11 190·5 (6 607·1 to 17 280·8)	8957·1 (5273·2 to 13969·7)	-20·0 (-28·4 to -11·8)*	160·9 (95·4 to 247·9)	120·1 (70·8 to 187·2)	-25·4 (-33·3 to -17·7)*
Chlamydial infection	337·0 (194·2 to 537·7)	369·8 (214·0 to 595·4)	9·7 (6·5 to 13·0)*	4·9 (2·8 to 7·8)	4·8 (2·8 to 7·7)	-2·3 (-5·0 to 0·5)
Gonococcal infection	382·8 (238·6 to 573·0)	469·8 (283·4 to 716·8)	22·7 (15·9 to 28·1)*	5·6 (3·5 to 8·4)	6·1 (3·7 to 9·3)	8·8 (2·8 to 13·8)*
Trichomoniasis	167·4 (67·1 to 354·8)	194·3 (77·8 to 412·1)	16·1 (15·0 to 17·2)*	2·5 (1·0 to 5·4)	2·6 (1·0 to 5·4)	1·0 (0·3 to 1·8)*
Genital herpes	197·8 (61·5 to 468·3)	236·4 (74·3 to 555·8)	19·5 (17·4 to 23·0)*	3·2 (1·0 to 7·6)	3·2 (1·0 to 7·5)	0·7 (-1·5 to 5·2)
Other sexually transmitted diseases	103·2 (74·9 to 139·2)	103·5 (74·3 to 141·2)	0·3 (-3·1 to 3·2)	1·6 (1·1 to 2·1)	1·4 (1·0 to 1·9)	-13·2 (-16·1 to -10·6)*

(Table 1 continues on next page)

	All-age DALYs (thousands)			Age-standardised rate (per 100 000)		
	2005	2015	Percentage change, 2005–15	2005	2015	Percentage change, 2005–15
(Continued from previous page)						
Hepatitis	6042·4 (5738·7 to 6344·8)	4925·3 (4634·8 to 5229·3)	-18·5 (-22·7 to -14·1)*	93·0 (88·6 to 97·5)	66·7 (62·8 to 70·8)	-28·3 (-31·9 to -24·6)*
Acute hepatitis A	1486·6 (1058·5 to 1896·9)	1030·3 (714·9 to 1335·9)	-30·7 (-39·7 to -20·8)*	21·5 (15·2 to 27·5)	13·8 (9·6 to 17·9)	-35·6 (-43·9 to -26·4)*
Hepatitis B	2693·5 (2298·5 to 3080·0)	2449·1 (2083·5 to 2799·2)	-9·1 (-15·4 to -2·0)*	44·0 (37·8 to 50·0)	33·4 (28·5 to 38·1)	-24·1 (-29·2 to -18·6)*
Hepatitis C	98·4 (26·2 to 220·4)	89·1 (25·5 to 201·9)	-9·5 (-23·3 to 7·6)	1·6 (0·4 to 3·7)	1·2 (0·3 to 2·8)	-25·9 (-37·5 to -11·0)*
Acute hepatitis E	1763·9 (1304·1 to 2265·3)	1356·8 (972·0 to 1794·4)	-23·1 (-31·6 to -14·0)*	25·9 (19·0 to 33·7)	18·3 (13·1 to 24·2)	-29·6 (-37·1 to -21·6)*
Other infectious diseases	6761·5 (4784·8 to 8106·7)	6536·0 (4527·7 to 7935·6)	-3·3 (-13·9 to 9·2)	101·3 (72·0 to 120·9)	89·0 (61·6 to 107·8)	-12·2 (-21·6 to -10·0)*
Non-communicable diseases	1322207·9 (1181179·7 to 1471448·8)	1473508·2 (1309803·2 to 1650147·4)	11·4 (9·7 to 13·0)*	23220·2 (20898·5 to 25689·0)	21062·4 (18790·6 to 23518·0)	-9·3 (-10·9 to -7·9)*
Neoplasms	187562·1 (183699·7 to 191539·2)	209359·2 (204155·5 to 214470·1)	11·6 (9·3 to 14·2)*	3424·3 (3356·2 to 3494·4)	3024·4 (2949·8 to 3098·9)	-11·7 (-13·4 to -9·7)*
Lip and oral cavity cancer	2954·1 (2866·9 to 3050·0)	3780·1 (3645·7 to 3928·9)	28·0 (23·9 to 32·6)*	53·7 (52·1 to 55·4)	53·7 (51·8 to 55·8)	0·1 (-3·1 to 3·7)
Nasopharynx cancer	1805·8 (1477·1 to 1912·4)	1911·7 (1562·1 to 2040·2)	5·9 (-2·2 to 12·4)	30·9 (25·4 to 32·7)	26·5 (21·6 to 28·2)	-14·3 (-20·6 to -9·0)*
Other pharynx cancer	1419·9 (1374·7 to 1468·8)	1715·7 (1635·4 to 1795·0)	20·8 (15·2 to 26·3)*	25·9 (25·1 to 26·8)	24·3 (23·2 to 25·4)	-6·3 (-10·7 to -2·1)*
Oesophageal cancer	10665·4 (10313·5 to 11043·9)	9854·4 (9465·3 to 10270·8)	-7·6 (-12·4 to -2·1)*	201·0 (194·5 to 207·8)	143·6 (137·9 to 149·5)	-28·6 (-32·2 to -24·4)*
Stomach cancer	18665·9 (18186·3 to 19175·5)	17439·6 (16876·7 to 18034·6)	-6·6 (-9·7 to -3·3)*	350·6 (341·9 to 360·0)	255·9 (247·9 to 264·5)	-27·0 (-29·4 to -24·5)*
Colon and rectum cancer	14409·5 (14108·0 to 14779·0)	17026·6 (16586·7 to 17504·7)	18·2 (15·7 to 20·9)*	274·0 (268·2 to 280·8)	251·3 (244·9 to 258·4)	-8·3 (-10·2 to -6·2)*
Liver cancer	19643·6 (16871·4 to 20641·2)	20578·0 (18937·8 to 21915·3)	4·8 (-1·4 to 15·5)	350·9 (302·8 to 368·5)	292·1 (269·1 to 311·0)	-16·8 (-21·4 to -8·7)*
Liver cancer due to hepatitis B	8420·5 (7092·2 to 9020·2)	8029·5 (7279·6 to 8795·5)	-4·6 (-11·2 to 8·2)	143·4 (121·3 to 153·7)	110·6 (100·4 to 121·2)	-22·9 (-27·9 to -12·9)*
Liver cancer due to hepatitis C	2897·4 (2607·7 to 3109·1)	3324·0 (3012·2 to 3573·6)	14·7 (10·3 to 20·2)*	55·5 (50·1 to 59·5)	49·3 (44·9 to 52·9)	-11·2 (-14·4 to -7·1)*
Liver cancer due to alcohol use	4786·9 (4075·0 to 5168·7)	5888·5 (5368·4 to 6441·0)	23·0 (14·8 to 36·1)*	89·2 (76·2 to 96·3)	84·8 (77·5 to 92·7)	-4·9 (-11·1 to 4·8)
Liver cancer due to other causes	3538·8 (3022·8 to 3829·6)	3336·0 (3019·1 to 3637·7)	-5·7 (-11·2 to 3·8)	62·9 (54·1 to 68·1)	47·4 (42·9 to 51·6)	-24·6 (-28·9 to -17·5)*
Gallbladder and biliary tract cancer	2448·7 (2359·2 to 2524·3)	2615·7 (2454·6 to 2740·8)	6·8 (2·2 to 11·6)*	47·3 (45·7 to 48·7)	39·0 (36·7 to 40·9)	-17·5 (-21·0 to -13·8)*
Pancreatic cancer	6528·1 (6433·9 to 6630·7)	8236·6 (8064·7 to 8420·0)	26·2 (23·3 to 29·0)*	125·2 (123·4 to 127·0)	121·8 (119·3 to 124·4)	-2·7 (-4·8 to -0·6)*
Larynx cancer	2364·1 (2290·8 to 2443·5)	2608·5 (2518·3 to 2705·0)	10·3 (7·1 to 13·9)*	44·0 (42·7 to 45·5)	37·6 (36·3 to 39·0)	-14·5 (-17·0 to -11·8)*
Tracheal, bronchus, and lung cancer	31802·0 (31117·2 to 32516·2)	36419·5 (35356·6 to 37615·6)	14·5 (11·0 to 19·1)*	604·5 (591·8 to 617·4)	535·9 (520·3 to 553·1)	-11·3 (-14·0 to -7·9)*
Malignant skin melanoma	1304·0 (1086·8 to 1655·2)	1596·3 (1293·4 to 1982·7)	22·4 (15·6 to 27·5)*	23·3 (19·4 to 29·6)	22·8 (18·4 to 28·3)	-2·3 (-7·7 to 1·7)
Non-melanoma skin cancer	794·3 (764·9 to 828·0)	1088·8 (1028·3 to 1150·1)	37·1 (32·0 to 41·9)*	15·2 (14·7 to 15·9)	16·3 (15·4 to 17·2)	6·8 (2·8 to 10·6)*
Non-melanoma skin cancer (squamous-cell carcinoma)	789·6 (761·2 to 822·3)	1082·9 (1025·0 to 1142·0)	37·1 (32·0 to 42·0)*	15·1 (14·6 to 15·8)	16·2 (15·3 to 17·1)	6·9 (2·9 to 10·6)*
Non-melanoma skin cancer (basal-cell carcinoma)	4·7 (2·2 to 8·7)	6·0 (2·8 to 11·1)	26·9 (23·4 to 30·2)*	0·1 (0·0 to 0·2)	0·1 (0·0 to 0·2)	-3·8 (-6·5 to -1·2)*

(Table 1 continues on next page)

	All-age DALYs (thousands)			Age-standardised rate (per 100 000)		
	2005	2015	Percentage change, 2005-15	2005	2015	Percentage change, 2005-15
(Continued from previous page)						
Breast cancer	12 939·2 (12 181·7 to 13 828·7)	15 410·6 (14 415·5 to 16 217·8)	19·1 (11·6 to 25·8)*	231·4 (218·3 to 246·9)	217·1 (203·2 to 228·7)	-6·2 (-11·9 to -1·1)*
Cervical cancer	6819·2 (6393·4 to 7236·4)	6963·0 (6526·2 to 7408·0)	2·1 (-4·5 to 10·5)	118·9 (111·8 to 126·1)	96·7 (90·7 to 102·7)	-18·7 (-23·9 to -12·2)*
Uterine cancer	2077·7 (1963·6 to 2201·8)	2230·7 (2098·3 to 2386·9)	7·4 (0·3 to 15·8)*	38·7 (36·6 to 40·9)	32·2 (30·3 to 34·5)	-16·6 (-22·0 to -10·1)*
Ovarian cancer	3496·3 (3401·0 to 3660·5)	4135·9 (3992·4 to 4298·0)	18·3 (13·4 to 23·1)*	63·2 (61·5 to 66·0)	58·6 (56·6 to 60·8)	-7·3 (-11·0 to -3·8)*
Prostate cancer	4793·4 (3992·1 to 6079·2)	6281·2 (5233·9 to 7904·2)	31·0 (26·9 to 35·2)*	100·0 (83·5 to 126·8)	99·6 (83·0 to 125·7)	-0·4 (-3·5 to 2·6)
Testicular cancer	411·3 (390·2 to 435·8)	442·6 (413·1 to 473·3)	7·6 (0·3 to 13·6)*	6·3 (6·0 to 6·7)	5·9 (5·5 to 6·3)	-6·4 (-12·6 to -1·2)*
Kidney cancer	2649·3 (2539·2 to 2779·3)	3340·3 (3189·6 to 3513·2)	26·1 (21·2 to 30·6)*	48·5 (46·8 to 50·6)	48·4 (46·2 to 50·8)	-0·3 (-3·8 to 3·0)
Bladder cancer	2832·7 (2754·6 to 2908·1)	3368·8 (3251·3 to 3496·5)	18·9 (15·4 to 22·4)*	56·3 (54·8 to 57·8)	51·3 (49·5 to 53·2)	-8·8 (-11·5 to -6·2)*
Brain and nervous system cancer	6738·3 (6079·6 to 7270·6)	7624·4 (6975·3 to 8218·8)	13·2 (5·0 to 21·0)*	111·4 (100·8 to 119·7)	105·6 (96·7 to 113·7)	-5·2 (-11·7 to 1·2)
Thyroid cancer	649·6 (605·3 to 707·6)	846·3 (754·1 to 929·0)	30·3 (19·2 to 37·4)*	12·1 (11·3 to 13·2)	12·4 (11·1 to 13·6)	2·2 (-6·2 to 7·8)
Mesothelioma	544·9 (529·2 to 560·6)	701·9 (678·8 to 722·2)	28·8 (24·3 to 33·3)*	10·1 (9·8 to 10·3)	10·3 (9·9 to 10·5)	2·0 (-1·5 to 5·4)
Hodgkin's lymphoma	949·3 (837·3 to 1120·9)	845·6 (758·4 to 1031·7)	-10·9 (-14·9 to -6·7)*	15·5 (13·7 to 18·3)	11·6 (10·5 to 14·2)	-24·7 (-28·2 to -21·1)*
Non-Hodgkin lymphoma	5063·1 (4581·8 to 5581·3)	6283·8 (5449·8 to 6632·3)	24·1 (11·5 to 31·7)*	88·3 (79·7 to 96·6)	89·5 (77·5 to 94·3)	1·3 (-8·5 to 7·0)
Multiple myeloma	1694·3 (1646·6 to 1751·4)	2182·4 (2096·5 to 2255·8)	28·8 (23·8 to 33·2)*	32·2 (31·2 to 33·2)	32·1 (30·8 to 33·1)	-0·4 (-4·1 to 2·9)
Leukaemia	11 272·2 (10 956·3 to 11 683·7)	12 040·4 (11 609·8 to 12 499·3)	6·8 (2·9 to 10·5)*	181·6 (176·6 to 187·8)	167·7 (161·8 to 173·9)	-7·6 (-10·8 to -4·6)*
Acute lymphoid leukaemia	4857·0 (4505·3 to 5452·8)	5057·8 (4658·4 to 5507·7)	4·1 (-1·7 to 9·9)	73·6 (68·3 to 82·4)	69·1 (63·6 to 75·2)	-6·1 (-11·3 to -1·0)*
Chronic lymphoid leukaemia	1183·2 (1107·9 to 1256·6)	1268·6 (1194·2 to 1349·3)	7·2 (1·8 to 12·8)*	21·8 (20·5 to 23·1)	18·7 (17·7 to 19·9)	-14·1 (-18·3 to -10·0)*
Acute myeloid leukaemia	4199·7 (3770·1 to 4553·5)	4769·0 (4345·7 to 5193·7)	13·6 (8·2 to 18·5)*	68·5 (62·1 to 74·0)	66·6 (60·8 to 72·5)	-2·8 (-7·0 to 1·1)
Chronic myeloid leukaemia	1032·3 (957·8 to 1131·8)	945·1 (885·8 to 1032·9)	-8·5 (-12·3 to -3·8)*	17·7 (16·5 to 19·3)	13·3 (12·5 to 14·5)	-24·8 (-27·8 to -21·2)*
Other neoplasms	9825·9 (9023·1 to 10 351·7)	11 789·9 (10 528·0 to 12 616·2)	20·0 (13·4 to 25·7)*	163·3 (150·1 to 171·6)	164·6 (146·8 to 175·8)	0·8 (-4·5 to 5·5)
Cardiovascular diseases	326 252·4 (318 365·8 to 334 285·8)	347 528·9 (337 220·3 to 358 093·5)	6·5 (4·1 to 8·7)*	6231·9 (6086·8 to 6378·1)	5179·7 (5026·3 to 5334·2)	-16·9 (-18·7 to -15·3)*
Rheumatic heart disease	11 594·7 (10 668·9 to 12 707·4)	10 513·2 (9 611·0 to 11 154·5)	-9·3 (-13·3 to -5·7)*	193·6 (178·4 to 210·6)	146·6 (134·2 to 160·1)	-24·3 (-27·5 to -21·3)*
Ischaemic heart disease	147 780·0 (144 845·9 to 151 651·5)	164 020·4 (159 621·3 to 169 088·2)	11·0 (8·3 to 13·3)*	2860·0 (2805·1 to 2927·3)	2452·6 (2388·9 to 2526·8)	-14·2 (-16·2 to -12·5)*
Cerebrovascular disease	118 566·2 (115 776·1 to 121 419·2)	118 626·7 (114 862·4 to 122 627·0)	0·1 (-2·5 to 2·7)	2283·2 (2230·3 to 2335·8)	1776·6 (1721·1 to 1834·8)	-22·2 (-24·1 to -20·1)*
Ischaemic stroke	44 104·6 (42 363·3 to 45 822·3)	45 208·5 (43 150·2 to 47 386·8)	2·5 (-0·4 to 5·2)	903·6 (869·6 to 936·9)	706·2 (675·6 to 739·0)	-21·8 (-24·0 to -19·8)*
Haemorrhagic stroke	74 461·5 (72 176·9 to 76 874·3)	73 418·2 (70 737·2 to 76 596·4)	-1·4 (-4·6 to 2·1)	1379·6 (1337·6 to 1423·6)	1070·4 (1032·1 to 1116·3)	-22·4 (-25·0 to -19·6)*
Hypertensive heart disease	14 852·4 (13 919·4 to 16 052·6)	17 484·6 (16 286·9 to 18 593·6)	17·7 (11·6 to 22·9)*	286·5 (268·8 to 309·8)	262·2 (243·2 to 279·1)	-8·5 (-13·4 to -4·5)*

(Table 1 continues on next page)

	All-age DALYs (thousands)			Age-standardised rate (per 100 000)		
	2005	2015	Percentage change, 2005–15	2005	2015	Percentage change, 2005–15
(Continued from previous page)						
Cardiomyopathy and myocarditis	9619·8 (9134·6 to 9993·7)	9220·3 (8765·6 to 9706·0)	-4·2 (-8·3 to -0·1)*	163·4 (155·8 to 169·5)	130·0 (123·7 to 136·7)	-20·5 (-23·7 to -17·2)*
Atrial fibrillation and flutter	3463·2 (2764·5 to 4281·3)	4433·7 (3541·7 to 5489·1)	28·0 (26·6 to 29·3)*	73·8 (59·3 to 90·5)	70·6 (56·5 to 86·9)	-4·2 (-5·3 to -3·3)*
Aortic aneurysm	2468·0 (2403·2 to 2547·3)	2930·2 (2846·0 to 3007·0)	18·7 (12·9 to 22·6)*	47·6 (46·4 to 49·1)	43·8 (42·6 to 45·0)	-7·9 (-12·3 to -5·0)*
Peripheral vascular disease	913·0 (686·4 to 1279·3)	1198·8 (898·0 to 1682·6)	31·3 (27·7 to 35·2)*	19·3 (14·5 to 26·9)	19·1 (14·3 to 26·8)	-1·1 (-3·9 to 1·7)
Endocarditis	1962·3 (1616·0 to 2190·5)	2210·3 (1838·9 to 2421·3)	12·6 (6·3 to 18·8)*	33·0 (27·7 to 36·7)	31·2 (26·1 to 34·2)	-5·5 (-10·2 to -0·9)*
Other cardiovascular and circulatory diseases	15 032·7 (13 427·8 to 16 820·7)	16 890·7 (14 824·7 to 19 104·2)	12·4 (8·4 to 15·6)*	271·6 (243·4 to 303·7)	247·0 (217·2 to 279·2)	-9·0 (-12·1 to -6·5)*
Chronic respiratory diseases	97 356·3 (90 607·8 to 104 559·9)	97 451·8 (89 829·7 to 105 527·6)	0·1 (-2·4 to 2·6)	1802·0 (1694·4 to 1915·3)	1443·4 (1336·9 to 1556·8)	-19·9 (-22·2 to -17·8)*
Chronic obstructive pulmonary disease	63 773·8 (60 981·4 to 66 186·6)	63 850·4 (61 215·3 to 66 288·6)	0·1 (-3·1 to 3·8)	1246·9 (1195·2 to 1291·9)	971·1 (932·7 to 1007·6)	-22·1 (-24·6 to -19·2)*
Pneumoconiosis	976·7 (816·2 to 1168·6)	1099·6 (902·7 to 1314·7)	12·6 (6·9 to 18·9)*	17·8 (15·0 to 21·1)	16·0 (13·2 to 19·1)	-9·9 (-14·8 to -4·7)*
Silicosis	271·5 (240·1 to 308·1)	270·2 (235·8 to 310·1)	-0·5 (-10·1 to 11·5)	5·0 (4·5 to 5·7)	4·0 (3·5 to 4·5)	-21·3 (-28·7 to -12·0)*
Asbestosis	76·3 (63·0 to 87·9)	92·0 (75·3 to 107·2)	20·5 (13·7 to 27·8)*	1·4 (1·2 to 1·6)	1·4 (1·1 to 1·6)	-3·8 (-8·9 to 1·7)
Coal worker's pneumoconiosis	59·7 (52·8 to 67·4)	57·5 (49·0 to 67·2)	-3·6 (-17·4 to 9·4)	1·2 (1·0 to 1·3)	0·9 (0·7 to 1·0)	-25·1 (-35·6 to -15·1)*
Other pneumoconiosis	569·3 (440·2 to 715·7)	680·0 (526·2 to 855·9)	19·5 (13·4 to 25·9)*	10·2 (8·0 to 12·7)	9·8 (7·6 to 12·3)	-3·4 (-9·0 to 2·3)
Asthma	26 859·7 (21 268·1 to 33 122·5)	26 168·8 (20 501·4 to 32 583·0)	-2·6 (-9·2 to 4·2)	439·8 (351·4 to 537·5)	365·6 (287·5 to 453·5)	-16·9 (-23·5 to -10·3)*
Interstitial lung disease and pulmonary sarcoidosis	1672·3 (1351·2 to 1954·1)	2352·6 (1933·8 to 2618·7)	40·7 (29·2 to 48·8)*	32·0 (25·7 to 37·2)	35·4 (28·9 to 39·3)	10·3 (1·5 to 16·7)*
Other chronic respiratory diseases	4073·7 (3295·2 to 4805·9)	3980·4 (3225·6 to 4569·7)	-2·3 (-11·0 to 7·3)	65·5 (53·3 to 76·6)	55·4 (44·9 to 63·5)	-15·3 (-22·4 to -7·5)*
Cirrhosis and other chronic liver diseases	37 101·5 (35 722·1 to 39 385·8)	38 973·3 (37 202·0 to 41 702·9)	5·0 (1·6 to 8·7)*	632·5 (609·4 to 669·2)	539·6 (515·4 to 576·8)	-14·7 (-17·4 to -11·8)*
Cirrhosis and other chronic liver diseases due to hepatitis B	10 297·6 (9 495·9 to 11 233·5)	10 754·7 (9 876·8 to 11 957·6)	4·4 (0·2 to 8·6)*	178·0 (164·1 to 194·8)	149·5 (137·3 to 165·9)	-16·0 (-19·2 to -12·7)*
Cirrhosis and other chronic liver diseases due to hepatitis C	8408·4 (7768·6 to 9069·5)	9161·7 (8443·3 to 9968·2)	9·0 (5·7 to 12·3)*	146·5 (135·6 to 157·9)	127·4 (117·6 to 138·4)	-13·0 (-15·6 to -10·4)*
Cirrhosis and other chronic liver diseases due to alcohol use	10 093·5 (9 424·0 to 10 841·4)	10 997·4 (10 197·4 to 11 875·1)	9·0 (5·0 to 13·7)*	171·9 (160·5 to 184·3)	150·7 (139·7 to 162·6)	-12·3 (-15·5 to -8·6)*
Cirrhosis and other chronic liver diseases due to other causes	8302·1 (7741·0 to 9165·9)	8059·5 (7452·2 to 8977·7)	-2·9 (-6·5 to 1·3)	136·1 (126·8 to 149·9)	112·0 (103·6 to 124·7)	-17·7 (-20·6 to -14·3)*
Digestive diseases	41 351·8 (37 701·8 to 45 778·3)	42 189·1 (38 101·1 to 47 087·5)	2·0 (-2·7 to 6·2)	716·3 (653·0 to 793·4)	600·3 (542·1 to 670·1)	-16·2 (-19·7 to -12·9)*
Peptic ulcer disease	10 164·0 (9 220·6 to 11 279·2)	8894·4 (7950·0 to 10 039·2)	-12·5 (-17·7 to -5·9)*	180·8 (163·9 to 201·3)	128·5 (114·6 to 145·3)	-28·9 (-33·2 to -23·6)*
Gastritis and duodenitis	5757·0 (4331·9 to 7638·9)	6145·5 (4517·3 to 8251·9)	6·7 (2·5 to 11·1)*	100·9 (76·0 to 134·0)	87·8 (64·6 to 118·0)	-13·1 (-16·4 to -9·7)*
Appendicitis	2197·3 (1823·7 to 2661·6)	2015·9 (1631·6 to 2381·2)	-8·3 (-21·5 to 6·4)	34·4 (28·7 to 41·2)	27·6 (22·4 to 32·6)	-19·6 (-30·7 to -6·8)*
Paralytic ileus and intestinal obstruction	6957·0 (6123·4 to 8236·1)	7135·5 (6464·3 to 8502·6)	2·6 (-5·5 to 11·2)	117·3 (104·1 to 138·5)	100·7 (91·4 to 119·7)	-14·2 (-20·1 to -7·5)*
Inguinal femoral and abdominal hernia	1689·6 (1147·1 to 1996·2)	1645·7 (1089·7 to 1969·1)	-2·6 (-9·8 to 7·0)	29·2 (20·1 to 34·2)	23·4 (15·6 to 27·9)	-19·7 (-26·1 to -11·8)*

(Table 1 continues on next page)

	All-age DALYs (thousands)			Age-standardised rate (per 100 000)		
	2005	2015	Percentage change, 2005–15	2005	2015	Percentage change, 2005–15
(Continued from previous page)						
Inflammatory bowel disease	3222.5 (2514.5 to 4041.9)	3588.0 (2797.3 to 4483.0)	11.3 (5.1 to 15.4)*	53.8 (42.3 to 67.1)	49.8 (39.0 to 62.0)	-7.5 (-12.2 to -4.3)*
Vascular intestinal disorders	1475.6 (1358.6 to 1619.3)	1741.6 (1607.3 to 1903.4)	18.0 (12.8 to 23.7)*	28.8 (26.6 to 31.5)	26.5 (24.5 to 28.8)	-8.2 (-12.2 to -3.9)*
Gallbladder and biliary diseases	2706.6 (2459.3 to 2994.3)	2878.7 (2597.4 to 3161.8)	6.4 (-0.2 to 11.9)	48.3 (44.1 to 53.0)	41.6 (37.6 to 45.6)	-13.8 (-18.7 to -9.6)*
Pancreatitis	3765.2 (3501.1 to 4071.3)	4373.2 (4000.1 to 4766.0)	16.1 (9.2 to 23.2)*	62.8 (58.5 to 67.8)	60.2 (55.1 to 65.5)	-4.1 (-9.6 to 1.6)
Other digestive diseases	3417.0 (2883.7 to 4228.0)	3770.5 (3180.0 to 4438.8)	10.3 (-0.1 to 22.2)	60.0 (51.0 to 73.9)	54.2 (45.9 to 63.7)	-9.7 (-18.0 to -0.4)*
Neurological disorders	74 456.9 (58 081.0 to 93 693.7)	87 082.1 (67 962.4 to 109 039.4)	17.0 (15.5 to 18.5)*	1291.0 (1025.5 to 1596.9)	1252.4 (987.9 to 1556.6)	-3.0 (-4.2 to -2.0)*
Alzheimer's disease and other dementias	17 905.9 (15 147.4 to 20 849.2)	23 779.2 (20 118.0 to 27 886.0)	32.8 (31.2 to 34.4)*	409.6 (348.0 to 479.2)	395.6 (333.5 to 463.8)	-3.4 (-4.4 to -2.5)*
Parkinson's disease	1548.9 (1381.7 to 1747.5)	2059.3 (1831.8 to 2320.9)	32.9 (29.7 to 35.7)*	32.8 (29.4 to 36.8)	33.1 (29.5 to 37.2)	1.1 (-1.3 to 3.1)
Epilepsy	12 892.1 (10 694.5 to 15 020.3)	12 417.9 (10 438.3 to 14 478.8)	-3.7 (-8.6 to 1.3)	194.1 (160.8 to 226.1)	167.6 (140.8 to 195.3)	-13.7 (-17.9 to -9.3)*
Multiple sclerosis	1076.7 (913.9 to 1248.8)	1233.7 (1033.4 to 1436.9)	14.6 (9.9 to 18.4)*	18.1 (15.4 to 21.0)	16.9 (14.2 to 19.7)	-6.6 (-10.3 to -3.6)*
Motor neuron disease	759.2 (728.7 to 806.3)	910.0 (872.2 to 958.5)	19.9 (13.6 to 23.2)*	13.6 (13.1 to 14.4)	13.1 (12.5 to 13.8)	-3.9 (-9.0 to -1.3)*
Migraine	28 538.7 (17 584.9 to 42 479.8)	32 898.8 (20 294.8 to 48 945.4)	15.3 (14.0 to 16.6)*	435.2 (268.4 to 647.8)	438.7 (271.0 to 653.8)	0.8 (-0.1 to 1.8)
Tension-type headache	1961.3 (915.9 to 3647.4)	2260.5 (1054.6 to 4193.3)	15.3 (14.0 to 16.7)*	30.0 (14.0 to 55.5)	30.2 (14.1 to 56.0)	0.6 (-0.1 to 1.3)
Medication overuse headache	7705.4 (5100.2 to 10997.3)	9164.7 (6089.3 to 13080.8)	18.9 (15.4 to 22.8)*	123.0 (82.3 to 175.3)	123.7 (82.2 to 177.0)	0.6 (-2.3 to 3.5)
Other neurological disorders	2070.4 (1941.6 to 2302.3)	2359.5 (2216.5 to 2565.4)	14.0 (9.4 to 17.5)*	34.6 (32.2 to 38.0)	33.6 (31.3 to 36.4)	-3.1 (-6.4 to -0.4)*
Mental and substance use disorders	141 375.1 (105 843.7 to 178 447.3)	162 442.3 (121 032.0 to 205 579.7)	14.9 (14.1 to 15.7)*	2189.2 (1646.0 to 2762.8)	2183.3 (1627.1 to 2766.3)	-0.3 (-1.0 to 0.3)
Schizophrenia	13 185.0 (9 635.3 to 16 203.0)	15 516.1 (11 279.2 to 19 137.0)	17.7 (16.5 to 18.8)*	210.4 (154.7 to 258.4)	207.5 (151.1 to 255.5)	-1.4 (-2.4 to -0.5)*
Alcohol use disorders	11 566.7 (9 617.6 to 13 834.9)	11 194.3 (9 136.5 to 13 870.9)	-3.2 (-7.0 to 0.6)	183.4 (153.5 to 218.3)	149.4 (122.1 to 184.8)	-18.5 (-22.0 to -15.1)*
Drug use disorders	13 671.4 (11 292.6 to 16 046.0)	16 909.5 (14 037.6 to 19 871.6)	23.7 (18.6 to 27.2)*	207.2 (171.5 to 242.7)	223.5 (185.8 to 262.2)	7.9 (3.5 to 10.9)*
Opioid use disorders	9864.0 (8127.4 to 11516.8)	12 068.1 (9 878.0 to 14 145.1)	22.3 (17.5 to 26.1)*	150.2 (123.7 to 175.3)	159.3 (130.7 to 186.6)	6.0 (1.8 to 9.2)*
Cocaine use disorders	729.3 (558.4 to 902.4)	999.3 (773.5 to 1233.9)	37.0 (29.2 to 47.0)*	11.3 (8.7 to 14.0)	13.4 (10.4 to 16.6)	18.7 (12.1 to 27.3)*
Amphetamine use disorders	1001.3 (706.2 to 1348.3)	1402.6 (1025.2 to 1846.9)	40.1 (26.1 to 55.2)*	14.6 (10.4 to 19.6)	18.4 (13.5 to 24.2)	25.8 (13.3 to 39.4)*
Cannabis use disorders	548.1 (351.9 to 780.7)	577.2 (371.8 to 817.6)	5.3 (3.7 to 7.1)*	7.9 (5.0 to 11.2)	7.6 (4.9 to 10.7)	-3.7 (-5.0 to -2.3)*
Other drug use disorders	1528.7 (1245.5 to 1861.2)	1862.2 (1501.6 to 2274.2)	21.8 (15.7 to 27.5)*	23.1 (19.0 to 28.0)	24.8 (20.0 to 30.2)	7.1 (1.7 to 12.0)*
Depressive disorders	45 916.0 (31 684.6 to 61 591.2)	54 255.4 (37 513.6 to 72 968.9)	18.2 (17.2 to 19.2)*	726.9 (503.8 to 975.9)	734.2 (508.2 to 986.9)	1.0 (0.5 to 1.5)*
Major depressive disorder	37 544.8 (24 983.0 to 51 134.6)	44 224.4 (29 542.6 to 60 430.5)	17.8 (16.6 to 19.0)*	590.9 (395.0 to 804.5)	597.2 (399.6 to 816.6)	1.1 (0.5 to 1.7)*
Dysthymia	8 371.3 (5 537.6 to 11 983.7)	10 031.0 (6 604.4 to 14 289.3)	19.8 (18.3 to 21.5)*	136.1 (89.8 to 194.6)	137.0 (90.0 to 195.3)	0.7 (-0.2 to 1.6)

(Table 1 continues on next page)

	All-age DALYs (thousands)			Age-standardised rate (per 100 000)		
	2005	2015	Percentage change, 2005–15	2005	2015	Percentage change, 2005–15
(Continued from previous page)						
Bipolar disorder	7838·8 (4787·8 to 11674·9)	9004·7 (5501·3 to 13396·4)	14·9 (13·9 to 15·9)*	119·0 (72·8 to 176·9)	119·6 (73·2 to 177·8)	0·5 (0·0 to 0·9)
Anxiety disorders	21474·8 (14 666·7 to 29 339·6)	24 643·0 (16 799·8 to 33 714·1)	14·8 (12·8 to 16·6)*	329·2 (225·4 to 448·4)	332·4 (227·0 to 453·8)	1·0 (−0·4 to 2·3)
Eating disorders	1198·1 (806·7 to 1669·8)	1421·7 (950·5 to 1978·6)	18·7 (16·5 to 20·8)*	17·1 (11·5 to 23·7)	18·7 (12·5 to 26·0)	9·6 (7·7 to 11·4)*
Anorexia nervosa	584·9 (396·7 to 829·1)	653·0 (439·3 to 929·4)	11·7 (8·7 to 14·6)*	8·2 (5·6 to 11·6)	8·6 (5·8 to 12·2)	4·6 (1·8 to 7·4)*
Bulimia nervosa	613·3 (394·9 to 892·6)	768·7 (493·0 to 1113·8)	25·3 (23·2 to 27·6)*	8·8 (5·7 to 12·9)	10·1 (6·5 to 14·6)	14·2 (12·2 to 16·2)*
Autistic spectrum disorders	8949·8 (5978·1 to 12 300·4)	10 051·5 (6718·9 to 13 804·4)	12·3 (11·9 to 12·8)*	134·7 (90·1 to 185·0)	135·5 (90·6 to 186·0)	0·6 (0·2 to 0·9)*
Autism	5632·7 (3617·2 to 7920·0)	6335·9 (4070·1 to 8919·0)	12·5 (11·9 to 13·1)*	84·8 (54·5 to 119·2)	85·4 (54·9 to 120·3)	0·7 (0·2 to 1·2)*
Asperger syndrome and other autistic spectrum disorders	3317·1 (2204·2 to 4849·0)	3715·6 (2467·7 to 5435·2)	12·0 (11·5 to 12·5)*	49·9 (33·2 to 72·9)	50·1 (33·3 to 73·2)	0·3 (−0·1 to 0·7)
Attention-deficit hyperactivity disorder	617·9 (368·9 to 946·0)	620·1 (369·8 to 948·7)	0·4 (−0·6 to 1·2)	8·6 (5·1 to 13·2)	8·3 (5·0 to 12·7)	−3·5 (−4·4 to −2·8)*
Conduct disorder	5740·5 (3459·2 to 8923·4)	5770·5 (3471·8 to 8956·9)	0·5 (−0·3 to 1·4)	78·2 (47·2 to 121·3)	79·3 (47·7 to 123·1)	1·4 (0·7 to 2·0)*
Idiopathic developmental intellectual disability	3119·9 (1386·3 to 5416·5)	3442·1 (1503·1 to 5999·6)	10·3 (8·2 to 11·4)*	46·0 (20·4 to 80·0)	46·2 (20·1 to 80·5)	0·2 (−1·5 to 1·2)
Other mental and substance use disorders	8095·9 (5632·9 to 10919·3)	9613·4 (6698·4 to 12966·0)	18·7 (17·7 to 19·8)*	128·4 (89·5 to 172·8)	128·8 (89·8 to 173·6)	0·3 (−0·6 to 1·1)
Diabetes, urogenital, blood, and endocrine diseases	122 917·1 (106 347·2 to 142 775·0)	146 780·6 (126 480·8 to 171 189·3)	19·4 (16·4 to 22·5)*	2123·6 (1847·2 to 2453·9)	2086·8 (1803·2 to 2422·8)	−1·7 (−4·0 to 0·6)
Diabetes mellitus	49 724·5 (41 868·2 to 58 982·1)	64 134·5 (53 489·7 to 76 112·8)	29·0 (26·2 to 31·7)*	911·2 (772·1 to 1 076·0)	925·8 (776·1 to 1 096·3)	1·6 (−0·5 to 3·7)
Acute glomerulonephritis	412·3 (321·1 to 459·3)	337·9 (228·4 to 378·2)	−18·1 (−31·7 to −10·4)*	6·9 (5·4 to 7·7)	4·8 (3·2 to 5·3)	−30·6 (−42·4 to −24·1)*
Chronic kidney disease	29 488·3 (26 950·4 to 31 604·2)	35 259·7 (32 008·1 to 37 763·0)	19·6 (16·0 to 23·2)*	524·4 (482·1 to 561·0)	508·8 (462·1 to 545·1)	−3·0 (−5·8 to −0·1)*
Chronic kidney disease due to diabetes mellitus	8713·0 (7991·5 to 9466·3)	11 258·0 (10 302·9 to 12 225·2)	29·2 (25·9 to 32·5)*	159·0 (146·2 to 172·6)	163·3 (149·6 to 177·1)	2·7 (0·2 to 5·3)*
Chronic kidney disease due to hypertension	10 366·1 (9 400·8 to 10 985·1)	12 737·4 (11 488·7 to 13 553·9)	22·9 (18·6 to 27·4)*	188·4 (172·1 to 199·7)	186·1 (168·1 to 197·9)	−1·2 (−4·6 to 2·3)
Chronic kidney disease due to glomerulonephritis	7720·0 (6930·5 to 8332·1)	8136·0 (7294·2 to 8861·4)	5·4 (1·3 to 9·3)*	130·7 (117·3 to 140·8)	114·6 (103·0 to 124·6)	−12·3 (−15·6 to −9·0)*
Chronic kidney disease due to other causes	2689·3 (2208·6 to 3237·9)	3128·3 (2517·9 to 3802·9)	16·3 (13·1 to 19·2)*	46·3 (37·7 to 55·8)	44·7 (36·0 to 54·1)	−3·4 (−5·6 to −1·5)*
Urinary diseases and male infertility	7958·9 (6720·4 to 9447·1)	9653·3 (8096·4 to 11537·2)	21·3 (17·2 to 24·8)*	147·3 (123·4 to 175·7)	142·5 (119·0 to 170·6)	−3·3 (−6·3 to −0·5)*
Interstitial nephritis and urinary tract infections	3399·6 (3134·3 to 3743·4)	3950·3 (3625·0 to 4408·2)	16·2 (11·4 to 21·5)*	60·6 (56·1 to 66·3)	57·6 (52·8 to 63·9)	−5·0 (−9·0 to −0·3)*
Urolithiasis	448·5 (382·6 to 535·4)	456·1 (398·9 to 568·2)	1·7 (−4·7 to 10·8)	7·9 (6·7 to 9·4)	6·5 (5·7 to 8·1)	−17·3 (−22·5 to −9·6)*
Benign prostatic hyperplasia	2922·0 (1867·0 to 4187·6)	3792·7 (2419·5 to 5453·5)	29·8 (27·6 to 32·2)*	58·6 (37·5 to 84·1)	57·9 (37·1 to 82·9)	−1·3 (−2·8 to 0·3)
Male infertility	141·8 (57·7 to 293·6)	173·9 (69·7 to 365·9)	22·7 (19·2 to 25·9)*	2·1 (0·8 to 4·3)	2·3 (0·9 to 4·8)	9·8 (6·9 to 12·5)*
Other urinary diseases	1047·1 (861·6 to 1213·3)	1280·2 (1044·9 to 1424·5)	22·3 (10·3 to 36·4)*	18·2 (15·1 to 21·1)	18·2 (14·9 to 20·2)	0·2 (−9·9 to 11·5)
Gynaecological diseases	9292·8 (6396·3 to 13190·5)	10 255·1 (7 041·0 to 14 603·3)	10·4 (9·0 to 11·7)*	140·3 (96·5 to 199·6)	134·9 (92·6 to 192·2)	−3·8 (−4·9 to −2·8)*

(Table 1 continues on next page)

	All-age DALYs (thousands)			Age-standardised rate (per 100 000)		
	2005	2015	Percentage change, 2005–15	2005	2015	Percentage change, 2005–15
(Continued from previous page)						
Uterine fibroids	2212·2 (1393·2 to 3481·1)	2452·6 (1514·9 to 3914·8)	10·9 (8·3 to 13·1)*	34·6 (21·7 to 54·4)	32·4 (20·0 to 51·8)	-6·1 (-8·1 to -4·4)*
Polycystic ovarian syndrome	514·7 (245·2 to 972·0)	560·0 (263·4 to 1068·1)	8·8 (4·8 to 10·9)*	7·5 (3·6 to 14·2)	7·4 (3·5 to 14·0)	-2·6 (-6·5 to -0·6)*
Female infertility	275·6 (106·1 to 606·6)	344·5 (133·4 to 748·2)	25·0 (19·7 to 30·1)*	4·0 (1·6 to 8·9)	4·5 (1·7 to 9·8)	11·7 (7·2 to 16·2)*
Endometriosis	894·2 (580·2 to 1240·9)	999·0 (649·1 to 1390·3)	11·7 (10·4 to 12·9)*	13·4 (8·7 to 18·6)	13·0 (8·5 to 18·2)	-2·4 (-3·4 to -1·5)*
Genital prolapse	449·1 (228·1 to 794·4)	519·3 (258·8 to 921·5)	15·6 (11·7 to 18·4)*	8·0 (4·1 to 14·0)	7·3 (3·6 to 12·9)	-8·4 (-11·8 to -5·9)*
Premenstrual syndrome	3287·8 (2048·5 to 4917·5)	3621·6 (2248·2 to 5435·6)	10·2 (8·0 to 12·0)*	48·3 (30·1 to 72·2)	47·3 (29·4 to 71·0)	-2·0 (-3·9 to -0·4)*
Other gynaecological diseases	1659·3 (1213·1 to 2208·2)	1758·0 (1278·9 to 2342·6)	6·0 (4·0 to 8·1)*	24·6 (18·0 to 32·6)	23·0 (16·8 to 30·6)	-6·3 (-8·1 to -4·5)*
Haemoglobinopathies and haemolytic anaemias	20 211·2 (16 303·9 to 25 551·6)	20 604·4 (16 002·3 to 27 392·4)	1·9 (-11·4 to 19·6)	301·0 (243·6 to 378·6)	279·3 (217·4 to 370·5)	-7·2 (-18·9 to 8·7)
Thalassaemias	1366·6 (1131·0 to 1683·8)	1034·6 (839·9 to 1272·9)	-24·3 (-39·0 to -2·4)*	19·8 (16·4 to 24·3)	14·0 (11·4 to 17·2)	-29·0 (-42·6 to -9·0)*
Thalassaemias trait	3696·4 (2473·4 to 5333·5)	3922·2 (2605·5 to 5657·2)	6·1 (4·8 to 7·4)*	55·2 (36·9 to 79·6)	53·2 (35·3 to 76·7)	-3·6 (-4·7 to -2·6)*
Sickle cell disorders	8662·7 (6150·3 to 13 061·9)	8973·8 (5936·8 to 14 752·8)	3·6 (-23·4 to 40·6)	124·4 (88·4 to 187·0)	120·3 (79·7 to 197·7)	-3·3 (-28·4 to 31·4)
Sickle cell trait	1552·9 (1049·6 to 2211·1)	1720·3 (1153·9 to 2463·9)	10·8 (7·3 to 13·4)*	23·0 (15·5 to 32·7)	23·3 (15·6 to 33·4)	1·4 (-1·9 to 3·6)
G6PD deficiency	1284·4 (1105·7 to 1525·4)	1440·5 (1225·4 to 1706·3)	12·1 (5·3 to 21·5)*	19·8 (17·0 to 23·4)	19·3 (16·4 to 22·9)	-2·4 (-8·4 to 5·3)
G6PD trait	26·9 (18·4 to 37·1)	28·0 (19·2 to 38·9)	4·3 (0·1 to 8·8)*	0·4 (0·3 to 0·6)	0·4 (0·3 to 0·5)	-5·3 (-9·0 to -1·2)*
Other haemoglobinopathies and haemolytic anaemias	3621·4 (2874·6 to 4634·1)	3485·1 (2770·9 to 4426·5)	-3·8 (-6·2 to -1·0)*	58·5 (47·2 to 73·7)	48·8 (39·1 to 61·7)	-16·5 (-19·2 to -13·8)*
Endocrine, metabolic, blood, and immune disorders	5829·0 (5157·9 to 6691·1)	6535·7 (5812·5 to 7435·2)	12·1 (7·6 to 17·0)*	92·6 (82·5 to 105·6)	90·7 (80·9 to 103·0)	-2·1 (-5·8 to 1·7)
Musculoskeletal disorders	123 801·0 (90 494·0 to 163 616·5)	148 986·5 (108 880·7 to 197 118·2)	20·3 (19·4 to 21·3)*	2082·9 (1530·5 to 2750·7)	2065·1 (1510·9 to 2730·7)	-0·9 (-1·5 to -0·2)*
Rheumatoid arthritis	5204·7 (3777·3 to 6833·5)	6333·4 (4574·6 to 8332·9)	21·7 (18·7 to 24·7)*	90·9 (66·3 to 118·9)	89·7 (65·0 to 117·6)	-1·3 (-3·6 to 1·2)
Osteoarthritis	9562·5 (6666·3 to 13 106·3)	12 886·2 (8 990·3 to 17 634·2)	34·8 (33·6 to 36·0)*	180·4 (125·9 to 247·2)	187·4 (130·8 to 256·4)	3·9 (3·0 to 4·8)*
Low back and neck pain	80 053·5 (57 147·1 to 108 282·6)	94 941·5 (67 745·5 to 128 118·6)	18·6 (17·6 to 19·6)*	1337·3 (958·3 to 1802·5)	1309·7 (936·1 to 1765·1)	-2·1 (-2·7 to -1·4)*
Low back pain	51 243·0 (36 375·0 to 70 714·6)	60 074·8 (42 682·1 to 82 419·2)	17·2 (16·4 to 18·1)*	852·0 (605·3 to 1176·4)	829·5 (589·2 to 1138·3)	-2·6 (-3·2 to -2·0)*
Neck pain	28 810·5 (19 224·7 to 39 466·1)	34 866·7 (23 343·7 to 47 754·3)	21·0 (18·9 to 23·2)*	485·3 (327·2 to 662·1)	480·2 (322·0 to 653·6)	-1·1 (-2·3 to 0·1)
Gout	1063·1 (719·2 to 1451·4)	1342·8 (908·5 to 1843·8)	26·3 (24·6 to 28·0)*	18·9 (12·9 to 25·8)	19·0 (12·9 to 26·1)	0·6 (-0·3 to 1·5)
Other musculoskeletal disorders	27 917·2 (19 404·3 to 38 408·8)	33 482·6 (23 135·4 to 45 865·5)	19·9 (16·7 to 23·0)*	455·4 (316·9 to 623·9)	459·2 (318·0 to 630·7)	0·8 (-1·5 to 3·2)
Other non-communicable diseases	170 032·0 (132 096·5 to 221 007·5)	192 712·9 (148 719·0 to 251 205·4)	13·3 (7·1 to 16·9)*	2726·4 (2098·4 to 3558·8)	2687·2 (2068·1 to 3508·3)	-1·4 (-6·1 to 1·4)
Congenital anomalies	57 594·6 (49 489·5 to 69 397·4)	58 355·3 (52 597·9 to 64 983·3)	1·3 (-12·2 to 11·2)	826·5 (711·2 to 994·5)	781·1 (704·2 to 869·9)	-5·5 (-17·9 to 3·5)
Neural tube defects	6925·7 (5218·4 to 9487·8)	5981·2 (4589·5 to 7574·7)	-13·6 (-32·7 to 5·7)	98·3 (74·1 to 134·8)	79·9 (61·3 to 101·2)	-18·8 (-36·7 to -0·6)*
Congenital heart anomalies	26 778·2 (22 186·7 to 31 821·1)	25 706·4 (22 612·9 to 28 639·2)	-4·0 (-18·1 to 7·6)	382·7 (316·9 to 454·7)	343·9 (302·5 to 383·1)	-10·2 (-23·3 to 0·4)

(Table 1 continues on next page)

	All-age DALYs (thousands)			Age-standardised rate (per 100 000)		
	2005	2015	Percentage change, 2005–15	2005	2015	Percentage change, 2005–15
(Continued from previous page)						
Cleft lip and cleft palate	355·3 (288·5 to 424·1)	196·0 (150·8 to 259·3)	-44·8 (-54·9 to -33·7)*	5·1 (4·1 to 6·1)	2·6 (2·0 to 3·5)	-48·3 (-57·6 to -38·1)*
Down's syndrome	2379·1 (1571·3 to 3712·7)	2401·1 (1739·7 to 3291·8)	0·9 (-23·2 to 29·4)	34·8 (23·1 to 53·9)	32·2 (23·3 to 44·0)	-7·5 (-29·2 to 17·8)
Turner's syndrome	6·0 (2·1 to 12·9)	6·6 (2·4 to 14·4)	11·4 (8·7 to 14·0)*	0·1 (0·0 to 0·2)	0·1 (0·0 to 0·2)	-0·1 (-2·4 to 2·3)
Klinefelter's syndrome	1·2 (0·4 to 2·8)	1·3 (0·5 to 3·2)	13·5 (10·6 to 15·6)*	0·0 (0·0 to 0·0)	0·0 (0·0 to 0·0)	0·2 (-2·2 to 2·0)
Other chromosomal abnormalities	2110·0 (1124·3 to 4575·8)	2354·5 (1418·6 to 3905·9)	11·6 (-19·2 to 35·1)	30·4 (16·4 to 65·4)	31·5 (19·0 to 52·2)	3·6 (-24·3 to 25·0)
Other congenital anomalies	19 039·2 (16 026·4 to 23 290·1)	21 708·1 (18 750·6 to 25 714·7)	14·0 (0·3 to 26·3)*	275·0 (231·4 to 336·5)	291·0 (251·2 to 344·6)	5·8 (-6·6 to 16·7)
Skin and subcutaneous diseases	42 069·1 (27 826·1 to 62 299·6)	47 222·7 (31 372·1 to 69 571·8)	12·3 (11·5 to 13·2)*	639·6 (426·1 to 942·5)	643·4 (428·2 to 946·7)	0·6 (0·1 to 1·2)*
Dermatitis	7733·8 (5244·5 to 10760·5)	8788·0 (5957·2 to 12275·5)	13·6 (12·9 to 14·4)*	117·8 (80·2 to 164·1)	119·7 (81·3 to 166·6)	1·6 (1·0 to 2·2)*
Psoriasis	5478·8 (3816·1 to 7443·3)	6438·3 (4487·4 to 8755·9)	17·5 (16·6 to 18·4)*	87·4 (60·9 to 118·7)	87·9 (61·3 to 119·5)	0·6 (0·0 to 1·2)
Cellulitis	406·1 (251·0 to 550·4)	498·2 (315·1 to 692·8)	22·7 (11·9 to 34·6)*	6·9 (4·3 to 9·4)	7·0 (4·5 to 9·8)	1·6 (-7·4 to 11·6)
Pyoderma	1070·5 (714·3 to 1585·5)	1361·7 (932·3 to 2020·6)	27·2 (17·0 to 38·8)*	17·5 (11·7 to 25·6)	19·1 (13·1 to 28·2)	9·2 (0·6 to 18·9)*
Scabies	4944·3 (2786·4 to 8050·4)	5268·9 (2947·0 to 8616·4)	6·6 (3·8 to 9·5)*	73·0 (41·2 to 118·8)	71·1 (39·8 to 116·0)	-2·5 (-4·5 to -0·5)*
Fungal skin diseases	2457·6 (976·3 to 5214·9)	2783·3 (1101·3 to 5914·6)	13·3 (12·4 to 14·1)*	37·2 (14·7 to 79·0)	37·8 (15·0 to 80·3)	1·7 (1·4 to 2·0)*
Viral skin diseases	4977·3 (3147·1 to 7362·6)	5396·9 (3412·0 to 7962·7)	8·4 (7·9 to 9·0)*	73·7 (46·7 to 109·3)	73·0 (46·2 to 107·9)	-1·0 (-1·4 to -0·6)*
Acne vulgaris	6553·4 (3137·7 to 12159·0)	6854·0 (3271·3 to 12744·4)	4·6 (3·5 to 5·7)*	91·1 (43·7 to 169·0)	91·9 (43·9 to 171·0)	0·8 (-0·1 to 1·8)
Alopecia areata	610·5 (376·1 to 911·7)	695·6 (427·5 to 1035·9)	13·9 (13·1 to 14·7)*	9·4 (5·8 to 14·0)	9·3 (5·7 to 13·8)	-1·0 (-1·7 to -0·4)*
Pruritus	631·4 (306·3 to 1152·1)	741·6 (359·8 to 1349·6)	17·5 (15·5 to 19·4)*	10·1 (4·9 to 18·5)	10·2 (4·9 to 18·6)	0·5 (-0·7 to 1·7)
Urticaria	3718·5 (2297·3 to 5286·3)	4115·7 (2560·6 to 5836·2)	10·7 (9·5 to 12·0)*	55·5 (34·5 to 78·7)	55·5 (34·6 to 78·7)	0·0 (-0·5 to 0·4)
Decubitus ulcer	494·1 (328·7 to 602·5)	609·1 (415·1 to 738·7)	23·3 (17·3 to 33·2)*	9·8 (6·5 to 11·9)	9·4 (6·4 to 11·3)	-4·3 (-8·9 to 3·7)*
Other skin and subcutaneous diseases	2992·7 (1486·8 to 5413·5)	3671·3 (1817·7 to 6638·7)	22·7 (21·9 to 23·5)*	50·2 (24·9 to 90·8)	51·5 (25·5 to 93·2)	2·6 (2·1 to 3·3)*
Sense organ diseases	54708·4 (38 076·6 to 75 346·3)	68515·2 (47 752·5 to 94 246·6)	25·2 (24·2 to 26·4)*	993·6 (694·9 to 1361·8)	999·9 (697·1 to 1370·3)	0·6 (-0·1 to 1·3)
Glaucoma	388·4 (268·1 to 538·9)	541·3 (369·5 to 748·0)	39·4 (37·0 to 41·9)*	8·0 (5·5 to 11·2)	8·4 (5·7 to 11·6)	4·3 (2·6 to 5·8)*
Cataract	3091·3 (2213·8 to 4152·6)	3879·7 (2766·1 to 5232·4)	25·5 (24·1 to 26·9)*	62·9 (45·1 to 84·3)	60·3 (43·0 to 81·0)	-4·2 (-5·2 to -3·2)*
Macular degeneration	313·2 (219·5 to 429·6)	462·4 (327·4 to 633·5)	47·7 (45·2 to 49·9)*	6·9 (4·9 to 9·5)	7·4 (5·2 to 10·1)	6·9 (5·3 to 8·6)*
Refraction and accommodation disorders	12 046·9 (7737·6 to 18 905·4)	14 593·8 (9388·5 to 22 942·4)	21·1 (20·2 to 22·1)*	207·8 (133·9 to 326·0)	207·3 (133·7 to 324·8)	-0·3 (-0·9 to 0·4)
Age-related and other hearing loss	32 119·0 (21 992·4 to 44 438·1)	40 596·8 (27 842·2 to 56 168·9)	26·4 (24·6 to 28·3)*	589·4 (405·6 to 817·1)	595·9 (409·6 to 825·0)	1·1 (0·0 to 2·1)
Other vision loss	1352·0 (957·1 to 1852·2)	1756·4 (1247·4 to 2393·6)	29·9 (27·8 to 32·1)*	23·9 (17·0 to 32·5)	25·2 (17·9 to 34·1)	5·6 (4·3 to 6·7)*
Other sense organ diseases	5397·7 (3358·7 to 7841·8)	6684·7 (4178·4 to 9722·8)	23·8 (22·9 to 24·8)*	94·7 (59·3 to 137·2)	95·5 (59·9 to 138·6)	0·9 (0·2 to 1·5)*

(Table 1 continues on next page)

	All-age DALYs (thousands)			Age-standardised rate (per 100 000)		
	2005	2015	Percentage change, 2005–15	2005	2015	Percentage change, 2005–15
(Continued from previous page)						
Oral disorders	13 868·2 (8387·2 to 21 285·6)	16 969·6 (10 280·0 to 26 059·7)	22·4 (21·6 to 23·2)*	241·3 (147·2 to 367·9)	240·8 (146·9 to 367·8)	-0·2 (-0·5 to 0·1)
Deciduous caries	141·4 (61·3 to 279·6)	147·2 (62·8 to 292·3)	4·1 (1·6 to 5·9)*	2·1 (0·9 to 4·1)	2·0 (0·9 to 4·0)	-2·7 (-4·9 to -0·8)*
Permanent caries	1536·3 (681·5 to 2960·1)	1743·4 (773·5 to 3358·0)	13·5 (12·6 to 14·4)*	23·7 (10·5 to 45·7)	23·6 (10·5 to 45·5)	-0·4 (-1·2 to 0·4)
Periodontal diseases	2808·4 (1084·9 to 5841·7)	3520·7 (1357·6 to 7373·3)	25·4 (24·1 to 26·5)*	48·0 (18·6 to 100·0)	48·6 (18·8 to 101·2)	1·2 (0·6 to 1·8)*
Edentulism and severe tooth loss	5999·5 (4012·3 to 8293·2)	7640·3 (5092·8 to 10572·8)	27·3 (26·9 to 27·7)*	114·6 (76·7 to 158·5)	113·6 (76·0 to 157·3)	-0·8 (-1·1 to -0·6)*
Other oral disorders	3382·7 (2093·2 to 5085·7)	3918·1 (2423·4 to 5894·1)	15·8 (15·3 to 16·4)*	53·0 (32·8 to 79·5)	53·0 (32·8 to 79·6)	0·0 (-0·2 to 0·3)
Sudden infant death syndrome	1791·7 (1458·5 to 2904·4)	1650·1 (1366·4 to 2367·7)	-7·9 (-23·0 to 8·9)	25·5 (20·7 to 41·3)	22·0 (18·2 to 31·6)	-13·5 (-27·7 to 2·3)
Injuries	263 084·3 (245 566·5 to 278 621·5)	249 791·3 (231 409·2 to 266 419·2)	-5·1 (-8·5 to -0·9)*	4044·0 (3764·8 to 4300·1)	3376·1 (3120·8 to 3607·0)	-16·5 (-19·4 to -13·0)*
Transport injuries	77 146·4 (73 768·7 to 80 734·9)	72 365·5 (68 372·4 to 76 237·9)	-6·2 (-11·1 to -1·8)*	1175·5 (1121·9 to 1231·3)	971·4 (917·0 to 1024·0)	-17·4 (-21·6 to -13·5)*
Road injuries	71 941·1 (68 596·1 to 75 312·5)	67 270·4 (63 538·6 to 70 859·9)	-6·5 (-11·4 to -2·2)*	1095·3 (1042·8 to 1145·8)	903·0 (854·0 to 951·5)	-17·6 (-21·8 to -13·8)*
Pedestrian road injuries	27 185·9 (25 567·5 to 29 487·1)	24 491·1 (22 747·9 to 26 770·6)	-9·9 (-16·4 to -3·8)*	417·2 (392·7 to 452·1)	330·1 (307·3 to 360·7)	-20·9 (-26·5 to -15·5)*
Cyclist road injuries	3721·2 (3357·8 to 4119·6)	3399·8 (3022·9 to 3807·0)	-8·6 (-15·1 to -2·1)*	58·5 (52·7 to 65·1)	46·3 (41·1 to 52·0)	-20·9 (-26·3 to -15·4)*
Motorcyclist road injuries	13 655·4 (12 181·3 to 14 651·4)	13 686·2 (12 440·3 to 15 369·7)	0·2 (-7·8 to 10·6)	204·9 (182·3 to 219·9)	182·4 (165·7 to 204·3)	-10·9 (-17·9 to -1·7)*
Motor vehicle road injuries	26 257·7 (24 307·0 to 28 943·8)	24 460·3 (22 362·4 to 26 719·4)	-6·8 (-11·6 to -1·8)*	397·6 (368·7 to 437·0)	327·5 (299·3 to 357·3)	-17·6 (-21·8 to -13·2)*
Other road injuries	1120·9 (782·6 to 1331·6)	1233·1 (858·6 to 1446·6)	10·0 (-5·2 to 31·3)	17·1 (12·1 to 20·2)	16·6 (11·6 to 19·4)	-2·9 (-16·0 to 15·4)
Other transport injuries	5205·3 (4772·9 to 5728·4)	5095·1 (4507·0 to 5967·9)	-2·1 (-10·5 to 7·5)	80·2 (73·6 to 88·4)	68·4 (60·6 to 80·2)	-14·7 (-21·8 to -6·7)*
Unintentional injuries	118 001·9 (105 534·5 to 129 998·5)	107 990·9 (95 660·4 to 120 900·3)	-8·5 (-11·9 to -3·4)*	1840·5 (1642·4 to 2033·1)	1481·0 (1310·3 to 1660·5)	-19·5 (-22·2 to -15·3)*
Falls	23 912·3 (20 767·3 to 27 587·9)	26 101·6 (22 363·6 to 30 779·5)	9·2 (3·6 to 14·6)*	403·3 (349·6 to 467·0)	368·3 (315·1 to 435·0)	-8·7 (-13·1 to -4·4)*
Drowning	24 207·3 (20 715·5 to 25 812·8)	17 864·3 (15 792·7 to 19 306·8)	-26·2 (-30·6 to -20·0)*	356·2 (305·1 to 379·5)	241·1 (213·0 to 260·6)	-32·3 (-36·3 to -26·7)*
Fire, heat, and hot substances	11 737·6 (9 928·4 to 12 948·0)	10 274·1 (8 701·9 to 11 443·4)	-12·5 (-17·4 to -5·1)*	179·7 (153·4 to 198·0)	139·0 (117·7 to 155·0)	-22·6 (-26·8 to -16·6)*
Poisonings	5746·4 (3978·7 to 7215·7)	4866·9 (3240·1 to 6070·3)	-15·3 (-28·9 to 3·6)	86·6 (60·4 to 107·8)	65·5 (43·6 to 81·5)	-24·4 (-36·0 to -7·5)*
Exposure to mechanical forces	13 753·4 (12 020·2 to 15 236·7)	13 118·4 (10 878·2 to 14 847·6)	-4·6 (-11·8 to 1·7)	210·2 (183·6 to 234·3)	177·4 (146·9 to 201·3)	-15·6 (-21·7 to -10·2)*
Unintentional firearm injuries	1655·9 (1188·2 to 1837·5)	1546·4 (1101·5 to 1727·2)	-6·6 (-11·9 to -0·6)*	24·9 (18·0 to 27·6)	20·7 (14·8 to 23·1)	-16·8 (-21·4 to -11·8)*
Unintentional suffocation	2647·9 (2075·2 to 3023·5)	2495·6 (1885·3 to 2883·4)	-5·8 (-17·6 to 7·8)	39·3 (30·8 to 44·8)	33·9 (25·7 to 39·1)	-13·7 (-24·3 to -1·8)*
Other exposure to mechanical forces	9449·6 (8051·7 to 10717·9)	9076·5 (7327·9 to 10524·0)	-3·9 (-11·4 to 3·1)	146·0 (124·4 to 166·3)	122·8 (99·1 to 142·7)	-15·9 (-22·2 to -10·1)*
Adverse effects of medical treatment	5671·1 (4245·4 to 6758·2)	5384·9 (4213·4 to 6360·5)	-5·0 (-10·5 to 2·7)	89·8 (67·8 to 106·9)	74·1 (58·2 to 87·7)	-17·4 (-21·6 to -11·7)*
Animal contact	6453·3 (4501·0 to 7223·6)	5661·2 (3760·0 to 7514·8)	-12·3 (-19·9 to 10·9)	97·6 (68·4 to 109·2)	76·6 (50·8 to 101·5)	-21·5 (-28·2 to -0·7)*
Venomous animal contact	5306·6 (3504·8 to 5989·8)	4648·0 (2901·8 to 6463·8)	-12·4 (-20·6 to 12·9)	80·1 (53·1 to 90·4)	62·8 (39·2 to 87·3)	-21·6 (-29·0 to 1·1)

(Table 1 continues on next page)

	All-age DALYs (thousands)			Age-standardised rate (per 100 000)		
	2005	2015	Percentage change, 2005–15	2005	2015	Percentage change, 2005–15
(Continued from previous page)						
Non-venomous animal contact	1146·7 (920·0 to 1358·6)	1013·2 (840·1 to 1180·4)	-11·6 (-20·1 to 5·0)	17·5 (14·1 to 20·6)	13·8 (11·4 to 16·1)	-21·2 (-28·5 to -6·9)*
Foreign body	8668·4 (6684·4 to 10942·0)	8244·8 (6786·9 to 9635·2)	-4·9 (-14·1 to 4·8)	132·2 (103·4 to 165·2)	112·4 (92·8 to 131·1)	-15·0 (-22·5 to -7·3)*
Pulmonary aspiration and foreign body in airway	6632·4 (4907·7 to 8950·1)	6321·8 (5088·3 to 7704·2)	-4·7 (-15·7 to 6·1)	101·0 (76·1 to 134·5)	86·2 (69·6 to 104·7)	-14·7 (-23·9 to -6·2)*
Foreign body in eyes	48·8 (20·8 to 87·9)	56·2 (24·0 to 101·1)	15·2 (13·8 to 16·7)*	0·8 (0·3 to 1·4)	0·8 (0·3 to 1·4)	0·5 (-0·9 to 1·3)
Foreign body in other body part	1987·3 (1331·1 to 2717·2)	1866·9 (1394·9 to 2247·3)	-6·1 (-19·8 to 6·4)	30·4 (20·8 to 41·4)	25·4 (19·1 to 30·6)	-16·4 (-28·8 to -6·5)*
Environmental heat and cold exposure	4503·5 (3641·3 to 5457·2)	4234·7 (3350·6 to 5294·9)	-6·0 (-10·1 to -1·4)*	74·1 (59·9 to 89·9)	58·7 (46·4 to 73·6)	-20·7 (-24·0 to -17·1)*
Other unintentional injuries	13 348·6 (11 406·7 to 15 619·1)	12 240·0 (10 234·9 to 14 692·0)	-8·3 (-12·9 to -3·8)*	210·9 (178·2 to 249·5)	167·9 (139·9 to 202·4)	-20·4 (-24·1 to -16·6)*
Self-harm and interpersonal violence	58 615·7 (52 794·8 to 60 379·5)	55 719·4 (50 909·8 to 58 127·7)	-4·9 (-9·0 to -0·8)*	889·6 (801·0 to 916·0)	741·1 (676·7 to 773·2)	-16·7 (-20·3 to -13·0)*
Self-harm	35 819·1 (31 467·1 to 37 112·4)	34 260·9 (30 852·1 to 36 042·4)	-4·4 (-10·4 to 1·6)	551·9 (484·7 to 571·2)	457·9 (412·2 to 481·5)	-17·0 (-22·3 to -11·9)*
Interpersonal violence	22 796·5 (20 747·8 to 23 566·7)	21 458·5 (19 354·8 to 22 540·1)	-5·9 (-9·0 to -2·0)*	337·6 (308·1 to 349·1)	283·3 (255·6 to 297·4)	-16·1 (-18·8 to -12·7)*
Assault by firearm	8856·3 (7747·9 to 9172·1)	9188·1 (7854·5 to 9689·2)	3·7 (0·0 to 8·2)	129·0 (113·0 to 133·6)	120·7 (103·2 to 127·3)	-6·4 (-9·9 to -2·4)*
Assault by sharp object	5425·2 (5050·4 to 5731·6)	4546·1 (4223·1 to 4937·8)	-16·2 (-20·6 to -11·2)*	80·5 (75·1 to 85·0)	59·9 (55·6 to 65·0)	-25·6 (-29·6 to -21·2)*
Assault by other means	8515·1 (7723·3 to 9065·4)	7724·3 (6927·0 to 8395·8)	-9·3 (-14·2 to -2·7)*	128·2 (116·6 to 136·2)	102·7 (92·2 to 111·6)	-19·8 (-24·2 to -14·1)*
Forces of nature, war, and legal intervention	9320·3 (6951·1 to 11 778·4)	13715·5 (8561·9 to 18 537·5)	47·2 (-1·9 to 100·5)	138·4 (103·8 to 174·8)	182·6 (114·4 to 246·3)	31·9 (-11·6 to 79·5)
Exposure to forces of nature	6203·9 (4037·7 to 8412·0)	1435·5 (1092·6 to 1846·4)	-76·9 (-81·2 to -69·4)*	91·9 (60·2 to 124·2)	19·6 (14·9 to 25·2)	-78·7 (-82·7 to -72·0)*
Collective violence and legal intervention	3116·4 (2504·2 to 3760·8)	12 279·9 (7208·0 to 17 075·2)	294·0 (175·8 to 394·3)*	46·5 (37·4 to 56·1)	163·0 (95·9 to 226·4)	250·5 (144·1 to 341·2)*

Data in parentheses are 95% uncertainty intervals. DALYs=disability-adjusted life-years. G6PD=glucose-6-phosphate dehydrogenase. *Percentage changes that are statistically significant.

Table 1: Global all-age DALYs and age-standardised DALY rates in 2005 and 2015 with median percentage change between 2005 and 2015 for all causes

with each recording large increases in burden from 1990 to 2005, but by 2015, all measures of DALYs and relative ranks for malaria and HIV/AIDS had fallen markedly. Trends for NCDs and injuries, both in terms of ranks and changes in disease burden, were more varied. Between 1990 and 2005, total DALYs and all-age DALY rates significantly increased for many NCDs, including ischaemic heart disease, low back and neck pain, lung cancer, chronic kidney disease, and migraine. For these NCDs, their relative ranks also climbed by 2005, yet their age-standardised DALY rates either significantly decreased or remained relatively unchanged, reflecting the effects of changes in population age structure. This pattern continued through 2015 for many NCDs, and was further exemplified by Alzheimer's disease and other dementias as it rose to the 29th leading cause of global DALYs amid a significant decrease in age-standardised DALY rates. From 1990 to 2005, all three measures of DALYs significantly increased for a subset of NCDs (ie, sense organ diseases,

diabetes, depressive disorders, and other musculoskeletal disorders), which contributed to their rises in relative ranking. More heterogeneous patterns emerged for injuries; for instance, road injuries and interpersonal violence each rose in reflective ranks from 2005 to 2015, despite significant reductions in total DALYs, all-age DALY rates, and age-standardised DALY rates from each cause.

For at least one of their leading causes of DALYs in 2015, most age groups under 40 years old had a more than 31% decrease in total burden (figure 3). The causes for which DALYs largely declined included lower respiratory infections, diarrhoeal diseases, malaria, preterm birth complications, and drowning among children younger than 5 years, and HIV/AIDS and malaria for people between the ages of 5 years and 40 years. Increases in cause-specific DALYs varied more by age, with rising DALYs due to depressive disorders and drug use disorders for populations at age 20–30 years. DALYs from low back and neck pain increased since 2005 for many age groups.

	2005, at birth				2015, at birth				2015, at age 65 years			
	Females		Males		Females		Males		Females		Males	
	Life expectancy	HALE										
Global	70.73 (70.48– 70.96)	61.37 (58.61– 63.76)	65.69 (65.45– 65.93)	57.96 (55.69– 59.98)	74.83 (74.39– 75.25)	64.88 (61.96– 67.46)	69.04 (68.63– 69.42)	60.91 (58.55– 63.01)	18.46 (18.31– 18.61)	14.19 (12.98– 15.28)	15.45 (15.31– 15.59)	11.90 (10.89– 12.81)
High SDI	80.26 (80.23– 80.28)	69.62 (66.53– 72.30)	73.25 (73.20– 73.29)	64.82 (62.36– 66.96)	82.34 (82.26– 82.41)	71.33 (68.13– 74.13)	76.21 (76.10– 76.31)	67.27 (64.70– 69.54)	20.96 (20.92– 21.01)	16.18 (14.83– 17.38)	17.66 (17.62– 17.70)	13.55 (12.38– 14.58)
High-middle SDI	75.26 (75.02– 75.51)	65.85 (63.11– 68.23)	69.26 (68.96– 69.53)	61.50 (59.22– 63.59)	79.13 (78.80– 79.43)	69.09 (66.18– 71.63)	72.69 (72.25– 73.10)	64.46 (62.01– 66.60)	19.20 (19.01– 19.40)	14.90 (13.69– 16.00)	15.81 (15.58– 16.01)	12.30 (11.29– 13.21)
Middle SDI	71.83 (71.49– 72.19)	62.72 (60.08– 65.04)	66.80 (66.43– 67.19)	59.31 (57.11– 61.26)	75.71 (75.21– 76.19)	66.02 (63.24– 68.52)	69.55 (68.99– 70.14)	61.74 (59.46– 63.75)	17.33 (17.03– 17.62)	13.36 (12.19– 14.40)	14.30 (14.01– 14.59)	11.13 (10.24– 11.95)
Low-middle SDI	62.59 (61.97– 63.21)	53.65 (51.01– 56.01)	60.30 (59.69– 60.86)	52.55 (50.22– 54.66)	67.72 (66.80– 68.61)	58.18 (55.33– 60.84)	63.83 (62.99– 64.62)	55.85 (53.44– 58.01)	14.75 (14.34– 15.16)	11.00 (9.95– 12.05)	13.39 (13.06– 13.70)	10.09 (9.18– 10.94)
Low SDI	56.19 (55.16– 57.24)	48.18 (45.46– 50.55)	54.49 (53.52– 55.45)	47.34 (45.02– 49.45)	62.30 (60.22– 64.27)	53.64 (50.41– 56.49)	59.13 (56.85– 61.28)	51.66 (48.91– 54.31)	13.35 (12.36– 14.29)	9.88 (8.61– 11.06)	12.78 (11.77– 13.68)	9.55 (8.41– 10.63)
High-income	82.09 (82.07– 82.11)	71.13 (67.95– 73.87)	76.35 (76.33– 76.37)	67.48 (64.90– 69.75)	83.42 (83.38– 83.46)	72.21 (68.97– 75.02)	78.10 (78.05– 78.15)	68.91 (66.27– 71.25)	21.68 (21.65– 21.70)	16.78 (15.39– 18.00)	18.24 (18.22– 18.27)	14.03 (12.82– 15.09)
High-income North America	80.43 (80.39– 80.47)	68.84 (65.64– 71.64)	75.33 (75.28– 75.37)	65.86 (63.17– 68.24)	81.75 (81.68– 81.82)	69.80 (66.51– 72.75)	77.02 (76.95– 77.10)	67.11 (64.32– 69.65)	20.80 (20.74– 20.86)	15.56 (14.12– 16.84)	18.17 (18.12– 18.21)	13.50 (12.21– 14.67)
Canada	82.53 (82.48– 82.60)	71.39 (68.19– 74.21)	77.81 (77.75– 77.86)	68.29 (65.47– 70.74)	83.86 (83.66– 84.08)	72.49 (69.32– 75.37)	79.50 (79.28– 79.72)	69.64 (66.80– 72.16)	21.87 (21.72– 22.02)	16.77 (15.33– 18.04)	18.81 (18.68– 18.94)	14.29 (13.00– 15.44)
Greenland	71.07 (70.24– 71.91)	61.63 (59.00– 64.06)	67.44 (66.72– 68.12)	59.74 (57.50– 61.80)	74.90 (73.77– 75.71)	64.81 (61.93– 67.47)	70.00 (69.06– 70.94)	61.90 (59.46– 64.13)	16.76 (16.23– 17.09)	12.65 (11.46– 13.71)	14.26 (14.01– 14.52)	10.79 (9.82– 11.66)
USA	80.21 (80.17– 80.25)	68.57 (65.38– 71.38)	75.06 (75.00– 75.11)	65.60 (62.93– 67.96)	81.52 (81.44– 81.60)	69.50 (66.21– 72.47)	76.74 (76.66– 76.83)	66.82 (64.05– 69.35)	20.68 (20.62– 20.74)	15.42 (13.99– 16.71)	18.09 (18.04– 18.14)	13.41 (12.12– 14.58)
Australasia	83.25 (83.20– 83.31)	71.66 (68.37– 74.50)	78.57 (78.52– 78.62)	68.83 (66.11– 71.26)	84.32 (84.21– 84.44)	72.57 (69.31– 75.54)	80.09 (79.98– 80.20)	70.08 (67.24– 72.60)	22.01 (21.93– 22.09)	17.06 (15.65– 18.29)	19.11 (19.05– 19.18)	14.66 (13.42– 15.79)
Australia	83.55 (83.49– 83.61)	71.88 (68.59– 74.74)	78.76 (78.70– 78.82)	68.94 (66.19– 71.38)	84.52 (84.41– 84.65)	72.72 (70.08– 75.71)	80.19 (79.26– 80.31)	70.12 (67.25– 72.66)	22.13 (22.04– 22.22)	17.15 (15.75– 18.40)	19.16 (19.08– 19.23)	14.67 (13.43– 15.82)
New Zealand	81.80 (81.67– 81.93)	70.59 (67.45– 73.42)	77.61 (77.48– 77.73)	68.32 (65.64– 70.68)	83.29 (83.03– 83.53)	71.83 (68.60– 74.82)	79.58 (79.30– 79.88)	69.94 (67.20– 72.40)	21.39 (21.20– 21.55)	16.57 (15.20– 17.82)	18.89 (18.74– 19.06)	14.59 (13.37– 15.68)
High-income Asia Pacific	84.71 (84.67– 84.76)	74.76 (71.79– 77.29)	77.77 (77.70– 77.83)	69.68 (67.35– 71.75)	85.95 (85.85– 86.05)	75.82 (72.83– 78.39)	79.39 (79.24– 79.53)	71.05 (68.64– 73.16)	23.39 (23.32– 23.46)	18.66 (17.29– 19.85)	18.67 (18.61– 18.73)	14.63 (13.48– 15.62)
Brunei	78.95 (78.49– 79.31)	69.48 (66.76– 72.02)	74.80 (74.36– 75.15)	66.88 (64.59– 69.00)	79.36 (78.84– 79.87)	69.96 (67.21– 72.39)	75.13 (74.43– 75.87)	67.20 (64.83– 69.39)	18.77 (18.05– 19.01)	14.81 (14.53– 15.83)	15.96 (15.61– 16.33)	12.45 (11.43– 13.43)
Japan	85.34 (85.31– 85.37)	75.43 (72.48– 77.97)	78.46 (78.44– 78.49)	70.32 (67.99– 72.38)	86.35 (86.28– 86.41)	76.28 (73.33– 78.85)	79.94 (79.88– 80.00)	71.54 (69.14– 73.67)	23.78 (23.73– 23.84)	19.04 (17.65– 20.25)	18.94 (18.90– 18.98)	14.85 (13.69– 15.88)
Singapore	82.17 (81.99– 82.35)	73.34 (70.67– 75.64)	77.88 (77.70– 78.05)	70.62 (68.52– 72.48)	83.99 (83.69– 84.35)	75.02 (72.28– 77.44)	79.83 (79.45– 80.18)	72.31 (70.06– 74.21)	21.19 (20.97– 21.46)	17.24 (16.02– 18.32)	18.19 (17.94– 18.42)	14.75 (13.69– 15.65)
South Korea	81.97 (81.79– 82.15)	71.92 (68.98– 74.42)	75.22 (74.95– 75.50)	67.35 (65.02– 69.41)	84.28 (83.91– 84.64)	73.97 (70.91– 76.61)	77.53 (76.98– 78.05)	69.38 (66.95– 71.51)	21.66 (21.39– 21.91)	16.97 (15.58– 18.18)	17.33 (17.02– 17.62)	13.49 (12.34– 14.50)

(Table 2 continues on next page)

	2005, at birth				2015, at birth				2015, at age 65 years			
	Females		Males		Females		Males		Females		Males	
	Life expectancy	HALE										
(Continued from previous page)												
Western Europe	82.60 (82.59– 82.62)	71.55 (68.33– 74.39)	76.99 (76.98– 77.01)	68.13 (65.50– 70.42)	83.97 (83.91– 84.03)	72.75 (69.44– 75.62)	78.93 (78.87– 79.00)	69.82 (67.16– 72.15)	21.64 (21.60– 21.67)	16.79 (15.38– 18.01)	18.29 (18.25– 18.32)	14.20 (12.99– 15.23)
Andorra	87.91 (87.62– 88.21)	75.88 (72.38– 78.96)	80.38 (79.98– 80.77)	70.89 (68.11– 73.38)	88.44 (88.12– 88.73)	76.30 (72.75– 79.42)	81.20 (80.80– 81.57)	71.57 (68.74– 74.03)	24.99 (24.73– 25.23)	19.38 (17.77– 20.82)	19.53 (19.32– 19.74)	15.16 (13.88– 16.31)
Austria	82.22 (82.15– 82.30)	71.19 (67.98– 74.02)	76.63 (76.55– 76.71)	67.65 (65.05– 69.96)	83.66 (83.52– 83.79)	72.42 (69.15– 75.31)	78.81 (78.66– 78.95)	69.59 (66.86– 71.97)	21.34 (21.25– 21.43)	16.56 (15.16– 17.79)	18.21 (18.13– 18.29)	14.12 (12.93– 15.18)
Belgium	81.97 (81.87– 82.07)	70.88 (67.69– 73.65)	76.18 (76.08– 76.29)	67.18 (64.52– 69.46)	83.19 (82.82– 83.56)	71.96 (68.66– 74.88)	77.83 (77.37– 78.27)	68.62 (65.99– 71.05)	21.22 (20.97– 21.49)	16.29 (14.84– 17.59)	17.62 (17.36– 17.88)	13.47 (12.27– 14.58)
Cyprus	82.15 (81.97– 82.33)	71.02 (67.78– 73.89)	75.90 (75.73– 76.07)	67.09 (64.50– 69.34)	85.05 (84.80– 85.29)	73.41 (69.97– 76.38)	78.69 (78.38– 78.99)	69.38 (66.65– 71.82)	22.49 (22.31– 22.66)	17.33 (15.81– 18.67)	17.95 (17.77– 18.13)	13.78 (12.58– 14.90)
Denmark	80.50 (80.39– 80.62)	69.73 (66.63– 72.49)	76.08 (75.98– 76.19)	67.24 (64.67– 69.51)	82.44 (82.15– 82.72)	71.39 (68.26– 74.21)	78.31 (78.03– 78.60)	69.21 (66.59– 71.56)	20.43 (20.24– 20.62)	15.83 (14.49– 17.01)	17.64 (17.47– 17.80)	13.74 (12.63– 14.71)
Finland	82.43 (82.30– 82.55)	71.05 (67.72– 74.00)	75.55 (75.41– 75.68)	66.44 (63.69– 68.82)	83.87 (83.60– 84.15)	72.37 (68.92– 75.31)	77.97 (77.62– 78.32)	68.59 (65.86– 70.98)	21.57 (21.39– 21.76)	16.59 (15.12– 17.84)	17.97 (17.79– 18.16)	13.83 (12.60– 14.90)
France	83.90 (83.87– 83.94)	72.66 (69.29– 75.57)	76.81 (76.78– 76.84)	68.01 (65.38– 70.31)	85.17 (85.01– 85.33)	73.83 (70.44– 76.73)	78.39 (78.17– 78.60)	69.30 (66.55– 71.64)	23.00 (22.89– 23.11)	17.91 (16.38– 19.21)	18.69 (18.58– 18.80)	14.32 (12.99– 15.43)
Germany	82.06 (82.03– 82.10)	70.86 (67.65– 73.74)	76.58 (76.54– 76.61)	67.72 (65.12– 70.01)	83.19 (83.07– 83.30)	71.85 (68.59– 74.75)	78.36 (78.23– 78.49)	69.31 (66.67– 71.65)	20.95 (20.88– 21.03)	16.23 (14.87– 17.40)	17.80 (17.73– 17.87)	13.88 (12.74– 14.88)
Greece	82.34 (82.25– 82.43)	71.62 (68.49– 74.37)	76.82 (76.73– 76.92)	68.22 (65.71– 70.46)	83.49 (83.25– 83.74)	72.64 (69.46– 75.42)	78.42 (78.10– 78.72)	69.60 (66.70– 71.88)	20.96 (20.78– 21.14)	16.38 (15.01– 17.54)	18.30 (18.12– 18.47)	14.30 (13.13– 15.34)
Iceland	83.64 (83.28– 84.06)	72.37 (69.13– 75.24)	79.65 (79.44– 79.92)	70.50 (67.82– 72.89)	85.82 (85.38– 86.30)	74.35 (70.99– 77.33)	80.93 (80.60– 81.25)	71.60 (68.91– 73.99)	23.00 (22.63– 23.43)	17.83 (16.29– 19.21)	19.25 (19.07– 19.43)	14.93 (13.69– 16.05)
Ireland	81.77 (81.49– 82.11)	70.80 (67.53– 73.65)	77.65 (77.39– 77.86)	68.43 (65.76– 70.81)	84.31 (83.60– 85.02)	72.94 (69.60– 76.00)	79.18 (78.94– 79.51)	69.74 (66.95– 72.16)	21.98 (21.36– 22.66)	17.09 (15.66– 18.46)	18.30 (18.08– 18.61)	14.14 (12.92– 15.20)
Israel	81.73 (81.57– 81.87)	70.97 (67.90– 73.67)	77.63 (77.46– 77.77)	68.60 (65.94– 70.92)	83.82 (83.55– 84.07)	72.69 (69.47– 75.47)	80.18 (79.92– 80.47)	70.72 (67.94– 73.16)	21.29 (21.09– 21.47)	16.44 (15.05– 17.64)	19.10 (18.93– 19.28)	14.72 (13.43– 15.83)
Italy	83.73 (83.69– 83.77)	72.34 (69.02– 75.27)	78.29 (78.26– 78.33)	69.36 (66.72– 71.67)	84.57 (84.44– 84.71)	73.10 (69.76– 76.01)	79.64 (79.48– 79.78)	70.58 (67.86– 72.95)	21.90 (21.80– 21.99)	16.87 (15.44– 18.12)	18.37 (18.27– 18.45)	14.27 (13.05– 15.31)
Luxembourg	82.51 (82.32– 82.73)	70.98 (67.62– 73.95)	77.24 (77.03– 77.44)	67.78 (64.99– 70.19)	84.29 (84.01– 84.62)	72.47 (69.05– 75.42)	79.86 (79.60– 80.12)	70.09 (67.20– 72.60)	21.79 (21.59– 22.05)	16.67 (15.16– 17.96)	18.69 (18.54– 18.83)	14.31 (13.05– 15.45)
Malta	82.73 (82.33– 83.10)	71.52 (68.23– 74.41)	78.32 (77.92– 78.69)	69.06 (66.33– 71.43)	84.48 (84.15– 84.84)	72.97 (69.61– 75.95)	79.65 (79.26– 80.06)	70.17 (67.29– 72.59)	22.05 (21.80– 22.33)	17.03 (15.60– 18.34)	18.67 (18.43– 18.92)	14.48 (13.25– 15.55)
Netherlands	81.63 (81.53– 81.73)	70.43 (67.22– 73.29)	77.33 (77.24– 77.42)	68.31 (65.65– 70.67)	83.40 (83.08– 83.72)	71.93 (68.64– 74.87)	79.17 (78.86– 79.52)	69.89 (67.22– 72.30)	21.22 (21.02– 21.47)	16.28 (14.85– 17.52)	17.78 (17.59– 18.00)	13.75 (12.58– 14.80)
Norway	82.42 (82.27– 82.58)	70.96 (67.71– 73.85)	77.81 (77.66– 77.95)	68.50 (65.80– 70.87)	84.08 (83.81– 84.34)	72.56 (69.27– 75.55)	79.94 (79.68– 80.21)	70.40 (67.62– 72.90)	21.54 (21.35– 21.73)	16.65 (15.28– 17.91)	18.57 (18.41– 18.74)	14.36 (13.14– 15.48)
Portugal	81.74 (81.66– 81.83)	70.60 (67.34– 73.41)	75.06 (74.96– 75.14)	66.36 (63.75– 68.59)	83.86 (83.69– 84.02)	72.40 (69.05– 75.36)	77.65 (77.47– 77.83)	68.61 (65.93– 70.95)	21.38 (21.26– 21.49)	16.48 (15.07– 17.73)	17.71 (17.62– 17.81)	13.72 (12.59– 14.76)

(Table 2 continues on next page)

	2005, at birth				2015, at birth				2015, at age 65 years			
	Females		Males		Females		Males		Females		Males	
	Life expectancy	HALE										
(Continued from previous page)												
Spain	83.63 (83.59– 83.67)	72.81 (69.58– 75.61)	76.91 (76.87– 76.95)	68.29 (65.70– 70.55)	85.27 (85.17– 85.38)	74.27 (70.99– 77.13)	79.84 (79.72– 79.96)	70.91 (68.26– 73.23)	22.50 (22.42– 22.58)	17.58 (16.12– 18.83)	18.86 (18.79– 18.92)	14.72 (13.49– 15.77)
Sweden	82.61 (82.50– 82.73)	71.97 (68.89– 74.70)	78.42 (78.30– 78.54)	69.24 (66.58– 71.58)	83.98 (83.74– 84.25)	73.04 (69.93– 75.82)	80.25 (80.01– 80.48)	70.70 (67.98– 73.16)	21.39 (21.22– 21.59)	16.59 (15.24– 17.78)	18.72 (18.57– 18.86)	14.36 (13.13– 15.47)
Switzerland	83.72 (83.59– 83.85)	72.15 (68.81– 75.09)	78.67 (78.54– 78.80)	69.23 (66.47– 71.66)	85.18 (84.89– 85.50)	73.55 (70.15– 76.54)	80.65 (80.33– 81.01)	71.16 (68.40– 73.63)	22.52 (22.31– 22.74)	17.40 (15.85– 18.73)	19.25 (19.06– 19.47)	14.99 (13.77– 16.09)
UK	81.14 (81.11– 81.18)	70.63 (67.46– 73.34)	76.73 (76.70– 76.77)	67.86 (65.24– 70.12)	82.81 (82.71– 82.93)	72.09 (68.88– 74.81)	79.03 (78.94– 79.13)	69.86 (67.26– 72.19)	20.85 (20.77– 20.95)	16.41 (15.13– 17.54)	18.35 (18.28– 18.41)	14.38 (13.28– 15.39)
England	81.38 (81.34– 81.41)	70.91 (67.74– 73.60)	77.05 (77.02– 77.09)	68.20 (65.59– 70.47)	83.05 (82.94– 83.16)	72.37 (69.18– 75.11)	79.34 (79.24– 79.44)	70.19 (67.61– 72.53)	21.01 (20.92– 21.10)	16.57 (15.29– 17.71)	18.51 (18.44– 18.58)	14.54 (13.43– 15.54)
Northern Ireland	80.78 (80.57– 80.97)	70.40 (67.34– 73.09)	75.84 (75.64– 76.04)	67.11 (64.54– 69.40)	82.30 (81.91– 82.68)	71.74 (68.69– 74.38)	77.94 (77.50– 78.37)	68.98 (66.37– 71.31)	20.55 (20.27– 20.82)	16.17 (14.89– 17.30)	17.67 (17.45– 17.89)	13.82 (12.71– 14.82)
Scotland	79.48 (79.35– 79.61)	68.60 (65.44– 71.42)	74.35 (74.23– 74.47)	65.43 (62.83– 67.70)	81.22 (80.86– 81.62)	70.03 (66.77– 72.90)	77.15 (76.79– 77.50)	67.83 (65.20– 70.28)	19.81 (19.57– 20.06)	15.35 (14.05– 16.49)	17.46 (17.28– 17.65)	13.54 (12.45– 14.58)
Wales	80.66 (80.53– 80.79)	69.95 (66.78– 72.73)	76.19 (76.06– 76.31)	66.91 (64.19– 69.30)	82.16 (81.86– 82.46)	71.20 (67.94– 74.08)	77.98 (77.69– 78.28)	68.39 (65.59– 70.88)	20.38 (20.18– 20.58)	15.88 (14.57– 17.06)	17.63 (17.47– 17.79)	13.62 (12.50– 14.65)
Southern Latin America	79.29 (79.23– 79.34)	69.60 (66.74– 72.09)	72.64 (72.58– 72.69)	64.83 (62.54– 66.87)	80.41 (80.19– 80.63)	70.54 (67.61– 73.11)	73.98 (73.74– 74.23)	66.02 (63.62– 68.10)	19.81 (19.67– 19.94)	15.78 (14.59– 16.82)	15.83 (15.71– 15.95)	12.47 (11.47– 13.33)
Argentina	78.69 (78.61– 78.76)	69.07 (66.24– 71.55)	71.78 (71.71– 71.85)	64.04 (61.79– 66.07)	79.76 (79.51– 80.01)	70.00 (67.06– 72.53)	73.05 (72.77– 73.31)	65.20 (62.88– 67.28)	19.55 (19.40– 19.69)	15.58 (14.41– 16.62)	15.37 (15.23– 15.50)	12.10 (11.12– 12.94)
Chile	80.97 (80.87– 81.08)	71.07 (68.14– 73.59)	75.36 (75.26– 75.47)	67.21 (64.82– 69.31)	82.04 (81.53– 82.54)	71.91 (69.01– 74.53)	76.56 (75.93– 77.19)	68.28 (65.76– 70.51)	20.50 (20.15– 20.84)	16.30 (15.06– 17.39)	17.22 (16.88– 17.57)	13.59 (12.47– 14.60)
Uruguay	79.15 (78.98– 79.31)	69.51 (66.62– 71.93)	71.80 (71.65– 71.96)	64.33 (62.09– 66.28)	80.51 (80.15– 80.90)	70.65 (67.77– 73.25)	72.93 (72.50– 73.37)	65.30 (62.97– 67.25)	19.85 (19.65– 20.05)	15.80 (14.63– 16.86)	15.05 (14.85– 15.24)	11.86 (10.90– 12.68)
Central Europe, eastern Europe, and central Asia	73.64 (73.53– 73.75)	64.25 (61.48– 66.73)	62.93 (62.80– 63.06)	55.92 (53.84– 57.72)	77.31 (77.09– 77.50)	67.31 (64.36– 69.91)	67.95 (67.71– 68.21)	60.13 (57.76– 62.14)	17.86 (17.77– 17.94)	13.74 (12.56– 14.77)	13.97 (13.90– 14.04)	10.57 (9.60– 11.43)
Eastern Europe	72.72 (72.63– 72.80)	63.16 (60.37– 65.68)	59.78 (59.65– 59.91)	53.18 (51.24– 54.89)	76.70 (76.44– 76.94)	66.44 (63.45– 69.08)	65.73 (65.39– 66.05)	58.15 (55.84– 60.13)	17.40 (17.27– 17.52)	13.30 (12.12– 14.33)	13.10 (12.98– 13.22)	9.84 (8.90– 10.68)
Belarus	74.89 (74.62– 75.15)	65.58 (62.91– 68.05)	62.76 (62.41– 63.09)	55.92 (53.92– 57.70)	77.07 (76.58– 77.58)	67.38 (64.53– 69.96)	65.02 (64.33– 65.74)	57.85 (55.63– 59.94)	17.05 (16.77– 17.32)	13.19 (12.06– 14.23)	11.75 (11.47– 12.04)	8.86 (8.00– 9.68)
Estonia	77.67 (77.51– 77.84)	67.79 (64.90– 70.36)	67.12 (66.93– 67.29)	59.50 (57.27– 61.47)	81.43 (80.99– 81.88)	71.08 (68.17– 73.78)	73.44 (73.07– 73.83)	64.89 (62.39– 67.14)	19.70 (19.35– 20.07)	15.32 (14.07– 16.50)	15.61 (15.39– 16.50)	11.87 (10.77– 12.85)
Latvia	76.63 (76.48– 76.78)	66.98 (64.16– 69.53)	65.47 (65.32– 65.62)	58.12 (55.95– 60.04)	79.73 (79.34– 80.06)	69.63 (66.64– 72.23)	70.59 (70.17– 70.99)	62.48 (60.12– 64.65)	19.01 (18.79– 19.20)	14.70 (13.42– 15.80)	14.55 (14.37– 14.72)	10.99 (9.97– 11.90)
Lithuania	77.71 (77.60– 77.82)	67.74 (64.85– 70.37)	65.71 (65.60– 65.82)	58.22 (55.95– 60.15)	80.39 (80.16– 80.61)	70.02 (67.03– 72.76)	69.72 (69.47– 70.02)	61.55 (59.11– 63.70)	19.63 (19.49– 19.76)	15.11 (13.80– 16.26)	14.75 (14.64– 14.87)	10.97 (9.87– 11.94)
Moldova	73.43 (73.13– 73.69)	64.40 (61.73– 66.76)	65.55 (65.21– 65.82)	58.37 (56.23– 60.27)	77.70 (77.29– 78.06)	67.96 (65.13– 70.54)	69.93 (69.47– 70.35)	62.06 (59.81– 64.18)	17.39 (17.21– 17.58)	13.53 (12.39– 14.53)	14.04 (13.91– 14.18)	10.71 (9.76– 11.57)

(Table 2 continues on next page)

	2005, at birth				2015, at birth				2015, at age 65 years			
	Females		Males		Females		Males		Females		Males	
	Life expectancy	HALE										
(Continued from previous page)												
Russia	72.21 (72.11– 72.30)	62.43 (59.56– 64.98)	58.71 (58.54– 58.87)	52.21 (50.31– 53.88)	76.57 (76.22– 76.90)	66.02 (62.89– 68.71)	65.30 (64.85– 65.77)	57.71 (55.43– 59.76)	17.48 (17.29– 17.64)	13.26 (12.03– 14.31)	13.12 (12.95– 13.30)	9.83 (8.86– 10.68)
Ukraine	73.16 (72.91– 73.37)	64.21 (61.62– 66.53)	61.36 (61.08– 61.60)	54.64 (52.67– 56.37)	76.45 (75.94– 76.93)	66.93 (64.21– 69.44)	66.26 (65.56– 66.93)	58.71 (56.38– 60.69)	16.95 (16.73– 17.18)	13.15 (12.05– 14.14)	12.97 (12.77– 13.18)	9.82 (8.90– 10.63)
Central Europe	77.90 (77.86– 77.94)	68.36 (65.55– 70.83)	70.48 (70.44– 70.52)	62.43 (60.05– 64.51)	80.45 (80.31– 80.58)	70.52 (67.59– 73.06)	73.43 (73.25– 73.59)	64.92 (62.42– 67.13)	19.05 (18.98– 19.11)	14.80 (13.54– 15.90)	15.33 (15.26– 15.39)	11.65 (10.59– 12.59)
Albania	79.08 (78.63– 79.52)	69.12 (66.18– 71.74)	73.49 (72.96– 74.00)	64.65 (62.01– 66.95)	81.32 (80.71– 81.95)	71.17 (68.10– 73.74)	74.98 (74.10– 75.87)	66.20 (63.43– 68.64)	20.06 (19.65– 20.47)	15.65 (14.28– 16.89)	16.09 (15.36– 16.99)	12.34 (11.10– 13.63)
Bosnia and Herzegovina	79.89 (79.75– 80.05)	69.96 (67.03– 72.56)	74.05 (73.91– 74.23)	65.08 (62.52– 67.36)	81.82 (81.45– 82.21)	71.51 (68.53– 74.24)	76.00 (75.72– 76.30)	66.69 (64.02– 69.04)	20.09 (19.81– 20.38)	15.56 (14.25– 16.76)	16.64 (16.51– 16.81)	12.59 (11.45– 13.61)
Bulgaria	76.22 (76.11– 76.34)	67.07 (64.38– 69.46)	69.07 (68.95– 69.17)	61.35 (59.08– 63.34)	78.36 (77.90– 78.80)	68.88 (66.09– 71.34)	71.37 (70.84– 71.87)	63.34 (60.91– 65.43)	17.54 (17.26– 17.79)	13.67 (12.53– 14.68)	14.14 (13.91– 14.37)	10.78 (9.80– 11.69)
Croatia	79.21 (79.07– 79.35)	69.34 (66.46– 71.92)	72.01 (71.89– 72.12)	63.67 (61.24– 65.84)	80.92 (80.61– 81.26)	70.74 (67.69– 73.39)	74.65 (74.31– 75.00)	65.87 (63.30– 68.12)	18.92 (18.71– 19.15)	14.64 (13.35– 15.72)	15.39 (15.20– 15.58)	11.70 (10.64– 12.64)
Czech Republic	79.37 (79.28– 79.47)	69.51 (66.58– 72.10)	72.98 (72.90– 73.06)	64.55 (62.06– 66.73)	81.65 (81.47– 81.83)	71.41 (68.37– 74.06)	75.92 (75.73– 76.11)	66.94 (64.35– 69.27)	19.46 (19.33– 19.59)	15.05 (13.78– 16.19)	16.06 (15.96– 16.17)	12.13 (11.00– 13.14)
Hungary	77.25 (77.14– 77.34)	67.48 (64.65– 70.00)	68.82 (68.73– 68.91)	61.06 (58.81– 63.06)	79.95 (79.55– 80.34)	69.76 (66.77– 72.32)	73.22 (72.78– 73.68)	64.71 (62.24– 66.94)	19.00 (18.75– 19.23)	14.59 (13.27– 15.70)	15.41 (15.20– 15.63)	11.65 (10.56– 12.61)
Macedonia	76.41 (76.25– 76.57)	67.48 (64.89– 69.82)	71.36 (71.20– 71.51)	63.45 (61.07– 65.54)	79.14 (78.63– 79.65)	69.66 (66.84– 72.23)	73.92 (73.30– 74.51)	65.54 (63.08– 67.75)	17.94 (17.65– 18.28)	13.95 (12.78– 15.05)	15.06 (14.67– 15.47)	11.50 (10.47– 12.50)
Montenegro	77.96 (77.70– 78.22)	68.56 (65.82– 71.10)	71.99 (71.77– 72.22)	63.70 (61.25– 65.88)	80.18 (79.69– 80.68)	70.44 (67.53– 73.02)	74.21 (73.61– 74.86)	65.59 (63.03– 67.89)	18.94 (18.63– 19.25)	14.76 (13.53– 15.90)	15.64 (15.26– 16.09)	11.95 (10.83– 12.96)
Poland	79.36 (79.31– 79.41)	69.57 (66.70– 72.07)	70.78 (70.73– 70.83)	62.67 (60.24– 64.74)	81.61 (81.39– 81.81)	71.53 (68.56– 74.11)	73.45 (73.23– 73.69)	64.96 (62.44– 67.15)	19.98 (19.86– 20.10)	15.51 (14.22– 16.67)	15.56 (14.46– 15.66)	11.82 (10.74– 12.79)
Romania	76.08 (76.00– 76.17)	66.92 (64.24– 69.30)	68.82 (68.75– 68.90)	61.04 (58.74– 63.06)	78.99 (78.59– 79.36)	69.44 (66.68– 71.90)	71.52 (71.03– 72.00)	63.40 (60.98– 65.54)	18.15 (17.92– 18.37)	14.25 (13.12– 15.26)	14.55 (14.34– 14.77)	11.14 (10.15– 12.04)
Serbia	75.94 (75.71– 76.15)	66.88 (64.26– 69.26)	70.58 (70.36– 70.80)	62.59 (60.22– 64.67)	78.87 (78.69– 79.05)	69.23 (66.44– 71.75)	73.48 (73.20– 73.71)	64.90 (62.35– 67.11)	17.88 (17.77– 17.99)	13.85 (13.68– 14.89)	15.18 (15.02– 15.31)	11.52 (10.46– 12.47)
Slovakia	78.18 (78.06– 78.30)	68.72 (65.92– 71.18)	70.28 (70.16– 70.39)	62.14 (59.72– 64.26)	80.92 (80.69– 81.14)	70.92 (68.01– 73.52)	73.98 (73.74– 74.21)	65.21 (62.65– 67.47)	19.18 (19.03– 19.33)	14.90 (13.67– 16.05)	15.38 (15.26– 15.49)	11.60 (10.53– 12.55)
Slovenia	81.07 (80.92– 81.20)	70.80 (67.77– 73.46)	73.89 (73.76– 74.01)	65.25 (62.75– 67.47)	83.84 (83.62– 84.05)	73.21 (70.12– 75.94)	77.94 (77.71– 78.18)	68.66 (65.91– 71.09)	21.31 (21.16– 21.47)	16.54 (15.13– 17.77)	17.55 (17.42– 17.68)	13.39 (12.17– 14.47)
Central Asia	71.47 (71.12– 71.80)	62.79 (60.19– 65.08)	63.62 (63.27– 63.98)	56.73 (54.69– 58.52)	75.30 (74.77– 75.76)	65.96 (63.27– 68.44)	67.41 (66.85– 67.91)	59.93 (57.71– 61.94)	16.95 (16.75– 17.17)	13.14 (12.03– 14.15)	13.48 (13.32– 13.66)	10.35 (9.44– 11.16)
Armenia	75.32 (74.82– 75.77)	65.81 (63.07– 68.24)	68.18 (67.64– 68.75)	60.65 (58.47– 62.68)	78.37 (77.89– 78.95)	68.40 (65.51– 70.99)	70.75 (70.05– 71.47)	62.84 (60.47– 65.01)	18.06 (17.85– 18.44)	13.83 (12.63– 14.95)	14.37 (14.13– 14.66)	10.99 (9.99– 11.87)
Azerbaijan	71.39 (70.61– 72.13)	62.63 (59.91– 65.16)	65.10 (64.27– 65.92)	58.10 (55.95– 60.12)	76.32 (75.59– 77.07)	66.67 (63.73– 69.41)	69.57 (68.51– 70.58)	61.78 (59.33– 63.98)	17.57 (17.25– 17.90)	13.56 (12.32– 14.64)	14.46 (13.77– 14.96)	11.12 (10.05– 12.10)

(Table 2 continues on next page)

	2005, at birth				2015, at birth				2015, at age 65 years			
	Females		Males		Females		Males		Females		Males	
	Life expectancy	HALE										
(Continued from previous page)												
Georgia	76.02 (75.39– 76.63)	67.04 (64.39– 69.40)	67.40 (66.60– 68.21)	60.26 (58.09– 62.26)	78.04 (77.27– 78.74)	68.48 (65.63– 71.10)	67.86 (66.87– 69.05)	60.52 (58.23– 62.66)	18.13 (17.73– 18.55)	14.12 (12.89– 15.23)	13.23 (12.89– 13.73)	10.15 (9.21– 11.06)
Kazakhstan	71.06 (70.70– 71.40)	62.23 (59.70– 64.57)	59.79 (59.41– 60.14)	53.24 (51.28– 54.98)	75.03 (74.05– 75.95)	65.52 (62.54– 68.31)	65.29 (64.11– 66.49)	57.91 (55.58– 60.11)	16.49 (16.00– 16.99)	12.70 (11.53– 13.82)	12.66 (12.21– 13.13)	9.60 (8.68– 10.47)
Kyrgyzstan	70.68 (70.16– 71.20)	62.06 (59.52– 64.39)	62.56 (61.96– 63.23)	55.92 (53.99– 57.76)	74.10 (73.44– 74.71)	64.92 (62.25– 67.31)	65.60 (64.83– 66.41)	58.62 (56.46– 60.57)	16.13 (15.81– 16.47)	12.55 (11.47– 13.49)	12.64 (12.31– 12.98)	9.79 (8.91– 10.58)
Mongolia	67.72 (67.00– 68.42)	60.10 (57.87– 62.07)	59.95 (59.16– 60.71)	53.54 (51.58– 55.43)	71.87 (71.03– 72.67)	63.55 (61.09– 65.93)	62.78 (61.90– 63.63)	55.92 (53.79– 57.96)	15.12 (14.80– 15.43)	11.73 (10.76– 12.66)	11.78 (10.76– 12.05)	8.94 (8.13– 9.73)
Tajikistan	70.16 (69.42– 70.88)	61.79 (59.20– 64.06)	65.92 (65.24– 66.65)	58.68 (56.48– 60.63)	74.80 (73.94– 75.53)	65.67 (63.00– 68.18)	70.12 (69.20– 70.96)	62.17 (59.71– 64.38)	16.91 (16.48– 17.22)	13.20 (12.13– 14.21)	15.04 (14.71– 15.31)	11.61 (10.61– 12.52)
Turkmenistan	69.51 (68.21– 70.74)	61.31 (58.75– 63.67)	62.12 (60.77– 63.27)	55.49 (53.35– 57.52)	73.69 (72.71– 74.46)	64.83 (62.09– 67.28)	66.33 (65.35– 67.15)	59.11 (56.86– 61.15)	16.78 (16.53– 17.01)	13.09 (11.99– 14.06)	13.57 (13.32– 13.79)	10.47 (9.58– 11.31)
Uzbekistan	71.50 (70.65– 72.29)	62.93 (60.29– 65.29)	65.48 (64.61– 66.40)	58.39 (56.15– 60.45)	75.20 (74.34– 76.03)	66.03 (63.32– 68.49)	68.37 (67.51– 69.27)	60.85 (58.49– 63.00)	16.77 (16.40– 17.33)	13.10 (11.98– 14.12)	13.69 (13.42– 14.00)	10.59 (9.67– 11.42)
Latin America and Caribbean	76.49 (76.28– 76.69)	66.42 (63.49– 69.02)	70.45 (70.24– 70.68)	62.25 (59.82– 64.38)	78.25 (77.87– 78.62)	67.88 (64.80– 70.50)	72.20 (71.73– 72.62)	63.76 (61.28– 65.91)	18.85 (18.69– 19.02)	14.38 (13.08– 15.51)	16.25 (16.11– 16.40)	12.45 (11.37– 13.42)
Central Latin America	77.58 (77.40– 77.75)	67.68 (64.81– 70.20)	71.92 (71.72– 72.11)	63.73 (61.30– 65.83)	78.81 (78.44– 79.15)	68.69 (65.67– 71.26)	73.28 (72.84– 73.72)	64.88 (62.36– 67.07)	19.06 (18.95– 19.19)	14.51 (13.21– 15.68)	16.97 (16.86– 17.09)	13.05 (11.91– 14.05)
Colombia	78.26 (78.06– 78.46)	68.20 (65.23– 70.78)	72.02 (71.80– 72.23)	64.05 (61.65– 66.08)	80.80 (80.35– 81.25)	70.21 (67.06– 72.90)	75.15 (74.58– 75.65)	66.65 (64.15– 68.88)	20.06 (19.85– 20.28)	15.33 (15.52)	17.46 (17.28– 17.64)	13.55 (12.44– 14.57)
Costa Rica	81.47 (81.22– 81.69)	70.96 (67.85– 73.65)	76.56 (76.27– 76.80)	67.84 (65.19– 70.12)	82.60 (82.12– 83.05)	71.84 (68.61– 74.60)	78.08 (77.50– 78.58)	69.11 (66.46– 71.44)	21.24 (20.99– 21.51)	16.24 (14.85– 17.48)	18.90 (18.66– 19.12)	14.67 (13.44– 15.77)
El Salvador	77.02 (76.56– 77.45)	66.63 (63.59– 69.35)	68.79 (68.07– 69.49)	60.84 (58.45– 62.88)	78.94 (78.06– 79.85)	68.24 (64.99– 71.16)	70.57 (69.32– 71.94)	62.56 (60.03– 64.95)	19.55 (19.10– 20.06)	14.80 (13.36– 16.08)	17.79 (17.33– 18.24)	13.74 (12.56– 14.87)
Guatemala	73.16 (72.79– 73.54)	63.27 (60.37– 65.90)	66.41 (66.08– 66.75)	58.31 (55.91– 60.44)	75.20 (73.54– 77.05)	65.09 (61.95– 68.07)	69.81 (67.75– 71.79)	61.33 (58.45– 64.12)	17.38 (16.41– 18.60)	13.02 (12.33– 14.44)	17.18 (16.45– 17.95)	13.09 (11.85– 14.29)
Honduras	72.00 (69.46– 74.83)	63.00 (59.75– 66.13)	70.45 (67.22– 72.91)	62.58 (59.11– 65.60)	73.95 (61.55– 76.68)	64.71 (68.69– 76.01)	72.09 (70.73– 74.71)	64.07 (62.82– 67.10)	16.30 (15.18– 17.86)	12.48 (11.19– 13.99)	16.08 (14.54– 17.25)	12.49 (11.04– 13.84)
Mexico	77.72 (77.45– 77.95)	68.16 (65.42– 70.62)	72.54 (72.25– 72.81)	64.56 (62.22– 66.64)	78.32 (77.98– 78.60)	68.64 (65.82– 71.11)	73.43 (73.04– 73.75)	65.26 (62.82– 67.38)	18.65 (18.59– 18.70)	14.24 (15.37)	16.78 (16.74– 16.83)	12.91 (11.78– 13.91)
Nicaragua	79.26 (78.69– 79.76)	69.03 (65.97– 71.74)	73.74 (73.07– 74.37)	65.25 (62.65– 67.57)	80.76 (79.93– 81.68)	70.31 (67.22– 73.07)	75.01 (73.80– 76.26)	66.47 (63.72– 69.01)	20.76 (20.25– 21.30)	15.88 (14.42– 17.17)	18.30 (17.75– 18.85)	14.19 (12.90– 15.41)
Panama	79.72 (79.09– 80.35)	69.27 (66.25– 72.03)	74.77 (73.98– 75.48)	66.08 (63.42– 68.38)	81.01 (79.82– 82.22)	75.48 (73.89– 73.23)	66.63 (63.75– 76.95)	20.93 (20.19– 21.70)	15.90 (14.35– 17.37)	18.34 (17.55– 19.16)	14.11 (12.76– 15.39)	14.11 (12.76– 15.39)
Venezuela	78.22 (78.11– 78.32)	67.39 (64.28– 70.19)	71.17 (71.06– 71.27)	61.93 (59.15– 64.31)	79.16 (78.00– 80.36)	68.14 (64.68– 71.24)	70.69 (69.02– 72.41)	61.62 (58.62– 64.44)	19.37 (18.64– 20.13)	14.58 (13.09– 15.96)	16.05 (15.37– 16.77)	12.05 (10.85– 13.25)
Andean Latin America	76.76 (76.30– 77.23)	66.92 (64.06– 69.48)	73.09 (72.59– 73.57)	64.22 (61.57– 66.57)	79.01 (78.22– 79.82)	68.86 (65.87– 71.66)	75.40 (74.51– 76.28)	66.26 (63.43– 68.85)	19.24 (18.74– 19.76)	14.90 (13.63– 16.14)	17.61 (17.15– 18.06)	13.44 (12.14– 14.60)

(Table 2 continues on next page)

	2005, at birth				2015, at birth				2015, at age 65 years			
	Females		Males		Females		Males		Females		Males	
	Life expectancy	HALE										
(Continued from previous page)												
Bolivia	71.40 (69.86– 72.67)	62.29 (59.64– 64.88)	69.02 (67.91– 70.15)	60.74 (58.14– 63.17)	74.22 (72.07– 76.43)	64.65 (61.20– 67.81)	72.11 (69.61– 74.23)	63.35 (60.22– 66.40)	16.64 (15.46– 18.07)	12.80 (11.36– 14.26)	16.33 (15.07– 17.17)	12.41 (11.10– 13.67)
Ecuador	76.85 (76.33– 77.36)	66.94 (63.96– 69.56)	71.53 (70.79– 72.19)	62.95 (60.31– 65.29)	78.56 (77.59– 79.50)	68.39 (65.41– 71.20)	73.35 (72.06– 74.39)	64.57 (61.56– 67.16)	18.94 (18.40– 19.46)	14.57 (13.33– 15.81)	16.49 (15.80– 17.02)	12.48 (11.14– 13.66)
Peru	78.80 (78.19– 79.39)	68.71 (65.77– 71.33)	75.55 (74.73– 76.36)	66.30 (63.53– 68.76)	81.10 (80.01– 82.23)	70.74 (67.67– 73.66)	77.85 (76.38– 79.13)	68.37 (65.44– 71.13)	20.44 (19.71– 21.22)	15.93 (14.52– 17.30)	18.78 (18.02– 19.53)	14.42 (13.06– 15.70)
Caribbean	72.72 (71.92– 73.44)	63.17 (60.25– 65.76)	69.08 (68.51– 69.68)	60.88 (58.42– 63.12)	75.21 (73.88– 76.31)	65.16 (61.96– 67.91)	70.92 (69.92– 71.88)	62.43 (59.83– 64.66)	18.22 (17.84– 18.59)	14.04 (12.77– 15.10)	16.18 (15.97– 16.43)	12.47 (11.39– 13.43)
Antigua and Barbuda	77.20 (76.51– 78.00)	67.21 (64.20– 69.87)	72.45 (71.58– 73.41)	63.69 (61.07– 66.03)	78.85 (78.10– 79.78)	68.54 (65.44– 71.37)	73.78 (72.68– 75.33)	64.92 (62.16– 67.47)	18.43 (18.03– 19.10)	14.28 (13.02– 15.46)	15.57 (14.73– 16.84)	11.90 (10.54– 13.37)
The Bahamas	74.43 (73.70– 75.22)	64.60 (61.75– 67.14)	68.91 (67.92– 69.86)	60.65 (58.19– 62.97)	76.41 (75.13– 77.73)	66.24 (63.14– 69.13)	70.87 (69.42– 72.31)	62.47 (59.69– 65.04)	17.58 (16.97– 18.24)	13.49 (12.26– 14.70)	15.42 (14.86– 16.14)	11.79 (10.61– 12.96)
Barbados	76.96 (76.19– 77.73)	66.70 (63.63– 69.54)	72.90 (72.17– 73.53)	64.11 (61.48– 66.54)	77.56 (76.24– 78.67)	67.12 (63.92– 69.93)	73.63 (72.28– 74.85)	64.84 (62.02– 67.46)	18.05 (17.34– 18.64)	13.81 (12.54– 15.06)	16.09 (15.12– 16.78)	12.29 (10.97– 13.53)
Belize	73.08 (72.37– 73.77)	63.66 (60.85– 66.10)	67.27 (66.44– 68.11)	59.58 (57.17– 61.66)	74.73 (73.67– 75.77)	65.00 (62.11– 67.76)	69.05 (67.70– 70.29)	61.18 (58.61– 63.64)	16.75 (16.23– 17.30)	12.97 (11.80– 14.04)	14.44 (13.90– 15.11)	11.19 (10.17– 12.20)
Bermuda	79.60 (79.07– 80.17)	69.18 (66.21– 72.01)	73.35 (72.73– 74.01)	64.74 (62.22– 67.06)	82.23 (81.24– 83.29)	71.28 (68.14– 74.12)	75.00 (73.93– 75.97)	66.15 (63.61– 68.57)	20.38 (19.75– 21.11)	15.80 (14.45– 17.03)	16.27 (15.84– 16.62)	12.53 (11.43– 13.52)
Cuba	79.14 (79.04– 79.22)	68.94 (65.93– 71.55)	75.19 (75.09– 75.27)	66.68 (64.14– 68.92)	80.60 (80.24– 80.97)	70.06 (66.98– 72.80)	75.87 (75.52– 76.25)	67.22 (64.76– 69.45)	19.56 (19.34– 19.79)	15.16 (13.88– 16.31)	16.68 (16.50– 16.87)	13.01 (11.96– 13.95)
Dominica	77.15 (76.26– 77.93)	66.87 (63.91– 69.60)	71.36 (70.25– 72.27)	62.42 (59.76– 64.84)	76.81 (75.46– 78.06)	66.52 (63.28– 69.40)	69.71 (67.83– 71.50)	61.07 (58.09– 63.74)	18.29 (17.55– 18.86)	14.04 (12.66– 15.26)	14.93 (14.01– 15.94)	11.25 (9.97– 12.50)
Dominican Republic	76.16 (75.51– 76.65)	66.32 (63.46– 68.91)	70.75 (70.04– 71.40)	62.43 (59.94– 64.68)	77.93 (77.41– 78.30)	67.89 (64.83– 70.38)	72.82 (71.86– 73.45)	64.36 (61.78– 66.67)	19.04 (18.91– 19.11)	14.78 (13.55– 15.85)	16.79 (16.51– 16.90)	12.99 (11.86– 13.96)
Grenada	73.36 (72.61– 74.08)	63.76 (60.97– 66.16)	67.21 (66.38– 67.95)	59.21 (56.81– 61.43)	74.28 (72.93– 75.36)	64.40 (61.31– 67.09)	68.78 (67.50– 70.02)	60.56 (57.89– 62.93)	16.30 (15.74– 16.82)	12.50 (11.33– 13.54)	13.99 (13.53– 14.41)	10.61 (9.51– 11.52)
Guyana	67.83 (66.96– 68.68)	58.96 (56.30– 61.42)	61.28 (60.22– 62.33)	54.11 (51.88– 56.20)	70.47 (68.95– 71.86)	61.22 (61.70– 64.03)	63.47 (61.43– 65.26)	56.08 (53.43– 58.62)	14.88 (14.21– 15.47)	11.43 (10.35– 12.47)	12.55 (12.03– 13.14)	9.57 (8.67– 10.50)
Haiti	59.86 (57.33– 62.32)	51.93 (48.83– 54.97)	60.30 (58.18– 62.80)	52.84 (50.10– 55.64)	64.74 (61.22– 68.20)	55.81 (51.95– 59.24)	63.91 (59.20– 67.20)	55.75 (52.32– 59.08)	13.37 (11.94– 15.07)	10.09 (8.77– 11.46)	14.01 (12.88– 15.32)	10.54 (9.30– 11.84)
Jamaica	76.72 (75.84– 77.43)	66.45 (63.40– 69.28)	73.96 (72.94– 74.84)	64.93 (62.19– 67.48)	76.97 (75.72– 78.20)	66.55 (63.15– 69.48)	73.03 (71.74– 74.36)	64.18 (61.35– 66.78)	18.00 (17.35– 18.77)	13.85 (12.48– 15.08)	16.43 (15.55– 17.11)	12.65 (11.39– 13.78)
Puerto Rico	80.24 (80.07– 80.41)	69.47 (66.33– 72.22)	72.85 (72.67– 73.00)	63.94 (61.31– 66.24)	81.77 (81.16– 82.37)	70.70 (67.45– 73.61)	75.03 (74.35– 75.73)	65.70 (62.92– 68.26)	20.43 (20.03– 20.83)	15.71 (14.34– 16.94)	17.42 (17.11– 17.72)	13.17 (11.95– 14.31)
Saint Lucia	76.14 (75.26– 77.06)	66.13 (63.23– 68.75)	70.19 (69.24– 71.13)	61.74 (59.30– 63.99)	77.09 (75.75– 78.29)	66.91 (63.83– 69.68)	72.52 (71.02– 73.92)	63.78 (61.01– 66.41)	17.82 (17.13– 18.53)	13.74 (12.49– 14.93)	16.08 (15.41– 16.73)	12.35 (11.12– 13.45)
Saint Vincent and the Grenadines	73.00 (72.20– 73.75)	63.48 (60.67– 65.95)	68.03 (67.29– 68.71)	59.90 (57.52– 62.10)	73.95 (72.82– 75.00)	64.26 (61.26– 66.93)	68.77 (67.61– 69.89)	60.59 (58.00– 62.87)	16.11 (15.64– 16.62)	12.39 (11.28– 13.41)	14.09 (13.51– 14.61)	10.71 (9.59– 11.71)

(Table 2 continues on next page)

	2005, at birth				2015, at birth				2015, at age 65 years			
	Females		Males		Females		Males		Females		Males	
	Life expectancy	HALE										
(Continued from previous page)												
Suriname	72.74 (71.94– 73.56)	63.29 (60.51– 65.80)	67.08 (66.13– 68.21)	59.30 (56.87– 61.54)	75.34 (73.89– 76.55)	65.50 (62.51– 68.45)	68.82 (67.09– 70.46)	60.87 (58.15– 63.47)	17.81 (16.90– 18.46)	13.79 (12.48– 15.07)	14.97 (13.93– 15.94)	11.60 (10.37– 12.83)
Trinidad and Tobago	74.55 (74.07– 74.94)	64.63 (61.71– 67.22)	67.83 (67.34– 68.24)	59.72 (57.26– 61.89)	76.11 (74.90– 77.24)	65.86 (62.82– 68.63)	68.88 (67.40– 70.26)	60.56 (57.82– 63.02)	17.70 (17.11– 18.26)	13.58 (12.32– 14.73)	14.01 (13.42– 14.55)	10.64 (9.59– 11.68)
Virgin Islands	78.01 (77.51– 78.52)	67.81 (64.87– 70.40)	70.91 (70.26– 71.53)	62.41 (59.95– 64.67)	78.72 (77.88– 79.47)	68.35 (65.40– 71.18)	70.34 (68.90– 71.64)	62.04 (59.52– 64.54)	18.47 (18.11– 18.79)	14.26 (13.03– 15.36)	14.52 (13.95– 15.13)	11.03 (10.04– 12.07)
Tropical Latin America	76.21 (75.81– 76.58)	65.73 (62.71– 68.45)	68.77 (68.38– 69.19)	60.66 (58.25– 62.82)	78.18 (77.61– 78.70)	67.36 (64.20– 70.06)	70.75 (70.08– 71.38)	62.38 (59.89– 64.58)	18.72 (18.37– 19.08)	14.21 (12.88– 15.37)	15.33 (15.06– 15.65)	11.70 (10.66– 12.65)
Brazil	76.21 (75.81– 76.59)	65.73 (62.70– 68.46)	68.69 (68.28– 69.12)	60.58 (58.18– 62.74)	78.21 (77.63– 78.74)	67.38 (64.24– 70.09)	70.72 (70.02– 71.35)	62.35 (59.87– 64.54)	18.74 (18.38– 19.11)	14.23 (12.89– 15.40)	15.32 (15.05– 15.65)	11.70 (10.65– 12.65)
Paraguay	76.10 (75.16– 77.01)	65.86 (62.81– 68.66)	72.03 (71.11– 72.86)	63.47 (60.86– 65.87)	76.95 (75.54– 78.38)	66.44 (63.14– 69.64)	72.09 (70.20– 73.79)	63.54 (60.38– 66.19)	17.79 (17.02– 18.72)	13.48 (12.10– 14.92)	15.74 (14.84– 16.55)	12.01 (10.72– 13.22)
Southeast Asia, east Asia, and Oceania	73.95 (73.58– 74.33)	65.33 (62.87– 67.58)	68.52 (68.12– 68.95)	61.40 (59.27– 63.32)	78.38 (77.84– 78.93)	69.11 (66.39– 71.55)	71.90 (71.26– 72.50)	64.36 (62.09– 66.41)	18.48 (18.12– 18.87)	14.53 (13.45– 15.60)	14.95 (14.63– 15.29)	11.86 (10.97– 12.70)
East Asia	74.94 (74.46– 75.43)	66.54 (64.09– 68.73)	69.38 (68.86– 69.89)	62.44 (60.36– 64.35)	79.83 (79.19– 80.49)	70.71 (72.45– 73.12)	73.17 (72.45– 73.88)	65.74 (63.45– 67.78)	18.99 (18.51– 19.50)	15.07 (13.98– 16.18)	15.13 (14.73– 15.54)	12.09 (11.18– 12.92)
China	74.90 (74.41– 75.40)	66.53 (64.05– 68.73)	69.33 (68.82– 69.86)	62.40 (60.32– 64.31)	79.89 (79.24– 80.56)	70.78 (72.44– 73.23)	73.19 (72.44– 73.91)	65.75 (63.43– 67.77)	19.01 (18.52– 19.57)	15.11 (14.00– 16.22)	15.10 (14.68– 15.50)	12.07 (11.16– 12.90)
North Korea	72.55 (69.82– 75.12)	64.12 (60.50– 67.12)	67.53 (64.35– 71.01)	60.93 (57.56– 64.20)	74.89 (72.19– 77.52)	66.20 (62.62– 69.50)	68.94 (58.78– 71.93)	62.22 (58.78– 65.27)	16.59 (15.35– 17.90)	13.02 (11.60– 14.42)	13.94 (12.91– 15.25)	11.14 (10.00– 12.41)
Taiwan (province of China)	80.69 (80.57– 80.81)	70.76 (67.85– 73.30)	74.48 (74.38– 74.60)	66.78 (64.52– 68.76)	82.50 (81.18– 83.87)	72.18 (72.44– 75.14)	76.64 (75.07– 78.27)	68.54 (65.66– 71.16)	20.75 (19.81– 21.71)	16.11 (14.63– 17.56)	17.83 (17.04– 18.69)	14.14 (12.85– 15.40)
Southeast Asia	72.16 (71.46– 72.89)	62.99 (60.28– 65.42)	67.08 (66.24– 68.00)	59.39 (57.12– 61.51)	75.32 (74.30– 76.32)	65.77 (63.04– 68.45)	69.18 (67.96– 70.46)	61.37 (58.95– 63.67)	17.10 (16.64– 17.58)	13.05 (11.88– 14.14)	14.32 (13.85– 14.99)	11.02 (10.03– 12.01)
Cambodia	66.24 (65.34– 67.27)	57.67 (55.12– 60.05)	61.53 (60.48– 62.59)	54.41 (52.18– 56.48)	72.44 (70.47– 74.65)	63.20 (60.14– 66.32)	66.21 (64.31– 68.00)	58.76 (56.24– 61.37)	15.49 (14.72– 16.72)	11.80 (10.60– 13.12)	13.09 (12.36– 13.71)	10.05 (9.12– 11.01)
Indonesia	70.11 (69.16– 71.14)	61.40 (58.93– 63.80)	67.35 (66.18– 68.63)	59.63 (57.19– 61.78)	73.29 (71.23– 75.39)	64.06 (61.07– 66.97)	69.01 (66.64– 71.91)	61.21 (58.27– 64.28)	15.80 (15.03– 16.90)	11.97 (10.81– 13.15)	14.17 (13.25– 15.79)	10.88 (9.66– 12.36)
Laos	62.05 (60.33– 63.86)	54.26 (51.70– 56.65)	58.08 (56.59– 59.47)	51.31 (49.04– 53.42)	69.48 (66.71– 72.17)	60.75 (57.56– 63.91)	64.73 (62.17– 67.51)	57.20 (54.14– 60.31)	15.26 (13.95– 16.63)	11.60 (10.20– 13.00)	13.78 (12.78– 14.71)	10.49 (9.37– 11.60)
Malaysia	76.30 (76.22– 76.37)	66.76 (64.04– 69.16)	72.35 (72.27– 72.44)	64.08 (61.75– 66.23)	78.62 (77.72– 79.60)	68.60 (65.64– 71.42)	72.39 (70.75– 74.04)	64.21 (61.50– 66.73)	17.89 (17.46– 18.39)	13.71 (12.45– 14.86)	15.16 (14.29– 16.00)	11.79 (10.61– 12.90)
Maldives	78.22 (77.80– 78.65)	67.35 (64.31– 70.02)	75.73 (75.17– 76.25)	66.24 (63.45– 68.64)	82.26 (81.00– 83.52)	71.20 (68.08– 74.17)	78.06 (76.40– 79.63)	68.79 (65.73– 71.60)	20.80 (19.93– 21.72)	15.90 (14.43– 17.29)	18.29 (17.22– 19.27)	14.17 (12.75– 15.45)
Mauritius	75.67 (75.40– 75.92)	66.47 (63.85– 68.76)	69.17 (68.91– 69.42)	61.68 (59.50– 63.58)	77.70 (77.09– 78.26)	67.92 (65.14– 70.41)	71.40 (70.82– 72.07)	63.38 (60.98– 65.51)	17.89 (17.56– 18.24)	13.70 (12.52– 14.77)	14.77 (14.50– 15.10)	11.39 (10.39– 12.27)
Myanmar	66.02 (61.76– 70.84)	57.67 (53.57– 61.94)	60.74 (56.24– 61.94)	53.78 (49.53– 58.22)	71.20 (66.85– 75.28)	62.24 (58.07– 66.28)	64.92 (60.29– 69.96)	57.56 (53.22– 61.91)	15.05 (13.25– 17.14)	11.36 (9.68– 13.07)	12.90 (11.42– 15.11)	9.82 (8.39– 11.59)

(Table 2 continues on next page)

	2005, at birth				2015, at birth				2015, at age 65 years			
	Females		Males		Females		Males		Females		Males	
	Life expectancy	HALE										
(Continued from previous page)												
Philippines	73.03 (72.88– 73.18)	63.05 (60.16– 65.57)	65.43 (65.27– 65.58)	57.69 (55.43– 59.66)	74.41 (73.36– 75.31)	64.43 (61.55– 67.10)	67.71 (66.63– 68.76)	59.82 (57.37– 61.95)	16.55 (16.00– 17.05)	12.50 (11.34– 13.58)	13.19 (12.72– 13.67)	10.03 (9.09– 10.89)
Sri Lanka	77.97 (77.72– 78.23)	68.07 (65.20– 70.58)	70.10 (69.85– 70.36)	62.08 (59.74– 64.13)	81.19 (79.06– 83.24)	70.84 (67.31– 73.92)	74.05 (71.09– 76.93)	65.47 (61.81– 68.87)	19.88 (18.47– 21.33)	15.31 (13.60– 16.89)	16.64 (15.29– 18.11)	12.85 (11.28– 14.39)
Seychelles	75.88 (75.41– 76.46)	66.68 (64.02– 69.12)	68.23 (67.63– 68.81)	60.74 (58.52– 62.75)	77.93 (76.78– 78.97)	68.29 (65.52– 70.95)	70.55 (68.82– 72.14)	62.77 (60.08– 65.25)	17.98 (17.48– 18.44)	13.81 (12.63– 14.94)	14.50 (13.82– 15.26)	11.13 (10.05– 12.20)
Thailand	76.50 (75.95– 77.06)	66.91 (64.21– 69.39)	69.77 (68.82– 70.76)	61.78 (59.27– 64.01)	78.98 (77.47– 80.46)	69.11 (66.04– 72.23)	70.81 (68.74– 73.24)	62.92 (60.29– 65.67)	18.77 (18.08– 19.63)	14.48 (13.16– 15.80)	14.97 (14.25– 16.20)	11.59 (10.51– 12.79)
Timor-Leste	68.04 (66.22– 69.89)	59.73 (57.05– 62.44)	66.97 (65.35– 68.64)	58.74 (55.92– 61.30)	72.96 (70.01– 75.75)	64.02 (60.49– 67.35)	71.98 (68.85– 75.12)	63.08 (59.32– 66.79)	16.40 (14.96– 17.76)	12.60 (11.07– 14.05)	16.61 (15.28– 18.13)	12.61 (11.10– 14.17)
Vietnam	76.96 (74.35– 79.23)	67.23 (63.76– 70.40)	69.45 (66.39– 72.95)	61.89 (58.50– 65.51)	79.38 (77.02– 82.25)	69.76 (66.41– 73.20)	71.31 (67.83– 74.84)	63.80 (60.25– 67.14)	18.83 (17.64– 20.57)	14.65 (13.11– 16.41)	14.75 (13.69– 16.34)	11.58 (10.33– 13.06)
Oceania	60.94 (56.86– 65.01)	52.15 (48.22– 56.06)	59.03 (55.23– 63.02)	52.11 (48.32– 55.82)	63.72 (58.96– 67.73)	54.48 (49.99– 58.41)	61.42 (57.02– 65.32)	54.24 (50.11– 57.86)	11.91 (10.33– 13.38)	8.73 (7.39– 10.01)	11.74 (10.65– 12.99)	8.86 (7.77– 10.03)
American Samoa	74.63 (72.98– 75.91)	64.66 (61.63– 67.36)	69.55 (67.93– 71.11)	61.14 (58.40– 63.76)	74.85 (72.39– 77.07)	64.86 (61.72– 68.10)	70.87 (68.59– 73.17)	62.33 (59.32– 65.44)	16.17 (14.98– 17.31)	12.17 (10.85– 13.56)	14.46 (13.64– 15.44)	10.94 (9.80– 12.13)
Federated States of Micronesia	69.93 (64.32– 74.51)	61.17 (55.63– 65.47)	65.36 (60.00– 70.18)	58.15 (52.88– 62.29)	71.54 (65.30– 75.82)	62.52 (56.78– 66.79)	66.52 (60.42– 70.80)	59.17 (53.52– 63.30)	14.60 (11.87– 16.79)	11.12 (10.75– 13.02)	12.96 (10.83– 14.41)	9.94 (8.07– 11.30)
Fiji	67.13 (65.42– 68.78)	58.07 (55.19– 60.77)	62.69 (60.85– 64.52)	55.18 (52.48– 57.81)	68.13 (65.45– 70.87)	58.97 (60.45– 62.18)	63.69 (60.77– 66.48)	56.19 (55.62– 59.16)	13.44 (12.28– 14.61)	10.05 (8.82– 11.33)	11.87 (10.99– 12.73)	8.97 (7.99– 10.00)
Guam	76.52 (75.37– 77.64)	65.90 (62.55– 68.94)	70.07 (68.71– 71.26)	62.30 (60.02– 64.56)	75.31 (73.53– 77.29)	64.87 (61.61– 67.99)	68.49 (66.20– 70.79)	60.87 (58.19– 63.57)	16.69 (15.95– 17.72)	12.55 (11.35– 13.79)	13.65 (12.85– 14.37)	10.52 (9.51– 11.52)
Kiribati	64.94 (62.65– 67.10)	56.02 (52.94– 58.91)	56.94 (54.59– 59.28)	50.14 (47.27– 52.89)	67.44 (64.55– 70.21)	58.34 (54.91– 61.62)	58.85 (54.91– 61.78)	52.04 (48.74– 55.23)	13.39 (12.31– 14.42)	10.04 (8.86– 11.15)	10.74 (9.94– 11.52)	8.13 (7.19– 9.08)
Marshall Islands	66.83 (64.59– 68.99)	57.26 (54.03– 60.39)	62.73 (60.79– 64.89)	54.83 (52.07– 57.66)	69.50 (66.96– 71.91)	59.42 (55.77– 62.90)	64.57 (61.96– 67.29)	56.37 (53.19– 59.53)	13.79 (12.65– 14.83)	10.12 (8.77– 11.43)	12.42 (11.59– 13.18)	9.23 (8.14– 10.25)
Northern Mariana Islands	77.39 (76.65– 78.14)	67.32 (64.34– 69.95)	74.55 (73.60– 75.36)	65.91 (63.22– 68.16)	78.47 (77.15– 79.64)	68.26 (65.13– 71.21)	74.17 (72.71– 75.49)	65.64 (62.87– 68.11)	18.21 (17.55– 18.80)	13.88 (12.59– 15.11)	15.92 (15.26– 16.49)	12.28 (11.15– 13.36)
Papua New Guinea	58.54 (53.42– 63.88)	49.82 (45.14– 54.61)	57.25 (52.39– 62.71)	50.55 (45.95– 55.57)	61.85 (55.97– 67.04)	52.62 (47.33– 57.26)	60.13 (54.55– 57.26)	53.11 (47.99– 57.67)	11.07 (10.41– 13.12)	8.02 (6.41– 9.61)	11.34 (9.73– 13.22)	8.52 (7.14– 10.11)
Samoa	73.70 (71.66– 75.83)	64.10 (60.84– 67.21)	70.00 (67.91– 71.97)	61.81 (58.87– 64.79)	74.22 (71.79– 76.62)	64.45 (60.99– 67.91)	72.23 (69.74– 74.55)	63.58 (60.54– 66.60)	15.96 (14.76– 17.15)	12.06 (10.65– 13.56)	15.02 (14.00– 16.06)	11.45 (10.19– 12.72)
Solomon Islands	62.69 (57.49– 68.21)	54.94 (50.49– 59.82)	60.93 (56.04– 65.96)	54.44 (50.20– 58.93)	65.12 (59.19– 70.81)	56.98 (51.74– 61.79)	62.86 (56.96– 68.13)	56.07 (51.03– 60.69)	11.78 (9.54– 14.16)	8.95 (7.07– 10.81)	11.83 (9.87– 13.36)	9.10 (7.53– 10.52)
Tonga	72.70 (71.27– 73.99)	62.79 (59.60– 65.53)	67.29 (65.66– 68.82)	59.53 (56.92– 61.95)	74.68 (72.19– 77.07)	64.43 (60.83– 67.76)	68.40 (65.44– 71.10)	60.50 (57.30– 63.51)	16.40 (15.33– 17.46)	12.25 (10.86– 13.62)	13.64 (12.69– 14.66)	10.39 (9.31– 11.55)
Vanuatu	64.21 (59.00– 69.19)	56.64 (51.85– 61.42)	61.37 (56.42– 66.39)	54.95 (50.36– 59.62)	66.64 (61.23– 70.86)	58.58 (53.61– 62.91)	63.22 (58.12– 67.35)	56.41 (51.72– 60.55)	12.72 (10.52– 14.57)	9.79 (7.92– 11.44)	12.03 (10.43– 13.29)	9.32 (7.90– 10.69)

(Table 2 continues on next page)

	2005, at birth				2015, at birth				2015, at age 65 years			
	Females		Males		Females		Males		Females		Males	
	Life expectancy	HALE										
(Continued from previous page)												
North Africa and Middle East	72·55 (71·85– 73·21)	62·33 (59·42– 65·02)	68·64 (67·98– 69·34)	59·98 (57·44– 62·33)	74·94 (74·12– 75·69)	64·29 (61·31– 67·14)	70·07 (69·16– 70·94)	61·16 (58·55– 63·62)	17·58 (17·28– 17·87)	13·07 (11·80– 14·25)	15·37 (15·08– 15·65)	11·55 (10·45– 12·57)
Afghanistan	51·40 (47·39– 55·64)	44·45 (40·62– 48·10)	52·67 (48·93– 56·74)	45·55 (42·13– 49·21)	54·46 (50·19– 58·82)	47·11 (43·28– 51·24)	53·34 (49·22– 57·70)	46·27 (42·42– 50·19)	9·41 (8·21– 10·93)	6·91 (5·82– 8·22)	10·57 (9·49– 12·37)	7·68 (6·52– 9·09)
Algeria	76·07 (74·93– 77·12)	65·55 (62·46– 68·52)	73·51 (72·29– 74·79)	64·39 (61·45– 67·02)	77·44 (76·01– 78·52)	66·66 (63·44– 69·65)	75·37 (74·13– 76·79)	65·91 (63·15– 68·58)	18·29 (17·47– 18·76)	13·73 (12·34– 15·01)	17·48 (16·90– 18·23)	13·26 (11·99– 14·47)
Bahrain	78·06 (77·24– 78·93)	66·10 (62·55– 69·30)	74·63 (73·80– 75·59)	64·92 (62·04– 67·48)	80·34 (78·92– 81·83)	67·98 (64·36– 71·35)	78·21 (76·45– 79·99)	67·89 (64·67– 71·12)	19·23 (18·60– 20·06)	14·00 (12·51– 15·46)	17·76 (16·79– 18·90)	13·22 (11·71– 14·75)
Egypt	72·64 (72·38– 72·87)	62·41 (59·38– 65·08)	67·69 (67·45– 67·94)	59·42 (57·08– 61·59)	74·45 (73·88– 74·95)	63·84 (60·71– 66·58)	68·71 (68·13– 69·28)	60·23 (57·62– 62·46)	16·42 (16·23– 16·61)	12·16 (10·97– 13·28)	13·50 (13·34– 13·66)	10·12 (9·16– 11·03)
Iran	74·58 (72·51– 76·50)	64·28 (61·12– 67·51)	69·64 (67·24– 71·98)	60·98 (57·86– 64·21)	77·98 (75·87– 80·02)	67·13 (63·56– 70·62)	71·85 (69·18– 74·18)	62·99 (59·82– 66·40)	17·85 (16·78– 19·04)	13·34 (11·86– 14·92)	15·06 (14·08– 16·17)	11·40 (10·11– 12·80)
Iraq	68·24 (65·65– 70·65)	58·57 (55·18– 61·60)	63·26 (59·95– 66·53)	54·71 (51·15– 58·09)	70·66 (67·52– 73·64)	60·53 (56·47– 63·92)	64·33 (60·79– 68·46)	55·65 (51·90– 59·50)	14·80 (13·47– 16·11)	10·88 (9·40– 12·25)	13·05 (11·96– 15·04)	9·55 (8·38– 10·94)
Jordan	74·44 (73·52– 75·33)	63·69 (60·58– 66·46)	74·66 (73·91– 75·44)	65·14 (62·30– 67·75)	80·69 (79·65– 81·78)	68·76 (75·09– 72·06)	76·46 (73·79– 77·61)	66·68 (63·79– 69·52)	19·84 (19·18– 20·58)	14·67 (13·09– 16·13)	17·46 (16·64– 18·17)	13·12 (11·76– 14·44)
Kuwait	80·19 (79·91– 80·52)	68·91 (65·71– 71·77)	76·27 (75·99– 76·54)	66·90 (64·25– 69·36)	81·55 (80·47– 82·71)	70·19 (66·81– 73·29)	79·43 (77·94– 80·67)	69·56 (66·56– 72·40)	19·18 (18·38– 20·06)	14·41 (12·88– 15·84)	18·94 (18·09– 19·69)	14·40 (12·97– 15·80)
Lebanon	77·64 (75·78– 79·42)	65·80 (62·25– 69·26)	77·03 (74·16– 79·60)	65·45 (61·58– 69·00)	79·87 (77·76– 82·32)	68·01 (64·42– 71·62)	77·90 (75·08– 80·86)	66·81 (62·93– 70·47)	19·01 (18·09– 20·31)	14·11 (12·60– 15·68)	17·80 (16·14– 19·72)	13·07 (11·28– 15·01)
Libya	76·92 (76·07– 77·62)	66·28 (63·11– 69·04)	73·91 (73·06– 74·78)	64·74 (61·95– 67·21)	76·86 (75·26– 78·22)	66·09 (62·84– 69·25)	71·41 (68·93– 73·78)	62·45 (59·39– 65·54)	17·84 (16·99– 18·55)	13·34 (11·96– 14·72)	15·90 (14·76– 16·92)	11·99 (10·61– 13·33)
Morocco	73·80 (70·89– 76·16)	63·24 (59·71– 66·72)	71·62 (68·72– 74·08)	61·88 (58·64– 65·13)	75·83 (72·69– 79·05)	64·92 (61·12– 68·62)	73·32 (69·45– 76·58)	63·37 (59·35– 66·89)	17·20 (15·61– 19·00)	12·70 (11·04– 14·44)	16·39 (14·35– 18·01)	12·13 (10·31– 13·70)
Palestine	77·34 (76·55– 78·10)	66·84 (63·82– 69·70)	72·06 (70·51– 73·06)	62·49 (59·81– 65·14)	77·90 (75·78– 79·78)	67·44 (63·98– 70·52)	71·60 (68·64– 74·52)	62·50 (59·06– 65·67)	18·17 (17·04– 19·22)	13·84 (12·39– 15·21)	15·01 (13·92– 16·52)	11·35 (9·94– 12·77)
Oman	78·38 (77·34– 79·32)	66·81 (63·47– 69·91)	74·07 (72·92– 75·18)	64·15 (61·29– 66·85)	79·64 (78·29– 81·14)	67·75 (74·05– 71·05)	75·71 (76·93– 77·69)	65·38 (64·92– 68·65)	18·79 (18·11– 19·71)	13·71 (12·25– 15·17)	16·32 (15·48– 17·55)	12·03 (10·70– 13·57)
Qatar	78·31 (77·49– 79·11)	67·01 (63·71– 70·07)	75·24 (73·63– 76·87)	65·43 (62·30– 68·46)	81·20 (79·43– 82·88)	69·31 (65·82– 72·71)	79·11 (76·93– 81·25)	68·57 (64·92– 71·83)	19·76 (18·65– 20·98)	14·58 (12·92– 16·27)	18·59 (12·07– 19·91)	13·80 (12·07– 15·51)
Saudi Arabia	80·00 (79·43– 80·55)	68·99 (65·85– 71·89)	75·95 (75·29– 76·58)	66·78 (64·09– 69·25)	82·21 (81·29– 83·09)	70·87 (67·53– 74·08)	77·45 (76·34– 78·53)	68·15 (65·33– 70·73)	20·68 (19·98– 21·46)	15·72 (14·25– 17·18)	17·77 (16·98– 18·51)	13·67 (12·41– 14·90)
Sudan	66·07 (62·29– 70·01)	56·47 (52·50– 60·70)	63·09 (59·34– 67·21)	54·72 (50·75– 58·67)	69·27 (65·38– 73·59)	59·38 (55·50– 63·72)	65·81 (61·55– 69·91)	57·16 (53·08– 61·35)	15·03 (13·32– 17·27)	11·20 (9·59– 13·17)	14·27 (12·39– 15·93)	10·56 (8·92– 12·10)
Syria	77·37 (76·74– 77·97)	67·20 (64·25– 69·78)	73·85 (73·09– 74·68)	65·31 (62·81– 67·63)	74·81 (72·12– 77·75)	63·82 (59·86– 67·73)	62·58 (56·48– 70·19)	54·17 (48·63– 61·05)	17·98 (17·37– 18·82)	13·46 (12·11– 14·74)	15·47 (14·56– 16·66)	11·74 (10·46– 12·98)
Tunisia	78·95 (77·45– 80·46)	68·16 (64·92– 71·20)	73·66 (71·83– 75·41)	64·38 (61·36– 67·24)	80·73 (78·82– 82·71)	69·61 (66·03– 72·89)	74·63 (72·03– 77·02)	65·26 (61·88– 68·39)	19·77 (18·50– 21·22)	14·95 (13·25– 16·56)	16·18 (14·85– 17·69)	12·24 (10·67– 13·82)

(Table 2 continues on next page)

	2005, at birth				2015, at birth				2015, at age 65 years			
	Females		Males		Females		Males		Females		Males	
	Life expectancy	HALE										
(Continued from previous page)												
Turkey	79.52 (78.90– 80.10)	68.04 (64.90– 70.99)	72.91 (72.25– 73.57)	64.01 (61.39– 66.35)	82.37 (81.82– 82.94)	70.49 (66.94– 73.59)	75.90 (75.25– 76.55)	66.61 (63.86– 69.04)	21.21 (20.91– 21.55)	15.75 (14.19– 17.16)	17.35 (16.89– 17.67)	13.17 (11.92– 14.27)
United Arab Emirates	77.32 (76.47– 78.00)	66.22 (63.07– 69.16)	74.16 (73.26– 74.93)	64.47 (61.53– 67.14)	78.02 (75.54– 79.89)	67.00 (63.40– 70.34)	74.52 (71.82– 77.08)	65.04 (61.38– 68.31)	17.99 (16.84– 18.95)	13.46 (11.93– 14.80)	15.92 (14.70– 17.04)	11.98 (10.50– 13.33)
Yemen	64.72 (59.58– 70.44)	56.24 (51.57– 61.26)	64.41 (59.19– 69.89)	56.78 (52.15– 61.46)	66.57 (60.83– 71.63)	57.71 (52.65– 62.10)	65.39 (59.70– 70.23)	57.40 (52.34– 61.68)	13.55 (11.14– 16.14)	10.28 (8.32– 16.14)	14.03 (11.65– 12.21)	10.71 (8.75– 12.50)
South Asia	65.22 (64.66– 65.79)	55.61 (52.88– 58.11)	62.43 (61.92– 62.97)	54.26 (51.89– 56.37)	69.56 (68.83– 70.27)	59.52 (56.53– 62.18)	65.42 (64.74– 66.10)	57.15 (54.76– 59.35)	15.10 (14.74– 15.47)	11.19 (10.11– 12.26)	13.39 (13.11– 13.69)	10.04 (9.15– 10.90)
Bangladesh	68.70 (67.58– 69.89)	58.62 (55.71– 61.43)	65.67 (64.60– 66.76)	57.34 (54.71– 59.68)	72.52 (70.56– 74.46)	62.18 (58.79– 65.30)	68.52 (66.50– 70.35)	60.06 (57.26– 62.74)	16.16 (14.93– 17.32)	12.13 (10.66– 13.46)	14.58 (13.76– 15.33)	11.12 (10.01– 12.19)
Bhutan	70.15 (67.77– 72.32)	59.82 (56.53– 63.05)	68.68 (66.42– 70.67)	59.58 (56.50– 62.30)	74.26 (71.54– 76.54)	63.08 (59.42– 66.49)	71.50 (69.05– 74.20)	61.95 (58.52– 65.20)	17.80 (16.35– 19.13)	13.11 (11.52– 14.60)	16.35 (15.06– 17.75)	12.22 (10.73– 13.75)
India	65.03 (64.41– 65.66)	55.38 (52.60– 57.89)	62.22 (61.65– 62.83)	54.01 (51.64– 56.15)	69.53 (68.68– 70.43)	59.44 (56.46– 62.17)	65.24 (64.46– 66.00)	56.97 (54.53– 59.20)	15.06 (14.65– 15.51)	11.14 (10.03– 12.17)	13.21 (12.89– 13.53)	9.89 (9.00– 10.75)
Nepal	68.83 (66.99– 70.57)	59.54 (56.41– 62.32)	65.96 (64.42– 67.66)	57.63 (54.90– 60.03)	71.23 (68.75– 73.92)	61.58 (58.13– 64.95)	68.09 (65.20– 70.91)	59.53 (56.39– 62.74)	15.56 (14.22– 17.16)	11.81 (10.30– 13.35)	14.76 (13.48– 16.01)	11.25 (9.97– 12.59)
Pakistan	63.24 (61.42– 65.05)	54.31 (51.48– 57.18)	61.29 (59.54– 63.19)	53.51 (50.86– 55.94)	67.32 (65.22– 69.82)	57.93 (54.76– 61.01)	64.54 (62.22– 66.49)	56.40 (53.59– 59.28)	14.39 (13.34– 15.78)	10.73 (9.41– 12.06)	13.59 (12.72– 14.32)	10.21 (9.14– 11.34)
Sub-Saharan Africa	55.92 (55.05– 56.74)	47.94 (45.32– 50.16)	54.26 (53.46– 55.10)	47.13 (44.90– 49.12)	63.61 (61.83– 65.17)	54.63 (51.52– 57.44)	60.02 (58.35– 61.58)	52.35 (49.72– 54.90)	14.28 (13.44– 15.06)	10.57 (9.31– 11.69)	13.33 (12.62– 13.99)	9.96 (8.86– 10.99)
Southern sub-Saharan Africa	51.80 (50.49– 53.23)	44.71 (42.42– 46.86)	49.85 (48.74– 51.12)	43.62 (41.66– 45.44)	63.21 (61.67– 64.69)	54.06 (51.28– 56.71)	57.51 (56.00– 58.95)	50.14 (47.75– 52.40)	14.81 (14.01– 15.56)	10.80 (9.59– 11.98)	12.35 (11.68– 13.03)	9.05 (8.05– 9.98)
Botswana	50.78 (39.38– 64.33)	44.97 (34.78– 54.50)	47.92 (38.74– 60.22)	42.81 (34.01– 51.74)	62.01 (47.70– 72.30)	53.95 (41.57– 61.73)	55.78 (43.89– 61.73)	49.39 (39.01– 66.62)	13.82 (7.51– 57.39)	10.57 (7.51– 18.54)	12.01 (7.31– 13.91)	9.10 (5.40– 12.22)
Lesotho	44.77 (41.20– 48.58)	38.92 (35.56– 42.30)	41.21 (38.65– 44.59)	36.45 (34.07– 39.63)	50.41 (43.61– 58.50)	43.83 (37.87– 49.81)	44.11 (38.61– 51.77)	39.13 (34.27– 44.74)	11.49 (8.12– 15.20)	8.57 (5.97– 11.17)	9.26 (7.16– 12.77)	6.87 (5.22– 9.42)
Namibia	55.05 (51.33– 59.06)	47.65 (43.90– 51.40)	51.07 (48.07– 54.29)	44.91 (41.85– 48.01)	68.44 (61.62– 73.04)	58.70 (52.80– 63.47)	59.98 (53.61– 66.28)	52.56 (46.87– 58.13)	16.51 (12.83– 18.64)	12.31 (9.39– 14.52)	13.01 (10.40– 15.97)	9.79 (7.69– 12.19)
South Africa	53.62 (52.18– 55.38)	46.19 (43.83– 48.49)	51.67 (50.36– 53.14)	45.17 (43.09– 47.15)	63.99 (62.55– 65.18)	54.63 (51.81– 57.23)	58.64 (57.32– 59.91)	51.07 (48.74– 53.20)	14.97 (14.18– 15.69)	10.88 (9.65– 12.06)	12.56 (11.93– 13.21)	9.15 (8.16– 10.12)
Swaziland	43.17 (39.24– 47.45)	37.24 (33.86– 40.73)	41.68 (38.42– 45.84)	36.52 (33.44– 39.83)	54.87 (46.71– 63.97)	47.12 (40.22– 53.87)	48.96 (42.27– 58.26)	43.01 (37.10– 49.82)	12.24 (8.24– 16.65)	9.06 (5.84– 12.09)	10.36 (7.55– 14.68)	7.68 (5.39– 10.79)
Zimbabwe	45.56 (42.36– 49.34)	39.75 (36.90– 42.77)	44.61 (41.95– 48.39)	39.24 (36.79– 42.17)	62.54 (56.27– 68.35)	54.29 (48.68– 59.44)	56.28 (51.09– 62.63)	49.52 (44.79– 54.76)	14.81 (11.38– 18.26)	11.23 (8.38– 13.90)	12.20 (10.09– 15.22)	9.24 (7.37– 11.68)
Western sub-Saharan Africa	57.51 (56.07– 58.82)	49.35 (46.72– 51.71)	55.80 (54.46– 57.19)	48.44 (46.03– 50.65)	64.15 (60.80– 66.47)	55.25 (51.81– 58.49)	60.97 (58.24– 62.97)	53.19 (50.17– 55.91)	14.98 (13.35– 16.06)	11.21 (9.74– 12.54)	14.16 (13.01– 15.00)	10.62 (9.32– 11.78)
Benin	60.97 (56.43– 65.32)	52.53 (48.15– 56.53)	56.21 (52.01– 61.29)	49.70 (45.45– 53.99)	64.84 (55.71– 72.41)	56.44 (51.34– 62.89)	59.88 (58.24– 68.32)	53.59 (50.17– 60.42)	13.93 (12.54– 17.93)	10.67 (9.47– 13.80)	12.40 (9.11– 15.93)	9.66 (7.11– 12.34)

(Table 2 continues on next page)

	2005, at birth				2015, at birth				2015, at age 65 years			
	Females		Males		Females		Males		Females		Males	
	Life expectancy	HALE	Life expectancy	HALE								
(Continued from previous page)												
Burkina Faso	56·63 (53·42– 59·73)	48·94 (45·56– 52·15)	55·04 (52·19– 58·27)	48·19 (44·89– 51·56)	61·85 (54·32– 68·96)	54·25 (47·62– 60·31)	60·01 (52·54– 66·03)	53·28 (47·34– 58·69)	13·59 (10·13– 17·19)	10·51 (7·72– 13·28)	13·54 (10·32– 16·35)	10·47 (7·98– 12·83)
Cameroon	55·92 (52·80– 59·52)	48·21 (44·66– 51·56)	53·84 (50·74– 57·25)	46·89 (43·74– 50·19)	61·20 (53·95– 68·55)	53·35 (46·88– 59·61)	57·59 (50·44– 64·22)	50·85 (44·49– 56·32)	14·06 (10·39– 17·58)	10·68 (7·68– 13·40)	12·99 (9·83– 16·07)	9·86 (7·25– 12·30)
Cape Verde	76·02 (73·41– 76·91)	65·63 (62·25– 68·60)	63·22 (59·21– 68·01)	56·01 (52·27– 60·21)	76·55 (74·33– 77·59)	66·26 (62·96– 69·24)	69·13 (66·18– 71·78)	60·99 (57·64– 64·19)	18·09 (16·79– 18·64)	13·68 (12·17– 15·01)	14·85 (13·40– 16·22)	11·31 (9·78– 12·82)
Chad	54·49 (49·28– 60·02)	47·07 (42·17– 51·79)	51·52 (46·42– 57·50)	44·78 (40·30– 49·49)	58·88 (49·37– 67·61)	51·31 (43·16– 58·31)	55·92 (47·52– 63·60)	48·99 (41·63– 55·16)	13·57 (9·15– 18·17)	10·39 (6·83– 13·74)	12·82 (9·30– 16·41)	9·68 (6·76– 12·31)
Côte d'Ivoire	54·81 (51·77– 58·19)	47·31 (44·08– 50·42)	52·17 (49·40– 55·26)	45·74 (42·79– 48·69)	61·41 (55·01– 68·52)	53·25 (47·55– 58·86)	57·21 (49·99– 64·50)	50·62 (44·57– 56·49)	13·45 (10·36– 17·14)	10·18 (7·63– 12·89)	12·52 (9·58– 15·76)	9·58 (7·19– 12·09)
The Gambia	66·15 (63·52– 69·30)	57·47 (53·89– 60·95)	63·77 (60·62– 66·93)	56·24 (53·02– 59·47)	69·63 (64·33– 74·55)	60·60 (55·53– 65·49)	66·65 (61·40– 70·72)	58·85 (53·84– 63·05)	15·20 (12·74– 17·85)	11·66 (9·40– 13·93)	14·44 (12·06– 16·15)	11·12 (9·08– 12·84)
Ghana	62·22 (58·39– 66·35)	54·17 (50·38– 58·28)	59·44 (55·06– 63·57)	52·58 (48·64– 56·59)	67·55 (59·65– 74·25)	59·26 (52·16– 65·27)	63·06 (54·99– 69·81)	56·43 (49·27– 62·23)	14·65 (11·01– 18·42)	11·43 (8·29– 14·33)	13·26 (10·33– 16·14)	10·37 (7·79– 12·80)
Guinea	56·87 (54·19– 59·83)	49·13 (46·04– 52·16)	55·17 (52·35– 58·20)	48·39 (45·37– 51·39)	60·49 (54·52– 67·09)	52·65 (47·07– 58·36)	57·81 (50·80– 64·49)	51·10 (45·32– 56·48)	13·11 (10·29– 16·46)	10·04 (7·59– 12·67)	12·74 (9·77– 15·74)	9·76 (7·38– 12·08)
Guinea-Bissau	53·45 (41·87– 65·17)	47·62 (37·00– 56·49)	52·32 (41·45– 61·55)	46·95 (36·92– 54·28)	56·44 (42·35– 67·69)	50·29 (37·66– 58·94)	54·86 (41·87– 63·97)	49·30 (37·55– 56·40)	12·49 (37·14)	9·86 (7·38– 13·75)	12·60 (5·56– 16·67)	9·81 (7·97– 12·82)
Liberia	58·41 (55·78– 61·50)	49·34 (45·91– 52·69)	58·42 (55·80– 61·11)	49·59 (46·21– 52·76)	63·41 (57·41– 69·44)	54·28 (49·06– 59·54)	63·08 (57·54– 67·67)	54·32 (49·25– 58·93)	13·46 (11·01– 16·61)	9·89 (7·64– 12·38)	14·11 (11·59– 16·32)	10·32 (8·27– 12·33)
Mali	57·10 (54·19– 59·62)	48·99 (45·63– 52·25)	56·70 (54·17– 59·25)	49·03 (45·80– 51·97)	60·44 (54·41– 66·68)	52·48 (46·95– 57·55)	60·04 (54·18– 64·35)	52·64 (47·53– 56·71)	13·86 (10·84– 16·93)	10·51 (8·06– 12·88)	14·63 (12·17– 16·71)	11·06 (8·94– 12·93)
Mauritania	66·08 (63·12– 68·95)	56·59 (53·00– 60·00)	66·76 (63·89– 69·70)	57·74 (54·26– 60·96)	69·81 (63·59– 75·57)	60·13 (54·58– 65·16)	69·83 (63·66– 75·84)	60·72 (55·13– 65·85)	15·43 (12·47– 18·74)	11·57 (9·08– 13·97)	16·23 (13·54– 19·71)	12·20 (9·82– 14·85)
Niger	56·69 (53·93– 59·39)	49·23 (46·04– 52·19)	55·23 (52·52– 57·72)	48·42 (45·56– 51·12)	62·47 (56·63– 68·18)	54·62 (49·53– 59·73)	59·58 (52·81– 65·38)	52·71 (46·88– 57·41)	14·18 (11·30– 17·09)	10·82 (8·50– 13·07)	13·37 (10·41– 15·90)	10·20 (7·88– 12·23)
Nigeria	57·27 (54·37– 59·89)	48·85 (45·55– 51·94)	56·32 (53·71– 59·05)	48·49 (45·46– 51·58)	66·33 (59·47– 70·32)	56·70 (51·20– 61·03)	62·95 (57·35– 65·78)	54·38 (49·69– 57·95)	16·69 (13·14– 18·76)	12·42 (9·77– 14·48)	15·56 (13·16– 16·63)	11·54 (9·51– 13·09)
São Tomé and Príncipe	66·23 (62·95– 69·55)	57·52 (53·96– 60·98)	64·70 (61·02– 68·11)	56·82 (53·12– 60·37)	68·81 (61·31– 75·85)	59·88 (53·33– 66·12)	67·19 (59·36– 74·84)	59·20 (52·56– 65·53)	14·08 (10·84– 17·67)	10·72 (8·12– 13·39)	14·05 (11·23– 17·55)	10·75 (8·30– 13·43)
Senegal	63·34 (58·73– 68·24)	54·93 (50·49– 59·50)	60·96 (55·78– 65·54)	53·85 (49·19– 57·99)	66·70 (58·22– 74·46)	58·42 (50·73– 64·76)	63·84 (54·36– 70·63)	56·97 (48·79– 63·01)	13·78 (10·12– 17·89)	10·62 (7·54– 13·61)	13·27 (9·98– 16·37)	10·37 (7·55– 12·81)
Sierra Leone	53·25 (51·13– 55·51)	45·84 (43·07– 48·53)	51·31 (49·15– 53·49)	44·43 (41·65– 46·90)	57·73 (51·98– 63·30)	50·57 (45·66– 55·39)	56·49 (50·98– 62·48)	49·86 (44·73– 54·86)	12·49 (11·46– 15·38)	9·48 (8·12– 11·79)	12·90 (10·28– 15·60)	9·74 (7·58– 11·87)
Togo	59·02 (55·78– 62·50)	51·21 (47·63– 54·69)	54·59 (51·23– 58·28)	48·11 (44·83– 51·43)	64·61 (58·52– 70·49)	56·48 (50·97– 61·52)	58·47 (52·25– 65·69)	52·02 (46·39– 57·51)	14·56 (11·46– 17·65)	11·20 (8·60– 13·65)	12·32 (9·91– 15·47)	9·53 (7·35– 11·85)
Eastern sub-Saharan Africa	56·08 (54·83– 57·34)	48·45 (45·86– 50·76)	54·53 (53·31– 55·72)	47·61 (45·40– 49·71)	64·32 (61·38– 67·07)	55·84 (52·48– 58·99)	60·50 (57·72– 63·16)	53·13 (50·12– 56·22)	13·98 (12·61– 15·37)	10·53 (9·15– 11·90)	13·06 (11·85– 14·21)	9·88 (8·61– 11·14)

(Table 2 continues on next page)

	2005, at birth				2015, at birth				2015, at age 65 years			
	Females		Males		Females		Males		Females		Males	
	Life expectancy	HALE	Life expectancy	HALE	Life expectancy	HALE						
(Continued from previous page)												
Burundi	55·27 (52·28– 58·06)	48·44 (45·16– 51·56)	54·16 (50·88– 57·52)	47·83 (44·43– 51·02)	62·69 (54·73– 70·23)	55·80 (48·75– 62·27)	60·10 (53·06– 67·68)	53·82 (47·80– 59·82)	12·82 (9·59– 16·50)	10·08 (7·37– 12·88)	12·63 (9·89– 15·75)	9·88 (7·68– 12·18)
Comoros	66·70 (63·68– 69·99)	57·99 (54·66– 61·50)	65·33 (62·26– 68·60)	57·09 (53·78– 60·39)	69·29 (63·40– 75·47)	60·62 (54·98– 65·59)	66·88 (59·97– 73·41)	58·93 (52·88– 64·41)	14·53 (11·92– 17·65)	11·10 (8·78– 13·54)	14·20 (11·45– 17·21)	10·82 (8·68– 13·10)
Djibouti	63·91 (53·86– 71·72)	55·49 (47·06– 61·84)	57·84 (47·80– 66·54)	51·35 (42·10– 58·41)	66·98 (56·90– 74·05)	58·00 (50·10– 64·10)	60·88 (50·23– 69·17)	53·93 (44·93– 60·78)	14·82 (10·34– 18·71)	11·17 (7·77– 13·99)	12·68 (9·03– 16·33)	9·67 (6·85– 12·37)
Eritrea	60·25 (51·43– 69·82)	52·69 (45·09– 59·93)	59·80 (49·75– 67·61)	52·23 (44·29– 58·27)	61·27 (51·78– 71·86)	54·02 (45·38– 61·97)	60·30 (50·84– 69·23)	53·13 (44·87– 60·06)	11·72 (8·22– 16·59)	9·11 (6·24– 12·36)	12·28 (9·13– 15·98)	9·37 (6·69– 12·12)
Ethiopia	56·14 (53·45– 58·72)	49·32 (46·48– 52·07)	56·04 (53·51– 58·96)	49·30 (46·65– 52·30)	66·80 (59·33– 73·42)	58·77 (52·50– 64·20)	63·57 (55·62– 69·63)	56·28 (49·52– 61·63)	14·27 (10·72– 17·84)	11·00 (8·27– 13·63)	13·56 (10·54– 16·34)	10·42 (7·90– 12·58)
Kenya	58·53 (57·14– 59·88)	49·67 (46·13– 52·44)	55·96 (54·64– 57·25)	48·76 (46·34– 50·93)	67·57 (65·77– 69·31)	57·66 (53·64– 61·01)	62·82 (61·09– 64·62)	54·92 (52·29– 57·47)	15·31 (14·17– 16·53)	11·45 (9·89– 12·97)	13·87 (13·01– 14·77)	10·59 (9·37– 11·68)
Madagascar	62·33 (59·40– 65·25)	53·64 (50·13– 57·00)	60·13 (56·80– 62·98)	52·03 (48·41– 55·18)	65·04 (57·77– 72·35)	56·75 (50·52– 62·59)	61·91 (54·45– 69·95)	54·43 (48·02– 60·83)	13·15 (10·10– 16·69)	9·99 (7·55– 12·60)	12·64 (9·95– 15·97)	9·56 (7·46– 11·87)
Malawi	49·32 (46·27– 52·66)	42·60 (39·74– 45·60)	47·72 (44·73– 51·41)	41·63 (38·71– 44·92)	62·98 (56·85– 68·70)	54·97 (49·46– 59·86)	58·15 (51·92– 63·72)	51·35 (46·05– 56·40)	14·81 (11·53– 17·83)	11·32 (8·60– 13·68)	13·51 (10·59– 16·30)	10·37 (7·97– 12·57)
Mozambique	54·79 (51·04– 58·72)	46·48 (42·43– 50·10)	50·88 (47·90– 54·71)	44·16 (41·06– 47·64)	59·86 (52·11– 67·28)	51·69 (45·15– 57·69)	54·28 (47·35– 61·94)	47·80 (41·60– 53·86)	13·85 (9·99– 17·92)	10·32 (7·24– 13·27)	11·87 (9·20– 15·65)	8·84 (6·56– 11·72)
Rwanda	60·97 (58·02– 64·00)	52·57 (48·95– 56·16)	58·08 (55·32– 61·18)	50·39 (47·21– 53·62)	67·68 (60·39– 73·73)	59·11 (52·82– 64·58)	63·73 (61·94)	56·16 (56·40– 61·32)	15·06 (11·69– 18·23)	11·56 (8·91– 14·25)	13·95 (10·78– 16·52)	10·74 (8·17– 12·89)
Somalia	52·51 (41·10– 66·66)	45·86 (35·54– 55·66)	51·66 (41·01– 63·08)	46·33 (36·49– 54·83)	54·98 (42·35– 67·96)	47·96 (37·04– 57·19)	53·40 (41·87– 64·12)	47·76 (37·41– 55·73)	10·69 (7·12– 16·09)	8·10 (5·21– 11·73)	11·26 (7·63– 15·67)	8·71 (5·75– 11·88)
South Sudan	55·74 (43·07– 67·56)	48·44 (37·79– 57·07)	54·87 (43·09– 64·34)	47·89 (37·96– 55·43)	56·87 (43·88– 68·72)	49·77 (38·27– 58·38)	55·45 (43·16– 64·87)	48·64 (38·05– 56·08)	11·97 (7·37– 17·66)	8·98 (5·34– 13·00)	12·33 (7·95– 16·51)	9·12 (5·83– 12·18)
Tanzania	57·17 (53·50– 60·79)	49·44 (46·06– 59·05)	57·12 (53·46– 60·80)	49·74 (46·17– 53·19)	65·47 (56·90– 72·65)	57·40 (50·02– 63·49)	62·69 (54·76– 68·51)	55·32 (48·37– 60·49)	14·46 (10·45– 18·66)	11·18 (7·94– 14·12)	13·93 (10·64– 16·46)	10·71 (8·00– 12·82)
Uganda	55·34 (52·66– 58·44)	47·78 (44·69– 50·97)	51·71 (49·09– 54·57)	45·36 (42·49– 48·23)	64·35 (56·89– 71·57)	56·25 (51·27– 62·36)	58·36 (45·44– 66·03)	51·76 (51·69)	14·11 (10·60– 18·08)	10·90 (7·98– 13·80)	12·40 (9·73– 15·74)	9·54 (7·25– 12·13)
Zambia	48·46 (45·68– 51·26)	42·41 (39·54– 45·15)	45·78 (43·60– 48·33)	40·36 (37·94– 42·72)	59·83 (53·63– 67·00)	52·38 (46·92– 58·21)	53·97 (48·48– 61·10)	47·77 (42·64– 53·66)	12·74 (9·89– 16·41)	9·81 (7·34– 12·57)	10·61 (8·78– 13·49)	8·07 (6·51– 10·38)
Central sub-Saharan Africa	56·41 (52·60– 59·67)	46·71 (41·51– 50·81)	54·60 (51·21– 57·42)	46·50 (42·70– 49·68)	61·04 (55·98– 66·25)	50·71 (44·61– 55·69)	58·94 (53·90– 63·87)	50·56 (46·12– 54·91)	12·79 (10·79– 15·11)	8·87 (6·84– 10·89)	12·61 (10·67– 14·58)	9·12 (7·53– 10·85)
Angola	55·19 (43·52– 67·89)	47·66 (37·79– 56·61)	55·20 (44·08– 64·69)	48·52 (38·95– 56·00)	59·75 (45·75– 71·66)	51·73 (39·66– 60·53)	59·55 (41·43– 68·63)	52·43 (37·41– 59·55)	12·10 (7·38– 17·92)	8·97 (5·42– 12·88)	12·72 (8·13– 16·57)	9·57 (6·13– 12·44)
Central African Republic	47·63 (39·73– 57·62)	41·10 (34·21– 48·26)	43·62 (37·13– 52·61)	38·33 (32·47– 45·21)	51·75 (42·48– 63·18)	45·12 (37·16– 53·72)	47·55 (39·54– 59·15)	42·15 (34·69– 50·66)	10·90 (7·53– 16·52)	8·25 (5·45– 12·08)	9·75 (7·41– 14·86)	7·34 (5·32– 10·57)
Congo	55·61 (52·62– 58·83)	47·40 (43·62– 50·61)	58·08 (54·72– 61·61)	50·39 (46·85– 54·00)	60·01 (52·72– 68·01)	51·32 (45·16– 57·62)	63·40 (55·94– 68·74)	55·23 (48·88– 60·46)	11·54 (8·70– 68·74)	8·42 (6·22– 11·06)	14·00 (11·01– 16·37)	10·51 (8·10– 12·58)

(Table 2 continues on next page)

	2005, at birth				2015, at birth				2015, at age 65 years			
	Females		Males		Females		Males		Females		Males	
	Life expectancy	HALE	Life expectancy	HALE	Life expectancy	HALE						
(Continued from previous page)												
Democratic Republic of the Congo	57·64 (55·14–60·61)	47·07 (40·13–51·19)	55·25 (52·58–57·86)	46·61 (42·75–49·94)	62·14 (56·77–68·11)	51·04 (43·96–56·67)	59·37 (53·21–65·66)	50·66 (45·33–55·99)	13·20 (10·70–16·25)	9·04 (6·57–11·48)	12·70 (10·21–15·37)	9·14 (7·12–11·15)
Equatorial Guinea	56·51 (44·74–67·67)	49·44 (38·84–57·56)	55·55 (44·50–63·43)	48·93 (39·34–55·10)	61·90 (48·89–71·63)	53·99 (42·57–61·32)	60·27 (48·44–67·38)	53·03 (42·56–58·90)	13·77 (8·32–18·74)	10·47 (6·04–13·86)	13·88 (9·07–16·60)	10·47 (6·73–12·86)
Gabon	61·28 (58·52–64·16)	51·68 (47·81–55·07)	59·26 (55·55–63·45)	51·16 (47·61–55·01)	68·45 (61·48–74·32)	57·02 (50·37–62·57)	63·58 (56·80–69·02)	55·00 (48·91–60·04)	14·41 (11·30–17·40)	10·16 (7·60–12·55)	13·98 (11·11–16·34)	10·34 (8·00–12·41)

Data in parentheses are 95% uncertainty intervals. HALE=healthy life expectancy. SDI=Socio-demographic Index.

Table 2: Global, regional, and national or territory life expectancy and HALE at birth, by sex, in 2005 and 2015, and HALE at age 65, by sex, in 2015

For people 60 years and older, several causes, including ischaemic heart disease, chronic kidney disease, diabetes, hearing loss, and Alzheimer's disease and other dementias, ranked among the leading causes of DALYs in 2015 and caused more DALYs than in 2005.

Regional and country-specific HALE

HALE at birth was highest for men in Singapore (72·3 years [95% UI 70·1–74·2]) and for women in Andorra (76·3 years [72·8–79·4]) in 2015. It was lowest in Lesotho for both men (39·1 years [34·3–44·7]) and women (43·8 years [37·9–49·8]; table 2; appendix p 12). HALE at birth in 2015 exceeded 70 years in only 14 geographies for men, while 59 countries and territories surpassed this threshold for women. 13 countries and territories had HALE lower than 50 years for either sex. Since 2005, 121 countries and territories had significant increases in HALE at birth for men and 139 for women, led in both cases by Zimbabwe, whereas HALE at birth worsened for two countries (Syria and Libya) driven mainly by decreases in life expectancy (table 2). HALE at age 65 years was highest in Andorra for both men (15·2 years [UI 13·9–16·3]) and women (19·4 years [17·8–20·8]) in 2015, whereas the lowest HALE for men was in Lesotho (6·9 years [5·2–9·4]) and for women in Afghanistan (6·9 years [5·8–8·2]).

Epidemiological transition

HALE at birth increased continuously and in a largely linear manner with increasing SDI for both sexes (figure 4A). At a SDI of 0·20, average HALE was 46·2 years for men and 47·1 years for women; by a SDI of 0·90, average HALE was 69·8 years for men and 73·8 years for women. Among high SDI regions, North America was the furthest below expected HALE at birth for both sexes, whereas high-income Asia Pacific remained above expected HALE at birth for both sexes and over time. In Australasia, male HALE was consistently close to expected levels, whereas female HALE remained below expected

levels since 1990. With the exception of the Caribbean in 2010 (the year of the Haitian earthquake) HALE at birth throughout Latin America and the Caribbean was generally higher than expected for both sexes. HALE at birth also exceeded expectations in east Asia and north Africa and the Middle East. By contrast, Oceania steadily remained below expected levels of HALE, and all regions within the central Europe, eastern Europe, and central Asia GBD super-region had HALE lower than expected over time, particularly for men. HALE trends in sub-Saharan Africa were heavily influenced by the HIV/AIDS epidemic, particularly in southern sub-Saharan Africa, where HALE was well below expected levels. Notably, after lagging below expected levels of HALE before 2005, eastern sub-Saharan Africa posted average HALE for women that exceeded expected levels. Less pronounced increases occurred for men in eastern sub-Saharan Africa, as HALE essentially equalled expected levels only around 2010 (figure 4A).

Years of functional health lost on average increased as countries developed (figure 4B). Among high-income regions, all regions except for high-income Asia Pacific consistently exceeded expected levels of functional health loss over time, and women generally experienced a higher gap than men. South Asia, north Africa and the Middle East, and central sub-Saharan Africa had generally higher-than-expected functional health loss, whereas a number of regions, including Oceania, southeast Asia, and east Asia, all recorded smaller gaps between life expectancy and HALE than was expected.

The ratio of years of functional health loss to life expectancy, the proportion of life expectancy spent with disability, declined slightly with increasing SDI (figure 4C). Among men and women, central sub-Saharan Africa had the highest proportion of life spent with disability in 2015, although high-income North America had the largest difference between observed and expected levels for that year. For both sexes, several regions showed higher-than-expected proportions of life expectancy spent in ill health (eg, south Asia, north Africa and the Middle East, and




Figure 2: Leading 30 Level 3 causes of global DALYs for both sexes combined, 1990, 2005, 2015, with percentage change in number of DALYs, and all-age, and age-standardised rates
 Causes are connected by lines between time periods; solid lines are increases and dashed lines are decreases. For the time period of 1990 to 2005 and for 2005 to 2015, three measures of change are shown: percent change in the number of DALYs, percent change in the all-age DALY rate, and percent change in the age-standardised DALY rate. Statistically significant changes are shown in bold.
 DALYs=disability-adjusted life-years. COPD=chronic obstructive pulmonary disease.


Australasia), whereas others experienced lower-than-expected levels over time (eg, southeast Asia, east Asia, eastern sub-Saharan Africa, and southern Latin America).

Expected age-standardised YLL rates for many communicable causes and neonatal conditions declined profoundly as SDI increased (figure 5A). At the same time, age-standardised YLD rates for the leading causes of YLDs such as mental and substance abuse disorders and musculoskeletal disorders demonstrate relatively little change. At higher SDIs, the composition of disease burden shifted toward YLDs as the primary driver of burden, mainly due to the differential pace of change. The combined effect of the change in age-specific rates and age-structure change that occur with development is shown in figure 5B, which provides expected all-age YLL and YLD rates for Level 2 causes. Demographic shifts in age-structure potentiate and accelerate the transition from Group 1 conditions toward NCDs in terms of the composition of the

burden of disease that health systems must handle. Of note, from an SDI of 0.8 onwards further declines in the age-standardised rates are matched or exceeded by increases in population age structure so that all-age rates for YLDs actually increase, as do YLL rates for some causes such as neurological conditions. These characterisations of the epidemiological transition demonstrate the double burden of communicable diseases and NCDs for populations with an SDI in the intermediate range.

Observed versus expected total and cause-specific burden

In 2015, the Maldives and Nicaragua had the lowest ratios of observed to expected all-ages DALY rates; many countries throughout Latin America also had lower-than-expected all-ages DALY rates (figure 6). Other regions where observed all-ages DALY rates fell below expected levels, included western Europe (eg, Portugal, Spain, France, Italy, and Sweden); western sub-Saharan Africa (eg, Burkina Faso,

**Figure 3: Leading ten Level 3 causes of global age-specific DALYs in 2015**

Each cause is coloured by the percentage change in age-specific DALYs from 2005 to 2015. NN Preterm=neonatal preterm birth complications. NN Sepsis=neonatal sepsis and other neonatal infections. LRI=lower respiratory infections. Iron=iron-deficiency anaemia. HIV=HIV/AIDS. Back & neck=low back and neck pain. IHD=ischaemic heart disease. NN Enceph=neonatal encephalopathy due to birth asphyxia and trauma. Diarrhoea=diarrhoeal diseases. Skin=skin and subcutaneous diseases. Depression=depressive disorders. Stroke=cerebrovascular disease. Congenital=congenital anomalies. Diabetes=diabetes mellitus. COPD=chronic obstructive pulmonary disease. Alzheimer's=Alzheimer's disease and other dementias. PEM=protein-energy malnutrition. Conduct=conduct disorder. Sense=sense organ diseases. Other NN=other neonatal disorders. Intest inf=intestinal infectious diseases. Violence=interpersonal violence. NN Haemol=haemolytic disease and other neonatal jaundice. Anxiety=anxiety disorders. TB=tuberculosis. Lung C=lung, bronchial, and tracheal cancers. STD=sexually transmitted diseases excluding HIV. Haemog=haemoglobinopathies and haemolytic anaemias. CKD=chronic kidney disease. Other MSK=other musculoskeletal disorders. Drugs=drug use disorders. Stomach C=stomach cancer. HTN HD=hypertensive heart disease. GBD=Global Burden of Disease. DALYs=disability-adjusted life-years.

Niger, and Senegal); eastern sub-Saharan Africa (eg, Burundi and Ethiopia); north Africa and the Middle East (eg, Jordan, Saudi Arabia, and Turkey); east Asia (eg, China); and a subset of countries in South and Southeast Asia (eg, Bangladesh, Sri Lanka, and Vietnam). By contrast, observed all-ages DALY rates exceeded expected levels in southern sub-Saharan Africa, much of central Asia and eastern Europe, and a number of countries in central sub-Saharan Africa. Notably, observed all-ages DALY rates surpassed expected levels in the USA.

Ischaemic heart disease and stroke were the leading two causes of DALYs worldwide in 2015, and 106 geographies also had one of these diseases as the leading cause of DALYs that year (figure 7). Four GBD super-regions showed deviations from this trend: Latin America and the Caribbean, where diabetes and interpersonal violence often resulted in the most DALYs; north Africa and the Middle East, where war was a primary cause of burden; south Asia, where neonatal disorders often ranked among the leading causes of DALYs; and sub-Saharan Africa, where HIV/AIDS or malaria was the leading driver of disease burden in 29 geographies.

Stroke resulted in the most countries (94) having lower observed DALYs than expected based on their SDI. Other leading causes for which observed DALYs were well below expected levels included ischaemic heart disease particularly in Latin America, east Asia, and southeast

Asia; road injuries in north Africa and the Middle East; and lower respiratory infections and diarrhoeal diseases in sub-Saharan Africa. Many high-income countries and territories also had lower-than-expected DALYs from ischaemic heart disease and Alzheimer's disease and other dementias. Road injuries accounted for lower burden than expected in 52 countries and territories, especially in Colombia. Although many Group 1 causes remained among the leading causes of DALYs, observed levels were often lower than expected on the basis of SDI (eg, lower respiratory infections in Ethiopia, diarrhoeal diseases in Afghanistan, and preterm birth complications in Kenya).

By contrast, diabetes was a leading cause for which observed burden exceeded expected levels in many geographies, especially in Oceania and the Caribbean. Observed DALYs due to chronic obstructive pulmonary disease were higher than expected in 30 geographies, as were those due to liver cancer and lung cancer for a subset of countries and territories. Drug use disorders led to more observed DALYs than expected in many high-income countries in 2015, particularly in the USA and Australia. A similar pattern occurred for self-harm, cirrhosis, alcohol use disorders, and drug use disorders throughout eastern Europe, and most prominently in Russia. Interpersonal violence was among the leading two causes of DALYs for six of 11 countries in central and tropical Latin America (Brazil, Colombia, El Salvador, Guatemala, Honduras, and




Figure 4: Co-evolution of HALE (A), functional health lost (life expectancy minus HALE; B), and life expectancy spent with disability (life expectancy minus HALE, divided by HALE; C) with SDI globally and for GBD regions, 1990 to 2015

Coloured lines show global and region values for each metric. Each point in a line represents 1 year starting at 1990 and ending at 2015. In all regions, SDI has increased year on year so progress in SDI is associated with later years for a given region. The black lines indicate trajectories for each geography expected on the basis of SDI alone. GBD=Global Burden of Disease. SDI=Socio-demographic Index. HALE=healthy life expectancy. DALYs=disability-adjusted life-years.

Venezuela), and each had observed burden far surpassing expected levels. Throughout sub-Saharan Africa, HIV/AIDS and malaria resulted in far more DALYs than expected based on SDI.

Heterogeneous trends across and within regions emerged in terms of both leading causes of DALYs and ratios of observed to expected levels. For instance, south Asia's disease burden landscape diverged from global patterns, with both ischaemic heart disease and neonatal disorders ranking among some of the leading causes of

burden and often resulting in higher-than-expected DALYs (figure 7). Stroke and lower respiratory infections resulted in fewer DALYs than expected for most countries in south Asia, yet other causes exacted more DALYs than expected on the basis of SDI (eg, tuberculosis in India and drowning in Bangladesh). Further, the 2015 Nepal earthquake resulted in forces of nature being its leading cause of DALYs that year. Many countries in central Asia experienced observed DALYs that surpassed expected levels due to both Group 1 causes (eg, lower respiratory




Figure 5: Expected relationship between age-standardised YLL and YLD rates and SDI (A) and all-age YLL and YLD rates (per 100 000) and SDI (B) for 21 GBD Level 2 causes

These stacked curves represent the average relationship between SDI and each cause observed across all geographies over the time period 1990 to 2015. In each figure, the y axis goes from lowest SDI to highest SDI. The left side shows rates for YLLs and the right side shows rates for YLDs; higher rates are further from the midline. The difference between (A) and (B) is the effect of shifts in population age structure expected with SDI. GBD=Global Burden of Disease. SDI=Socio-demographic Index. YLDs=years lived with disability. YLLs=years of life lost.




Figure 6: Ratio of observed versus expected age-standardised DALY rates (per 100 000) on the basis of SDI alone for both sexes combined, 2015

Ratios are colour-coded in terms of the magnitude of differences between observed and expected all-ages DALY rates. Blues indicate much lower observed DALYs than expected levels based on SDI, whereas reds reflect that observed DALYs far exceed expected levels given SDI. ATG=Antigua and Barbuda. VCT=Saint Vincent and the Grenadines. LCA=Saint Lucia. TTO=Trinidad and Tobago. TLS=Timor-Leste. FSM=Federated States of Micronesia. SDI=Socio-demographic Index. DALY=disability-adjusted life-years.

infections, preterm birth complications, and neonatal encephalopathy) and NCDs such as hypertensive heart disease. War was the leading cause of DALYs in five countries in north Africa and the Middle East in 2015, including Afghanistan, Iraq, Libya, Syria, and Yemen. Although neonatal sepsis frequently led to higher-than-expected DALYs in sub-Saharan Africa, burden from preterm birth complications fell below expected levels, on the basis of SDI, in most countries. Notably, NCDs such as diabetes ranked among the leading ten causes of DALYs for a subset of countries in sub-Saharan Africa (eg, South Africa), whereas nutritional deficiencies also remained among leading causes of burden in others (eg, Ghana and Zimbabwe); in both instances, observed DALYs generally exceeded expected levels.

Discussion

GBD 2015 results show that the world has become healthier in the past 25 years. Yet, this progress has not been universal. From 1990 to 2015, global HALE at birth increased from 56.7 years to 62.8 years, with 191 of 199 countries or territories recording improved HALE. Since 1990, global HALE at age 65 years also improved by 1.8 years, with an increase in 179 countries or territories. The global number of years of functional health lost grew

during this time, from 8.2 years to 9.1 years. With YLL rates falling at a much faster pace than YLD rates, non-fatal health loss accounted for an increasing proportion of global DALYs, rising from 21.2% in 1990, to 32.1% in 2015. Worldwide progress was largely driven by rapid reductions in DALYs from communicable, maternal, neonatal, and nutritional diseases, although declines in age-standardised DALY rates from NCDs and injuries also contributed to overarching gains. Despite reductions in age-standardised DALY rates, 137 causes had statistically significant increases in total DALYs since 2005, a trend with extensive implications for health systems. Mental and substance use disorders, musculoskeletal disorders, and a range of other conditions including idiopathic developmental intellectual disability, vision and hearing impairment, and neurological disorders all saw rising disease burden since 2005, and few saw any evidence of declining age-specific YLD rates.

Considerable research and policy attention has considered the existence of compression of morbidity, or whether people live healthier lives as their lifespans extend.³⁰ Beyond its profound consequences for financing health systems, compression of morbidity has considerable implications for societal structures and expectations about longevity of careers or timing of

retirement. Compression can be interpreted in both absolute and relative terms. Absolute compression implies that as people live longer lives, they lose fewer years due to functional health loss, whereas relative compression implies that as people live longer lives, the ratio of years of functional health lost to life expectancy declines. Although some evidence shows that compression occurs among people with specific diseases such as diabetes and dementia,^{31,32} national studies show more mixed results.^{33–35} This might not be surprising, as most national studies rely on self-reported health status and chronic conditions, which are then further complicated by variations in how individuals use the response scales and profound framing effects.^{36–39} By contrast, the GBD study provides a more comprehensive and comparable assessment of changes in functional health status by synthesising many types of data, by cause, and applying standardised disability weights to reflect the public's average views of severity of different conditions. GBD 2015 results unequivocally show that as life expectancy increases, people spend more time with reduced functional health status, and thus absolute expansion of morbidity has occurred. This trend is driven by marked declines in age-specific mortality at the same time minimal improvement, if any, has occurred for age-specific YLDs per capita. The proportion of lifespans spent in ill health has remained comparatively constant since 1990, and did not vary as a function of SDI; thus, we found nominal evidence of relative compression at the global level.

Drawing from our empirical characterisation of epidemiological transitions on the basis of SDI, we found that life expectancy and HALE increased linearly with SDI, whereas years of functional health lost climbed with rising SDI. Historically, increases in SDI are associated with a rapid decrease in burden from communicable, neonatal, maternal, and nutritional diseases, the leading killers of children, adolescent girls, and women. Efforts to increase income, provide more years of education, and reduce adolescent and total fertility rates thus might catalyse additional gains for life expectancy, HALE, and reduced disease burden, emphasising the critical role of policy interventions beyond more traditional health service delivery.

At different locations in this continuous process of change, we see evidence of a so-called double burden of disease: from a SDI of approximately 0·35 to 0·60, we expect NCDs and Group 1 causes to each account for at least 20% of disease burden. The average relationships with SDI imply that within a country with wide inequalities in SDI, we should also expect wide variation in disease burden patterns. Subnational disparities might be consistent with variations in subnational SDI, either recording higher or lower burden than that of the national level. Use of average patterns can help to benchmark a country against others, but such assessments can also help to provide insights as to whether public action or other factors are helping make inequalities narrower than

expected based on SDI alone. Given the complexity of health patterns identified for many causes and differential patterns by age and sex, providing some understanding of expected patterns on the basis of SDI alone can help to anchor the exploration of results and could provide some measure of the performance of health systems or the magnitude of avertable burden within each country or territory.

The *Lancet* Commission on Investing in Health⁴⁰ galvanised considerable interest in the notion of a “grand convergence”, such that levels of under-5 mortality, maternal mortality, and some infectious diseases could converge across all countries within a generation. Convergence can be achieved through progress on increasing SDI (ie, increasing per-capita income and average years of schooling, and reducing fertility) and reducing or inverting high ratios between observed and expected (based on SDI alone) health expectancies and health gaps. The Commission has argued that, within a generation, preventable deaths in children and mothers could largely be avoided through increased investment of development assistance for health and expanded national expenditure on health. The vision inspired some of the absolute threshold SDG targets, including reducing under-5 mortality to 25 deaths per 1000 livebirths, neonatal mortality to 12 deaths per 1000 livebirths, and maternal mortality ratio to 70 deaths per 100 000 livebirths. The relationships between these health outcomes, broader health measures, and SDI offer a framework by which the likelihood of such a grand convergence can be assessed. Based on GBD 2015 results, the historical relationship between SDI and health suggests that convergence in the sense of reduced absolute differences in rates is likely to occur with faster improvements in SDI. However, if convergence means smaller relative differences, then improvements in SDI alone might not be sufficient. Our findings show that continued SDI improvement does not appear to be historically associated with absolute convergence in life expectancy or HALE. The *Lancet* Commission also emphasised the importance of hastening progress through strategic investments by donors and governments in effective health technologies, an approach that has the potential to catalyse faster progress than what would be expected on the basis of SDI alone. Convergence in this scenario could then be interpreted as reduction in the ratio of observed to expected burden for low-income and lower-middle-income countries (LMICs), and could be used as an indirect, but summary, metric to monitor health system performance and overall progress toward the SDGs. Since history provides a perspective for identifying which countries have been able to reduce their ratio of observed to expected health outcomes, the comprehensive and longitudinal approach of the GBD is optimally suited for monitoring health system-driven convergence at a macro-level. The same tools can also therefore be used to generate insights on

	1	2	3	4	5	6	7	8	9	10
Global	IHD (0.92)	Stroke (0.93)	LRI (0.65)	Back & neck (0.96)	NN Preterm (0.72)	Diarrhoea (0.76)	Sense (1.00)	NN Enceph (1.17)	Road injuries (0.80)	HIV (0.66)
High income	IHD (0.91)	Back & neck (1.11)	Sense (0.94)	Lung C (0.98)	Stroke (0.59)	Diabetes (1.32)	Alzheimer's (0.77)	Depression (1.11)	COPD (1.24)	Skin (0.92)
High-income North America	IHD (1.79)	Back & neck (1.08)	Diabetes	Lung C (1.30)	Depression (1.31)	COPD (2.35)	Alzheimer's	Drugs (3.82)	Sense (0.89)	Other MSK (1.79)
Canada	Back & neck (1.31)	IHD (1.31)	Lung C (1.20)	Alzheimer's	Sense (0.91)	Diabetes (1.75)	Depression (0.98)	Other MSK (1.56)	Skin (1.02)	Stroke (0.55)
Greenland	Self-harm (4.70)	Lung C (2.65)	IHD (0.74)	Back & neck (1.10)	NN Preterm (1.60)	Sense (0.81)	Congenital (0.80)	COPD (1.81)	Other MSK (2.14)	Skin (1.04)
USA	IHD (1.84)	Back & neck (1.05)	Diabetes	Lung C (1.31)	Depression (1.34)	COPD (2.49)	Drugs (4.01)	Alzheimer's	Sense (0.98)	Other MSK (1.81)
Australasia	Back & neck (1.12)	IHD (1.01)	Depression	Other MSK (1.87)	Sense (0.84)	Diabetes (1.36)	Lung C (0.78)	Stroke	Skin (0.93)	Anxiety (1.72)
Australia	Back & neck (1.12)	IHD (1.02)	Depression	Other MSK (1.93)	Sense (0.84)	Diabetes (1.36)	Lung C (0.78)	Stroke	Skin (0.93)	Drugs (2.93)
New Zealand	Back & neck (1.15)	IHD (0.96)	Depression	Diabetes	Sense (0.85)	Lung C (0.79)	Stroke (0.53)	COPD (1.38)	Skin (0.93)	Other MSK (1.52)
High-income Asia Pacific	Back & neck (0.84)	Stroke (0.74)	IHD (0.52)	Sense (1.04)	Alzheimer's	Self-harm (1.25)	LRI (0.83)	Lung C (0.70)	Diabetes (0.93)	Depression (0.86)
Brunei	Diabetes (4.62)	IHD (2.13)	Back & neck (0.82)	Stroke (1.59)	Road injuries (2.44)	Skin (0.94)	Depression (0.90)	Congenital (1.56)	Sense (1.05)	Iron (1.06)
Japan	IHD (0.52)	Back & neck (0.82)	Sense	Stroke	Alzheimer's	LRI (0.67)	Lung C (0.89)	Self-harm (1.20)	Stomach C (1.15)	Colorect C (0.77)
Singapore	IHD (0.92)	LRI (1.88)	Back & neck (0.60)	Sense (1.01)	Stroke	Depression (0.61)	Skin (0.95)	Lung C (0.94)	CKD (1.13)	Colorect C (0.82)
South Korea	Back & neck (0.90)	Stroke	Self-harm	Diabetes	Sense (1.76)	IHD (1.02)	Lung C (0.46)	Liver C (0.85)	Depression (2.42)	Skin (0.83)
Western Europe	Back & neck (1.27)	IHD (0.75)	Sense	Lung C (0.94)	Stroke	Alzheimer's	Depression (0.49)	Diabetes (0.91)	COPD (0.99)	Migraine (1.17)
Andorra	Back & neck (1.41)	IHD (0.78)	Sense	Alzheimer's	Depression (0.66)	Lung C (1.12)	Stroke (0.66)	Falls (0.44)	Skin (0.87)	Migraine (1.15)
Austria	IHD (1.05)	Back & neck (1.22)	Sense	Lung C (0.93)	Alzheimer's	Stroke (0.87)	Depression (0.71)	Stroke (1.07)	Diabetes (1.09)	Migraine (1.29)
Belgium	Back & neck (1.39)	IHD (0.85)	Lung C	Sense	Alzheimer's	Stroke	COPD (0.55)	Stroke (1.41)	Self-harm (1.01)	Diabetes (1.02)
Cyprus	Back & neck (1.40)	IHD (1.03)	Diabetes	Sense	Depression (0.96)	Road injuries (1.07)	Stroke (1.54)	Migraine (0.49)	Skin (1.14)	Lung C (0.90)
Denmark	Back & neck (1.45)	IHD (1.08)	Lung C	COPD	Stroke	Alzheimer's	Sense (0.78)	Diabetes (0.79)	Depression (1.40)	Depression (1.10)
Finland	IHD (1.30)	Back & neck (1.38)	Alzheimer's	Stroke	Sense (0.64)	Depression (0.85)	Falls (1.23)	Lung C (0.69)	Self-harm (1.00)	Diabetes (0.92)
France	Back & neck (1.15)	IHD (0.43)	Sense	Lung C (0.92)	Depression	Alzheimer's	Stroke (0.63)	Self-harm (0.33)	Falls (0.96)	Diabetes (0.70)
Germany	IHD (1.27)	Back & neck (1.48)	Sense	Lung C (0.98)	Stroke	Depression (0.62)	Alzheimer's	Diabetes (0.61)	COPD (1.26)	Colorect C (0.79)
Greece	IHD (0.89)	Back & neck (1.28)	Stroke	Sense	Lung C (0.95)	Alzheimer's	Depression (1.00)	COPD (0.75)	Road injuries (0.97)	Diabetes (0.58)
Iceland	Back & neck (1.38)	IHD (0.96)	Sense	Lung C	Depression	Alzheimer's	Migraine (0.89)	Skin (0.77)	Stroke (0.41)	Anxiety (1.23)
Ireland	Back & neck (1.25)	IHD (1.12)	Sense	Lung C (0.92)	Depression	Lung C (1.07)	Migraine (0.88)	Stroke (0.90)	Anxiety (0.51)	COPD (1.28)
Israel	Back & neck (1.20)	IHD (0.54)	Diabetes	Sense	Depression	Skin (1.10)	Migraine (0.91)	Stroke (1.14)	Stroke (0.37)	Lung C (0.68)
Italy	Back & neck (1.31)	IHD (0.63)	Sense	Stroke	Alzheimer's	Lung C (0.81)	Diabetes (0.82)	Depression (1.10)	Migraine (1.06)	Skin (1.39)

(Figure 7 continues on next page)

progress, or lack thereof, on specific diseases and outcomes of interest. In some cases, there might be historical or geographical explanations for high burden for some conditions. In other cases, effective preventive and treatment measures have just not yet been implemented or are not functioning effectively.

Shifting from the MDGs to the SDGs dramatically broadened the global health agenda.²³ The SDGs include 17 goals, 169 targets, and 230 indicators; of these measures, 11 goals, 28 targets, and 46 indicators are health related.⁴¹ At present, 33 of the 46 health-related indicators are measured by the GBD study. Amid earlier

	1	2	3	4	5	6	7	8	9	10
Luxembourg	Back & neck (1.42)	IHD (0.90)	Sense (0.92)	Depression (1.11)	Lung C (0.95)	Stroke (0.59)	Alzheimer's (0.71)	Migraine (1.28)	Diabetes (1.23)	Skin (0.86)
Malta	IHD (0.67)	Back & neck (1.31)	Sense (0.89)	Diabetes	Depression (1.02)	Stroke (0.32)	Lung C (0.65)	Alzheimer's (0.76)	Skin (1.00)	Migraine (1.13)
Netherlands	Back & neck (1.50)	IHD (0.82)	Lung C (1.28)	Sense (0.90)	Alzheimer's (0.88)	COPD (1.60)	Diabetes (1.26)	Stroke (0.55)	Depression (0.99)	Skin (0.89)
Norway	Back & neck (1.59)	IHD (1.17)	Lung C (0.90)	Alzheimer's (0.80)	Sense (0.81)	Depression (1.03)	Stroke (0.68)	COPD (1.69)	Anxiety (1.87)	Diabetes (1.25)
Portugal	Back & neck (1.34)	Stroke (0.49)	IHD (0.32)	Sense (0.83)	Diabetes (0.75)	Depression (1.09)	Alzheimer's (0.73)	Lung C (0.74)	LRI (0.61)	Colorect C (0.94)
Spain	Back & neck (0.97)	IHD (0.45)	Sense (0.99)	Alzheimer's (0.84)	Depression (1.09)	Lung C (0.81)	Stroke (0.34)	Diabetes (0.70)	Migraine (1.15)	COPD (0.84)
Sweden	IHD (1.07)	Back & neck (1.25)	Stroke (0.57)	Alzheimer's (0.78)	Sense (0.78)	Diabetes (1.30)	Depression (1.05)	COPD (1.22)	Lung C (0.63)	Skin (0.90)
Switzerland	Back & neck (1.48)	IHD (0.95)	Sense (0.90)	Alzheimer's (0.76)	Depression (1.08)	Lung C (0.81)	Falls (1.38)	Skin (0.88)	Stroke (0.46)	Migraine (1.08)
UK	IHD (0.96)	Back & neck (1.17)	Lung C (1.00)	Stroke (0.63)	COPD (1.60)	Sense (0.79)	Alzheimer's (0.73)	Depression (0.96)	Skin (0.90)	LRI (0.72)
England	IHD (0.92)	Back & neck (1.17)	Lung C (0.95)	Stroke (0.61)	Sense (0.79)	COPD (1.56)	Alzheimer's (0.72)	Depression (0.95)	Skin (0.89)	LRI (0.71)
Northern Ireland	IHD (0.75)	Back & neck (1.11)	Lung C (1.05)	Stroke (0.47)	COPD (1.31)	Sense (0.73)	Depression (0.97)	Alzheimer's (0.79)	LRI (0.74)	Skin (0.88)
Scotland	IHD (1.49)	Back & neck (1.23)	Lung C (1.42)	Stroke (0.93)	COPD (2.12)	Migraine (1.75)	Sense (0.79)	Alzheimer's (0.75)	Depression (0.97)	Asthma (2.13)
Wales	IHD (1.07)	Back & neck (1.15)	Lung C (1.05)	Stroke (0.65)	Alzheimer's (0.87)	COPD (1.66)	Sense (0.79)	Depression (1.07)	Asthma (2.28)	LRI (0.77)
Southern Latin America	IHD (0.66)	Back & neck (0.88)	Stroke (0.54)	LRI (1.18)	Depression (1.07)	Sense (0.84)	Diabetes (0.99)	Road injuries (0.81)	Congenital (1.07)	Skin (0.97)
Argentina	IHD (0.75)	Back & neck (0.87)	LRI (1.45)	Stroke (0.52)	Depression (1.07)	Diabetes (1.02)	Sense (0.84)	Congenital (1.09)	Road injuries (0.81)	Skin (0.97)
Chile	Back & neck (0.88)	IHD (0.49)	Stroke (0.49)	Depression (0.59)	Sense (0.86)	Diabetes (1.00)	Skin (0.97)	Road injuries (0.81)	Anxiety (1.57)	Self-harm (0.78)
Uruguay	IHD (0.49)	Stroke (0.57)	Back & neck (0.89)	Lung C (1.19)	Sense (0.80)	COPD (1.05)	Depression (1.03)	Diabetes (0.68)	Self-harm (1.20)	Road injuries (0.76)
Central Europe, eastern Europe, and central Asia	IHD (2.45)	Stroke (1.71)	Back & neck (1.20)	Sense (1.16)	Self-harm (1.42)	Lung C (1.05)	Depression (1.10)	LRI (1.06)	Road injuries (1.20)	Diabetes (0.93)
Eastern Europe	IHD (3.25)	Stroke (2.12)	Back & neck (1.21)	Self-harm (1.83)	Sense (1.25)	CMP (7.41)	Road injuries (1.74)	Drugs (6.23)	Depression (1.17)	Alcohol (8.90)
Belarus	IHD (4.09)	Stroke (2.00)	Back & neck (1.19)	Self-harm (1.95)	Sense (1.18)	Road injuries (2.00)	Lung C (1.28)	Falls (1.90)	Drugs (6.47)	Depression (1.15)
Estonia	IHD (1.89)	Back & neck (1.16)	Stroke (0.92)	Sense (1.08)	HTN HD (7.31)	Depression (1.25)	Lung C (0.89)	Alcohol (8.56)	Diabetes (0.96)	Alzheimer's (0.59)
Latvia	IHD (2.50)	Stroke (1.73)	Back & neck (1.24)	Sense (1.22)	CMP (6.87)	Self-harm (1.20)	Lung C (0.95)	Diabetes (1.26)	Depression (0.97)	Alcohol (6.81)
Lithuania	IHD (2.21)	Stroke (1.15)	Back & neck (1.23)	Self-harm (2.03)	Sense (1.17)	Depression (1.16)	Lung C (0.90)	Road injuries (1.26)	Diabetes (0.89)	Alcohol (7.02)
Moldova	IHD (1.45)	Stroke (1.05)	Back & neck (1.20)	Sense (0.96)	Depression (1.00)	Cirr Alc (4.21)	Self-harm (1.06)	LRI (0.75)	Skin (0.94)	Road injuries (0.51)
Russia	IHD (3.28)	Stroke (2.49)	Back & neck (1.20)	Self-harm (2.00)	Sense (1.30)	CMP (9.71)	Road injuries (2.09)	Alcohol (10.72)	Drugs (5.90)	Depression (1.14)
Ukraine	IHD (3.35)	Stroke (1.52)	Back & neck (1.26)	Sense (1.17)	Drugs (8.36)	Depression (1.29)	Self-harm (1.36)	Lung C (0.84)	HIV (1.64)	Road injuries (1.12)
Central Europe	IHD (1.46)	Stroke (1.19)	Back & neck (1.29)	Lung C (1.27)	Sense (1.10)	Diabetes (1.09)	Depression (1.06)	Colorect C (1.03)	Self-harm (0.86)	Road injuries (0.93)
Albania	IHD (0.90)	Stroke (0.99)	Back & neck (1.37)	Sense (0.94)	Depression (0.98)	Lung C (0.93)	Other Cardio (1.93)	Skin (0.81)	Iron (1.25)	Congenital (0.77)

(Figure 7 continues on next page)

discussions and negotiations over SDG 3 indicators, HALE was proposed as an indicator of overarching health status and progress;² this proposal was not ultimately adopted in the final set of indicators. HALE provides a strong summary measure of overall health status because it accounts for functional health loss in

addition to age-specific mortality. Other summary development measures, such as the Human Development Index, have considered replacing life expectancy with HALE as an input to the overall assessment.⁴² The GBD study measures health outcomes that are both amenable to intervention and could be risk

	1	2	3	4	5	6	7	8	9	10
Bosnia and Herzegovina	IHD (0.62)	Stroke (0.72)	Back & neck (1.16)	Diabetes	Sense (0.98)	Lung C (1.16)	CMP (5.19)	Depression (0.98)	Skin (0.86)	Colorect C (0.91)
Bulgaria	IHD (1.74)	Stroke (1.59)	Back & neck (1.28)	Sense	Lung C (1.12)	Diabetes (1.15)	HTN HD (5.12)	Other Cardio (2.40)	Depression (1.06)	Colorect C (1.06)
Croatia	IHD (1.00)	Back & neck (1.37)	Stroke (0.85)	Lung C (1.21)	Sense (0.97)	Diabetes	Depression (1.04)	Colorect C (1.13)	Alzheimer's (0.69)	COPD (0.77)
Czech Republic	IHD (2.15)	Back & neck (1.38)	Stroke (1.09)	Sense (1.09)	Lung C (1.11)	Diabetes	Depression (1.48)	Colorect C (1.13)	Alzheimer's (0.62)	Skin (0.87)
Hungary	IHD (1.82)	Back & neck (1.38)	Stroke (1.04)	Lung C (1.61)	Sense (1.05)	Diabetes	Colorect C (1.31)	Depression (1.07)	COPD (1.46)	Self-harm (0.96)
Macedonia	Stroke (1.53)	IHD (0.98)	Back & neck (1.26)	Diabetes	Sense (0.99)	Lung C (1.11)	Depression (1.01)	CMP (3.63)	Skin (0.86)	Migraine (0.98)
Montenegro	IHD (1.18)	Stroke (1.64)	Back & neck (1.27)	Lung C (1.43)	Sense (1.04)	Diabetes	Depression (1.04)	Skin (0.88)	Road injuries (0.72)	CMP (3.14)
Poland	IHD (1.67)	Back & neck (1.21)	Stroke (1.02)	Lung C (1.45)	Sense (1.27)	Diabetes	Depression (1.07)	Self-harm (1.06)	Road injuries (1.37)	Colorect C (1.03)
Romania	IHD (1.39)	Stroke (1.53)	Back & neck (1.33)	Sense (1.03)	Lung C (1.04)	Depression	HTN HD (3.57)	LRI (0.73)	Diabetes (0.64)	Road injuries (0.81)
Serbia	IHD (1.03)	Stroke (1.27)	Back & neck (1.31)	Lung C (1.38)	Diabetes	Sense (1.23)	Depression (1.02)	Colorect C (1.13)	Self-harm (0.96)	Road injuries (0.71)
Slovakia	IHD (2.05)	Back & neck (1.34)	Stroke (1.05)	Sense (1.06)	Lung C (1.02)	Depression	Diabetes	Colorect C (1.24)	Skin (0.88)	LRI (0.83)
Slovenia	Back & neck (1.35)	IHD (0.76)	Stroke (0.63)	Sense (0.99)	Lung C (0.86)	Depression	Diabetes	Alzheimer's (0.62)	Self-harm (0.84)	Colorect C (0.85)
Central Asia	IHD (1.95)	Stroke (1.44)	LRI (2.00)	Back & neck (1.01)	NN Preterm (1.00)	NN Enceph (2.17)	Congenital	Road injuries (0.70)	Sense (0.98)	Depression (0.97)
Armenia	IHD (1.52)	Stroke (0.87)	Diabetes	Back & neck (1.00)	Sense (1.03)	Lung C (1.29)	Depression	Congenital (1.21)	COPD (0.93)	Road injuries (0.64)
Azerbaijan	IHD (2.31)	LRI (3.36)	Stroke (1.25)	Back & neck (0.98)	NN Enceph (7.12)	Diabetes	NN Preterm (2.17)	Sense (1.08)	Depression	Congenital (1.07)
Georgia	IHD (1.70)	Stroke (1.55)	Back & neck (0.93)	Diabetes	Sense (1.26)	Road injuries	Depression	Lung C (0.95)	COPD (1.07)	HTN HD (3.42)
Kazakhstan	IHD (2.77)	Stroke (2.18)	Back & neck (1.04)	Self-harm	Road injuries (2.14)	NN Preterm (1.46)	NN Enceph (2.88)	Iron	LRI (1.76)	Sense (1.10)
Kyrgyzstan	IHD (1.62)	Stroke (1.54)	LRI (1.19)	NN Preterm (0.98)	NN Enceph (1.61)	Back & neck	Congenital	Road injuries (0.69)	Iron	Depression (0.91)
Mongolia	IHD (1.82)	Stroke (2.67)	Self-harm	LRI (3.02)	Liver C (2.19)	NN Enceph (10.57)	Back & neck	Road injuries (0.98)	Congenital	NN Preterm (0.91)
Tajikistan	LRI (1.27)	IHD (1.19)	NN Preterm (0.66)	Stroke	Back & neck (1.06)	Diarrhoea	Congenital	NN Enceph (0.70)	Iron	Haemog (3.79)
Turkmenistan	IHD (2.96)	LRI (6.42)	Stroke	Congenital (2.21)	Back & neck (0.94)	NN Preterm (2.59)	NN Enceph (5.53)	Diabetes	Depression	Sense (1.06)
Uzbekistan	IHD (1.71)	LRI (2.53)	Stroke (1.12)	Back & neck (1.00)	NN Enceph (2.60)	HTN HD (5.41)	Depression	Diabetes	Road injuries (0.89)	Sense (0.93)
Latin America and Caribbean	IHD (0.63)	Violence (4.14)	Back & neck (0.87)	Diabetes	Road injuries	Sense	Stroke	Depression	LRI (0.75)	CKD (1.34)
Central Latin America	Violence (4.78)	IHD (0.60)	Diabetes	Back & neck (1.55)	CKD	Sense	Congenital	Road injuries	LRI (0.68)	
Colombia	Violence (6.04)	IHD (0.50)	Back & neck (0.94)	Sense	Depression	Skin	Road injuries	Congenital	Diabetes	Stroke (0.35)
Costa Rica	Back & neck (0.89)	IHD (0.43)	Sense	Depression	Road injuries	CKD	Congenital	Skin	Diabetes	Iron (1.01)
El Salvador	Violence (8.82)	IHD (0.57)	CKD	Back & neck (2.01)	Diabetes	Road injuries	Sense	Iron	LRI (0.60)	Depression (0.90)
Guatemala	LRI (1.04)	Violence (4.70)	Diabetes	Diarrhoea	Back & neck (0.90)	IHD	Congenital	Iron	CKD	Sense (0.94)

(Figure 7 continues on next page)

standardised, thus offering a useful set of metrics to monitor progress toward specific SDG targets, such as the aim of target 3.8 to achieving effective universal health coverage.

DALYs and other health-gap metrics are one of many potential inputs when setting health policy and investment

priorities, but major research organisations and funders such as the US National Institutes of Health and others note the use of DALYs to inform budgeting decisions.^{43–46} Beyond health metrics, many other inputs are required for decision making, ranging from the effectiveness of different adoptable policies and programmes, to key

	1	2	3	4	5	6	7	8	9	10
Honduras	Violence (5.08)	IHD (0.96)	Stroke (0.83)	Congenital (1.03)	Back & neck (0.98)	Sense (0.95)	Skin (1.04)	Diabetes (0.72)	Depression (0.88)	NN Preterm (0.40)
Mexico	Diabetes (2.22)	IHD (0.59)	CKD (2.59)	Back & neck (0.79)	Sense (1.17)	Congenital (1.09)	Violence (3.13)	Road injuries (0.76)	Depression (0.91)	Skin (0.93)
Nicaragua	CKD (1.76)	Back & neck (0.96)	IHD (0.44)	Diabetes (0.77)	Sense (0.95)	Congenital (0.71)	LRI (0.38)	Depression (0.87)	Skin (0.94)	Road injuries (0.39)
Panama	Back & neck (0.91)	IHD (0.48)	Diabetes (1.40)	Violence (4.60)	Sense (1.19)	Congenital (1.02)	Road injuries (0.83)	Depression (0.99)	Stroke (0.46)	LRI (1.04)
Venezuela	Violence (9.65)	IHD (0.86)	Iron (3.05)	Road injuries (1.15)	Back & neck (0.90)	Diabetes (1.35)	Sense (1.12)	Congenital (0.93)	Stroke (0.54)	CKD (1.64)
Andean Latin America	LRI (1.31)	Back & neck (0.93)	IHD (0.39)	Sense (1.07)	Depression (1.09)	Road injuries (0.67)	Iron (1.34)	Skin (1.01)	Diabetes (0.74)	Congenital (0.67)
Bolivia	LRI (1.04)	IHD (0.51)	Back & neck (1.00)	Stroke (0.62)	Iron (1.51)	NN Enceph (1.16)	Road injuries (0.60)	NN Preterm (0.55)	Sense (0.97)	CKD (1.31)
Ecuador	IHD (0.45)	LRI (1.18)	Back & neck (0.91)	Road injuries (0.95)	Diabetes (1.20)	Sense (1.02)	CKD (1.58)	Depression (1.08)	Congenital (0.78)	Iron (1.32)
Peru	LRI (1.58)	Back & neck (0.92)	Sense (1.14)	Depression (1.11)	IHD (0.32)	Skin (1.03)	Iron (1.28)	Road injuries (0.56)	Congenital (0.65)	Migraine (1.07)
Caribbean	IHD (0.93)	Stroke (0.82)	Diabetes (1.58)	HIV (1.09)	LRI (0.80)	Back & neck (0.89)	Sense (1.06)	Congenital (1.06)	Road injuries (0.85)	Depression (1.11)
Antigua	Diabetes (3.67)	IHD (1.07)	Stroke (1.19)	Back & neck (0.82)	Depression (1.20)	Sense (1.21)	Skin (1.12)	LRI (1.49)	CKD (2.03)	NN Preterm (3.06)
The Bahamas	IHD (1.20)	Diabetes (2.82)	Stroke (1.15)	Back & neck (0.82)	HIV (2.74)	Violence (12.84)	Sense (1.19)	Depression (1.18)	LRI (1.67)	HTN HD (9.97)
Barbados	Diabetes (2.95)	IHD (0.61)	Stroke (0.73)	Back & neck (0.81)	Sense (1.07)	LRI (1.22)	Depression (1.15)	Skin (1.20)	CKD (1.39)	Prostate C (3.69)
Belize	HIV (1.07)	Diabetes (1.73)	Violence (3.59)	IHD (0.70)	Back & neck (0.95)	Back & neck (0.84)	Road injuries (0.72)	Stroke (0.68)	NN Preterm (0.62)	Depression (1.04)
Bermuda	IHD (1.82)	Back & neck (0.89)	Diabetes (2.83)	Depression (1.25)	Sense (1.19)	Stroke (0.97)	Skin (1.04)	HIV (2.89)	Iron (1.32)	Lung C (0.99)
Cuba	IHD (0.78)	Stroke (0.57)	Back & neck (0.82)	Sense (1.11)	Lung C (1.12)	Depression (1.14)	Diabetes (0.82)	LRI (0.82)	Skin (1.08)	Alzheimer's (0.85)
Dominica	Diabetes (2.80)	IHD (0.67)	Stroke (0.72)	Back & neck (0.83)	CKD (2.21)	LRI (1.38)	Sense (1.07)	NN Preterm (1.91)	Depression (1.12)	Skin (1.16)
Dominican Republic	IHD (0.83)	NN Preterm (1.19)	Stroke (0.69)	Road injuries (0.92)	Back & neck (0.82)	HIV (0.75)	Congenital (0.89)	Sense (0.98)	Depression (1.07)	LRI (0.79)
Grenada	Diabetes (3.04)	IHD (0.91)	Stroke (1.16)	LRI (2.14)	Back & neck (0.80)	CKD (2.26)	Depression (1.11)	Sense (1.06)	Skin (1.18)	Road injuries (0.75)
Guyana	IHD (1.34)	HIV (2.34)	Diabetes (2.27)	Stroke (1.51)	Self-harm (2.36)	NN Preterm (0.97)	LRI (1.01)	Back & neck (0.84)	Road injuries (0.80)	Violence (2.80)
Haiti	HIV (1.53)	LRI (0.72)	Diarrhoea (0.71)	IHD (1.14)	Stroke (1.07)	Congenital (1.26)	Road injuries (1.00)	NN Preterm (0.46)	NN Enceph (0.76)	Diabetes (1.65)
Jamaica	Diabetes (2.28)	Stroke (0.85)	IHD (0.43)	Violence (4.13)	Back & neck (0.83)	NN Preterm (1.48)	Sense (1.00)	Depression (1.10)	CKD (1.39)	Skin (1.11)
Puerto Rico	Diabetes (3.97)	IHD (1.19)	Back & neck (0.88)	Sense (1.20)	Violence (22.36)	Depression (1.23)	CKD (1.97)	Stroke (0.61)	Skin (1.09)	LRI (1.07)
St Lucia	Diabetes (2.38)	Stroke (0.86)	IHD (0.55)	Back & neck (0.83)	Sense (1.05)	Depression (1.11)	LRI (1.16)	Violence (3.79)	Skin (1.14)	CKD (1.46)
Saint Vincent and the Grenadines	Diabetes (3.41)	IHD (0.99)	Stroke (1.00)	Back & neck (0.82)	Violence (4.84)	NN Preterm (1.94)	LRI (1.34)	Sense (1.05)	HIV (0.97)	Depression (1.10)
Suriname	IHD (0.75)	Stroke (1.09)	Diabetes (1.58)	NN Preterm (1.83)	HIV (0.96)	Self-harm (1.97)	Back & neck (0.82)	Road injuries (0.89)	LRI (1.19)	Congenital (1.09)
Trinidad and Tobago	Diabetes (6.79)	IHD (1.81)	Stroke (1.30)	Back & neck (0.81)	Violence (13.68)	Sense (1.17)	Depression (1.18)	CKD (2.22)	Road injuries (1.36)	HIV (1.81)
Virgin Islands	IHD (2.41)	Diabetes (3.58)	Stroke (1.12)	Back & neck (0.89)	Violence (28.20)	Sense (1.19)	CKD (2.14)	Prostate C (4.76)	Depression (1.24)	Skin (1.16)

(Figure 7 continues on next page)

social, cultural, and ethical considerations. Nonetheless, DALYs and other summary health measures might have an even more prominent role in the future in setting research and development priorities within the health sector, particularly in the absence of robust information on the effectiveness or likely success of various research

projects and programmes.⁴⁷ As global health research funders increasingly use DALYs to shape priority-setting processes, health challenges faced by populations with less health-care market power, namely poor people, will inevitably receive more attention. Shifting to the use of disease burden for programme design and evaluation

	1	2	3	4	5	6	7	8	9	10
Tropical Latin America	IHD (0·67)	Violence (4·65)	Stroke (0·68)	Road injuries (0·98)	Back & neck (0·87)	Depression (1·23)	Diabetes (0·93)	Sense (0·99)	Anxiety (2·14)	LRI (0·67)
Brazil	IHD (0·67)	Violence (4·75)	Stroke (0·68)	Road injuries (0·98)	Back & neck (0·86)	Depression (1·23)	Diabetes (0·93)	Sense (0·99)	Anxiety (2·15)	LRI (0·67)
Paraguay	IHD (0·66)	Road injuries (0·93)	Stroke (0·76)	Back & neck (0·97)	Diabetes (1·11)	Depression (1·24)	Congenital (0·83)	NN Preterm (0·63)	Sense (0·97)	Skin (1·13)
Southeast Asia, east Asia, and Oceania	Stroke (1·17)	IHD (0·59)	Back & neck (0·86)	Road injuries (0·82)	COPD (1·21)	Sense (0·94)	Diabetes (0·67)	Lung C (1·25)	Skin (0·98)	Depression (0·74)
East Asia	Stroke (1·14)	IHD (0·53)	Back & neck (0·89)	COPD (1·29)	Road injuries (0·87)	Sense (0·91)	Lung C (1·36)	Liver C (3·16)	Depression (0·73)	Diabetes (0·47)
China	Stroke (1·14)	IHD (0·53)	Back & neck (0·88)	COPD (1·28)	Road injuries (0·86)	Sense (0·91)	Lung C (1·37)	Liver C (3·13)	Depression (0·74)	Diabetes (0·45)
North Korea	Stroke (1·52)	IHD (0·62)	COPD (1·68)	Back & neck (1·05)	Road injuries (0·94)	Liver C (4·38)	Sense (0·77)	Lung C (2·11)	Stomach C (2·43)	LRI (0·37)
Taiwan (province of China)	Diabetes (2·76)	Back & neck (0·99)	IHD (0·63)	Stroke (0·77)	Sense (1·08)	Liver C (3·50)	Lung C (0·88)	Road injuries (1·47)	Other MSK (1·57)	Skin (0·93)
Southeast Asia	Stroke (1·23)	IHD (0·75)	Diabetes (1·20)	LRI (0·89)	Back & neck (0·81)	Road injuries (0·74)	Sense (1·01)	TB (3·43)	NN Preterm (0·64)	Skin (1·07)
Cambodia	IHD (1·04)	LRI (0·63)	Stroke (1·00)	NN Preterm (0·66)	Road injuries (0·69)	Back & neck (0·94)	Congenital (0·74)	Iron (0·94)	HIV (0·37)	Sense (0·95)
Indonesia	Stroke (1·70)	IHD (0·93)	Diabetes (1·51)	TB (6·35)	Road injuries (0·76)	Back & neck (0·76)	NN Preterm (0·76)	Sense (0·99)	Diarrhoea (2·93)	LRI (0·68)
Laos	LRI (1·34)	NN Preterm (1·23)	IHD (1·12)	Stroke (1·08)	Congenital (1·08)	Road injuries (0·94)	Diarrhoea (0·78)	NN Enceph (0·76)	Back & neck (0·90)	Diabetes (1·13)
Malaysia	IHD (1·23)	LRI (2·28)	Stroke (0·94)	Road injuries (1·18)	Back & neck (0·75)	Diabetes (1·49)	Skin (1·15)	Sense (1·00)	Depression (0·84)	Other MSK (1·84)
Maldives	IHD (0·49)	Back & neck (0·82)	Sense (0·98)	Congenital (0·72)	Iron (1·35)	Skin (1·06)	Depression (0·75)	Stroke (0·34)	Diabetes (0·41)	Self-harm (0·69)
Mauritius	Diabetes (3·86)	IHD (0·93)	CKD (3·13)	Stroke (0·76)	Back & neck (0·82)	Sense (1·15)	Skin (1·10)	Depression (0·84)	Other MSK (1·72)	Asthma (1·58)
Myanmar	Stroke (1·14)	LRI (0·86)	TB (2·15)	Diabetes (1·07)	IHD (0·45)	Sense (1·13)	Back & neck (0·79)	COPD (1·12)	NN Enceph (1·08)	Congenital (0·84)
Philippines	IHD (1·11)	LRI (1·43)	Stroke (1·25)	Diabetes (1·30)	Back & neck (0·95)	Congenital (0·81)	NN Preterm (0·61)	Sense (1·05)	TB (3·97)	Skin (1·11)
Sri Lanka	IHD (0·72)	Diabetes (1·46)	Back & neck (0·81)	Self-harm (2·11)	Sense (1·14)	Stroke (0·56)	Asthma (2·09)	Skin (1·03)	Depression (0·81)	COPD (0·78)
Seychelles	IHD (0·82)	LRI (2·14)	Back & neck (0·79)	HTN HD (7·91)	Stroke (0·63)	CKD (2·20)	Sense (1·18)	Diabetes (1·26)	Skin (1·07)	Depression (0·84)
Thailand	IHD (0·44)	Stroke (0·63)	Road injuries (1·29)	LRI (1·61)	Diabetes (0·96)	Back & neck (0·73)	Sense (1·04)	Liver C (3·94)	CKD (1·34)	HIV (0·71)
Timor-Leste	LRI (0·60)	NN Preterm (0·59)	Congenital (0·78)	IHD (0·68)	Stroke (0·66)	Measles (3·94)	Diarrhoea (0·31)	NN Enceph (0·44)	Back & neck (0·87)	Skin (1·04)
Vietnam	Stroke (1·07)	IHD (0·40)	Road injuries (0·78)	Back & neck (0·85)	Sense (0·92)	Diabetes (0·60)	Lung C (1·88)	Skin (1·06)	Congenital (0·79)	Depression (0·77)
Oceania	LRI (1·07)	IHD (2·05)	Stroke (1·82)	Diabetes (3·91)	COPD (3·33)	Asthma (3·18)	Road injuries (0·96)	NN Preterm (0·42)	Iron (1·06)	CKD (2·38)
American Samoa	Diabetes (3·50)	IHD (1·11)	Skin (1·64)	Back & neck (0·90)	Stroke (0·99)	CKD (2·20)	Sense (1·03)	LRI (0·83)	Iron (0·91)	Congenital (0·43)
Federated States of Micronesia	IHD (1·54)	Diabetes (2·35)	Stroke (1·38)	LRI (0·73)	Skin (1·41)	CKD (1·91)	Back & neck (0·91)	Self-harm (1·42)	Road injuries (0·49)	COPD (1·26)
Fiji	Diabetes (7·08)	IHD (1·95)	LRI (2·35)	Stroke (1·27)	CKD (3·13)	Asthma (3·05)	Back & neck (0·84)	NN Preterm (1·19)	Congenital (1·04)	Skin (1·40)
Guam	IHD (3·95)	Diabetes (4·34)	Stroke (1·91)	Back & neck (0·94)	Self-harm (1·45)	Lung C (1·54)	Skin (1·38)	Sense (1·27)	LRI (2·13)	CKD (2·92)
Kiribati	Stroke (2·16)	Diabetes (4·10)	IHD (1·46)	LRI (0·68)	NN Preterm (0·53)	TB (2·01)	Violence (2·70)	Asthma (2·61)	NN Enceph (0·74)	Diarrhoea (0·46)

(Figure 7 continues on next page)

would benefit poor people, but also potentially increase overall efficiency of health research.⁴⁸ For instance, individuals who have historically underfunded conditions, such as mental health disorders, substance use, and musculoskeletal conditions, would benefit from the greater use of DALYs in decision-making processes.

Global progress has been especially rapid in reducing disease burden due to a number of communicable diseases, including diarrhoeal diseases, lower respiratory infections, tuberculosis, syphilis, typhoid, paratyphoid, and vaccine-preventable infections such as hepatitis B, measles, tetanus, and *Haemophilus influenzae* type b. To

	1	2	3	4	5	6	7	8	9	10
Marshall Islands	Diabetes (3.56)	IHD (1.61)	Stroke (1.46)	CKD (2.85)	LRI (0.78)	Skin (1.39)	Neck & back (0.88)	NN Preterm (0.48)	Iron (1.23)	Road injuries (0.51)
Northern Mariana Islands	Diabetes (4.03)	Neck & back (0.86)	IHD (1.39)	Skin (1.34)	Stroke (1.62)	Depression (0.82)	Road injuries (1.04)	Sense (1.28)	Self-harm (0.64)	Other MSK (1.60)
Papua New Guinea	LRI (1.02)	IHD (1.85)	Stroke (1.71)	COPD (3.50)	Diabetes (3.03)	Asthma (3.13)	Road injuries (0.98)	Diarrhoea (0.43)	Iron (1.06)	NN Preterm (0.38)
Samoa	IHD (0.94)	Diabetes (2.09)	Stroke (0.87)	Skin (1.41)	Neck & back (0.90)	CKD (1.78)	LRI (0.54)	Sense (0.92)	Congenital (0.45)	Iron (0.84)
Solomon Islands	IHD (2.00)	Stroke (1.67)	Diabetes (3.69)	LRI (0.46)	CKD (2.65)	COPD (1.66)	NN Preterm (0.34)	Skin (1.32)	Road injuries (0.65)	Neck & back (0.95)
Tonga	Diabetes (2.23)	IHD (0.90)	Stroke (0.75)	LRI (0.72)	Skin (1.44)	Neck & back (0.92)	NN Preterm (0.53)	Sense (0.83)	COPD (1.03)	Asthma (1.67)
Vanuatu	IHD (1.88)	Stroke (1.81)	Diabetes (2.60)	LRI (0.88)	NN Preterm (0.56)	COPD (1.53)	Skin (1.42)	Road injuries (0.63)	CKD (1.70)	Neck & back (0.91)
North Africa and Middle East	IHD (1.14)	War (796.25)	Congenital (1.18)	NN Preterm (0.78)	Road injuries (0.97)	Neck & back (1.07)	Stroke (0.84)	LRI (0.52)	Diabetes (1.26)	Depression (0.93)
North Africa and Middle East	IHD (1.14)	War (796.25)	Congenital (1.18)	NN Preterm (0.78)	Road injuries (0.97)	Neck & back (1.07)	Stroke (0.84)	LRI (0.52)	Diabetes (1.26)	Depression (0.93)
Afghanistan	War (1033.96)	LRI (0.70)	IHD (4.31)	Congenital (1.56)	Stroke (2.15)	NN Preterm (0.77)	Road injuries (2.38)	Other Unint (9.71)	Diarrhoea (0.24)	Violence (4.00)
Algeria	IHD (0.65)	Diabetes (1.10)	NN Preterm (0.75)	Neck & back (1.04)	Stroke (0.69)	Road injuries (0.69)	Congenital (0.93)	Depression (0.95)	Sense (0.82)	Skin (0.83)
Bahrain	Diabetes (3.31)	Neck & back (0.95)	IHD (0.83)	Depression (1.11)	Skin (1.04)	Road injuries (0.57)	Other MSK (1.92)	Sense (1.02)	Congenital (0.90)	Migraine (1.07)
Egypt	IHD (1.32)	Congenital (1.26)	Stroke (1.04)	LRI (0.77)	Diabetes (1.18)	Neck & back (1.08)	Cirr HepC (7.30)	Iron (1.53)	NN Preterm (0.48)	Sense (0.93)
Iran	IHD (1.24)	Road injuries (1.69)	Neck & back (1.03)	Stroke (0.76)	Diabetes (1.20)	Congenital (1.02)	Depression (1.05)	Sense (0.94)	Other Cardio (3.76)	Drugs (4.09)
Iraq	War (1932.51)	IHD (1.83)	IHD (1.22)	NN Preterm (0.70)	Diabetes (1.78)	Stroke (1.14)	NN Sepsis (2.44)	Road injuries (0.68)	Neck & back (1.08)	LRI (0.40)
Jordan	Congenital (1.09)	IHD (0.69)	Neck & back (1.00)	Diabetes (1.61)	NN Preterm (0.70)	Road injuries (0.67)	Depression (1.01)	Skin (0.91)	Iron (0.97)	Sense (0.92)
Kuwait	IHD (1.94)	Neck & back (0.88)	Depression (1.18)	Congenital (1.62)	Road injuries (1.32)	Diabetes (2.18)	Migraine (1.06)	Skin (0.88)	Sense (1.09)	Iron (1.18)
Lebanon	IHD (1.01)	War (5581.30)	Diabetes (1.72)	Neck & back (0.83)	Depression (1.02)	Sense (0.90)	Stroke (0.44)	Congenital (1.00)	Skin (0.93)	Iron (1.35)
Libya	War (1966.75)	IHD (0.92)	Neck & back (1.04)	Congenital (1.09)	Road injuries (0.83)	Stroke (0.74)	Diabetes (0.98)	NN Preterm (0.61)	Depression (1.00)	Other Trans (10.71)
Morocco	Diabetes (1.70)	IHD (0.61)	Neck & back (1.14)	NN Preterm (0.57)	Stroke (0.49)	Road injuries (0.62)	Congenital (0.78)	Sense (0.79)	Depression (0.88)	LRI (0.27)
Palestine	IHD (1.31)	NN Preterm (0.69)	Congenital (0.86)	Neck & back (1.06)	Depression (1.29)	Road injuries (0.49)	Stroke (0.70)	LRI (0.26)	Skin (0.81)	CKD (1.22)
Oman	Road injuries (1.60)	Other Cardio (9.17)	Neck & back (0.95)	Diabetes (2.26)	IHD (0.79)	Depression (1.07)	Congenital (0.71)	Sense (1.01)	Migraine (1.10)	Skin (0.86)
Qatar	Road injuries (1.48)	Neck & back (0.94)	Diabetes (2.92)	Depression (1.15)	IHD (0.67)	Migraine (1.12)	Skin (0.94)	NN Preterm (0.61)	Sense (0.96)	Anxiety (1.05)
Saudi Arabia	Neck & back (0.99)	IHD (0.84)	Road injuries (1.23)	Congenital (1.29)	Depression (1.11)	Skin (0.95)	Migraine (1.27)	Diabetes (1.15)	Sense (1.01)	NN Preterm (1.01)
Sudan	NN Preterm (1.21)	Congenital (1.54)	IHD (1.59)	LRI (0.52)	Road injuries (1.58)	Diarrhoea (0.52)	Stroke (1.05)	Neck & back (1.07)	Iron (0.81)	Diabetes (1.26)
Syria	War (8475.03)	IHD (1.32)	Stroke (0.88)	Neck & back (1.05)	Congenital (0.77)	Depression (0.94)	LRI (0.33)	Sense (0.78)	Road injuries (0.37)	Skin (0.78)
Tunisia	Diabetes (1.32)	Neck & back (1.02)	IHD (0.46)	Stroke (0.59)	Depression (1.00)	Sense (0.82)	Road injuries (0.60)	Congenital (0.83)	NN Preterm (0.60)	Lung C (1.21)
Turkey	Neck & back (1.13)	IHD (0.53)	Diabetes (0.97)	Congenital (0.97)	Sense (0.85)	Stroke (0.40)	Depression (0.92)	NN Preterm (0.82)	Lung C (1.18)	Road injuries (0.51)

(Figure 7 continues on next page)

these successes, the last decade has seen profound declines in the burden from malaria and HIV/AIDS. NCD trends have been much more complicated. For the leading cause of disability, low back and neck pain, a lack of knowledge about risks limits the opportunity for prevention. Occupational ergonomic factors and high

body-mass index (BMI) are estimated to be responsible for 30.9% (29.2–32.5) and 5.5% (3.4–7.6) of YLDs due to low back pain, respectively.⁴⁹ The highest occupational risk is found in service industries and manual labour, especially agriculture.^{50,51} The relatively small proportion of low back pain that is caused by high BMI is amenable to

	1	2	3	4	5	6	7	8	9	10
United Arab Emirates	IHD (2-63)	Road injuries (3-09)	Back & neck (0-99)	Diabetes (3-48)	Stroke (2-11)	Depression (1-18)	Other MSK (1-90)	Migraine (1-14)	Skin (0-93)	Drugs (1-93)
Yemen	War (1090-72)	NN Preterm (1-05)	IHD (2-03)	Congenital (1-34)	Road injuries (1-54)	LRI (0-38)	Stroke (1-16)	Diarrhoea (0-30)	Back & neck (1-11)	Diabetes (1-28)
South Asia	IHD (1-18)	NN Enceph (2-15)	NN Preterm (1-07)	LRI (0-74)	COPD (2-04)	Diarrhoea (1-07)	Stroke (0-82)	TB (1-98)	Iron (1-39)	Back & neck (0-91)
South Asia	IHD (1-18)	NN Enceph (2-15)	NN Preterm (1-07)	LRI (0-74)	COPD (2-04)	Diarrhoea (1-07)	Stroke (0-82)	TB (1-98)	Iron (1-39)	Back & neck (0-91)
Bangladesh	Stroke (1-15)	IHD (0-91)	NN Enceph (1-51)	LRI (0-51)	Back & neck (1-06)	Drowning (2-59)	COPD (1-18)	Other MSK (2-60)	Diabetes (0-99)	Sense (0-95)
Bhutan	NN Preterm (1-24)	IHD (0-74)	NN Enceph (1-88)	Back & neck (1-05)	Iron (1-49)	LRI (0-51)	Diarrhoea (1-31)	COPD (1-20)	Diarrhoea (0-87)	Sense (0-98)
India	IHD (1-15)	NN Preterm (1-25)	NN Enceph (2-08)	COPD (2-29)	LRI (0-81)	Diarrhoea (1-31)	Stroke (0-78)	TB (2-41)	Iron (1-60)	Back & neck (0-89)
Nepal	Disaster (1942-61)	LRI (0-58)	IHD (0-89)	NN Enceph (1-36)	TB (1-33)	COPD (1-49)	Back & neck (1-13)	Diarrhoea (0-43)	Stroke (0-49)	Road injuries (0-64)
Pakistan	NN Enceph (2-86)	IHD (1-73)	NN Preterm (0-85)	LRI (0-61)	Diarrhoea (0-88)	Stroke (0-86)	Iron (1-02)	TB (1-32)	Diabetes (1-31)	Other Unint (4-33)
Sub-Saharan Africa	HIV (2-70)	Malaria (5-04)	LRI (0-65)	Diarrhoea (0-68)	NN Sepsis (2-39)	NN Enceph (0-83)	NN Preterm (0-48)	Congenital (0-92)	PEM (1-24)	Meningitis (1-10)
Southern sub-Saharan Africa	HIV (14-25)	LRI (1-74)	TB (10-72)	Diarrhoea (3-65)	Diabetes (2-35)	Road injuries (1-42)	Violence (4-87)	IHD (0-75)	Stroke (0-94)	Back & neck (1-03)
Botswana	HIV (14-16)	TB (15-79)	LRI (1-74)	Diabetes (2-21)	IHD (0-92)	Road injuries (1-10)	Diarrhoea (3-93)	Stroke (1-10)	Self-harm (1-95)	Back & neck (0-98)
Lesotho	HIV (19-51)	Diarrhoea (3-49)	LRI (1-86)	TB (10-10)	NN Preterm (1-09)	Road injuries (1-75)	Other NN (5-08)	Violence (4-64)	IHD (1-21)	Stroke (1-34)
Namibia	HIV (8-67)	LRI (1-38)	Diarrhoea (3-47)	TB (8-56)	NN Preterm (0-76)	Road injuries (1-00)	Other NN (3-68)	IHD (0-67)	NN Enceph (1-09)	Diabetes (1-24)
South Africa	HIV (16-28)	Diabetes (2-67)	Violence (7-21)	LRI (2-39)	TB (17-11)	Road injuries (1-70)	IHD (0-76)	Back & neck (1-04)	Stroke (0-92)	Diarrhoea (7-09)
Swaziland	HIV (21-40)	LRI (2-58)	Diarrhoea (6-52)	Road injuries (2-06)	TB (14-05)	Diabetes (2-54)	IHD (1-17)	Stroke (1-42)	Other NN (3-78)	NN Preterm (0-74)
Zimbabwe	HIV (9-51)	Diarrhoea (2-40)	LRI (1-06)	NN Enceph (1-54)	NN Preterm (0-60)	TB (4-29)	NN Sepsis (2-32)	PEM (3-20)	Stroke (0-93)	Road injuries (0-69)
Western sub-Saharan Africa	Malaria (8-72)	Diarrhoea (0-96)	LRI (0-73)	HIV (1-80)	NN Sepsis (3-59)	NN Enceph (0-98)	NN Preterm (0-52)	Meningitis (1-45)	PEM (1-52)	Congenital (0-86)
Benin	Malaria (11-33)	LRI (0-70)	Diarrhoea (0-46)	NN Sepsis (2-64)	NN Preterm (0-53)	NN Enceph (0-78)	Meningitis (1-20)	PEM (1-43)	Congenital (0-80)	Stroke (0-88)
Burkina Faso	Malaria (4-96)	LRI (0-52)	Diarrhoea (0-45)	NN Sepsis (1-77)	Meningitis (1-00)	NN Enceph (0-71)	PEM (0-87)	Congenital (0-88)	NN Preterm (0-36)	TB (0-73)
Cameroon	HIV (3-90)	Malaria (7-07)	LRI (1-17)	Diarrhoea (1-08)	NN Sepsis (4-62)	NN Enceph (0-98)	NN Preterm (0-53)	Congenital (1-04)	Meningitis (1-91)	PEM (2-64)
Cape Verde	Stroke (0-85)	LRI (0-63)	Back & neck (1-01)	IHD (0-48)	HIV (0-53)	NN Preterm (0-51)	NN Enceph (0-93)	Depression (1-05)	Iron (1-10)	
Chad	Diarrhoea (1-22)	LRI (0-95)	Malaria (3-72)	HIV (1-87)	NN Sepsis (2-61)	NN Preterm (1-67)	NN Enceph (0-64)	Meningitis (1-47)	NN Enceph (0-74)	Congenital (0-90)
Côte d'Ivoire	Malaria (32-28)	HIV (2-73)	LRI (0-92)	NN Sepsis (4-37)	Diarrhoea (0-68)	NN Preterm (0-68)	NN Enceph (1-00)	Congenital (0-93)	Stroke (0-96)	Meningitis (1-15)
The Gambia	LRI (0-36)	NN Sepsis (2-23)	Diarrhoea (0-30)	HIV (0-96)	NN Preterm (0-38)	NN Enceph (0-53)	Meningitis (0-67)	PEM (0-78)	Iron (0-72)	
Ghana	Malaria (81-63)	LRI (0-98)	NN Sepsis (5-51)	HIV (1-20)	NN Preterm (0-50)	Stroke (1-03)	NN Enceph (0-84)	Congenital (0-82)	IHD (0-76)	PEM (2-99)
Guinea	Malaria (6-85)	LRI (0-69)	Diarrhoea (0-37)	NN Sepsis (2-22)	NN Preterm (0-62)	NN Enceph (0-85)	HIV (1-09)	Haemog (3-00)	Meningitis (0-96)	PEM (0-88)
Guinea-Bissau	LRI (0-85)	Diarrhoea (0-88)	HIV (2-52)	Malaria (2-60)	Meningitis (1-80)	NN Sepsis (2-87)	NN Preterm (0-71)	STD (1-74)	STD (3-39)	NN Enceph (0-85)

(Figure 7 continues on next page)

intervention, but the continued escalation of obesity rates indicates that these measures might have little effectiveness. With increasing SDI, the proportion of the workforce in agriculture would be expected to become smaller, which would have some effect on the burden of low back pain. Yet, based on our analyses, nearly 65% of

the burden would remain. The management of most low back and neck pain is largely focused on pain relief and prevention of worsening outcomes through physical therapy and exercise; given the very large burden and the associated economic consequences of lost work time, low back and neck pain should be a priority for research to

	1	2	3	4	5	6	7	8	9	10
Liberia	Malaria (2.88)	LRI (0.46)	Diarrhoea (0.43)	HIV (1.23)	NN Sepsis (1.98)	NN Encephalitis (0.68)	NN Preterm (0.48)	TB (1.07)	Ebola (58898.20)	Meningitis (0.78)
Mali	Malaria (5.95)	Diarrhoea (0.40)	PEM (1.48)	NN Preterm (0.87)	NN Sepsis (2.48)	LRI (0.29)	NN Encephalitis (0.91)	Meningitis (0.95)	HIV (1.07)	Congenital (0.82)
Mauritania	LRI (0.60)	NN Sepsis (3.54)	Diarrhoea (0.53)	NN Preterm (0.50)	NN Encephalitis (0.81)	Iron (1.25)	Congenital (0.89)	Back & neck (1.07)	STD (2.22)	Meningitis (0.89)
Niger	Malaria (1.30)	Diarrhoea (0.54)	LRI (0.39)	Meningitis (1.09)	PEM (0.54)	NN Preterm (0.52)	NN Sepsis (1.02)	NN Encephalitis (0.54)	Congenital (0.60)	TB (0.55)
Nigeria	Malaria (104.66)	Diarrhoea (2.09)	HIV (2.16)	LRI (1.00)	NN Sepsis (5.39)	NN Encephalitis (1.25)	Haemog (6.32)	NN Preterm (0.48)	Congenital (0.90)	Meningitis (2.19)
São Tomé and Príncipe	LRI (0.67)	NN Sepsis (2.33)	Congenital (0.83)	Stroke (0.97)	Diarrhoea (0.43)	NN Encephalitis (0.61)	NN Preterm (0.27)	IHD (0.63)	Back & neck (1.04)	CKD (2.03)
Senegal	Diarrhoea (0.52)	LRI (0.42)	NN Sepsis (2.04)	NN Encephalitis (0.59)	NN Preterm (0.39)	TB (1.21)	Congenital (0.77)	Meningitis (0.95)	Iron (1.04)	Malaria (1.89)
Sierra Leone	Malaria (14.10)	LRI (0.90)	Haemog (7.62)	Diarrhoea (0.61)	Ebola (97400.61)	NN Sepsis (2.97)	NN Encephalitis (1.00)	NN Preterm (0.66)	Meningitis (1.60)	HIV (1.15)
Togo	Malaria (17.88)	HIV (2.21)	LRI (0.68)	Diarrhoea (0.54)	NN Sepsis (3.19)	NN Encephalitis (0.88)	NN Preterm (0.55)	Congenital (0.80)	Haemog (3.11)	Stroke (0.83)
Eastern sub-Saharan Africa		HIV (2.56)	LRI (0.58)	Diarrhoea (0.54)	Malaria (2.27)	Congenital (1.03)	NN Encephalitis (0.71)	NN Preterm (0.42)	NN Sepsis (1.49)	PEM (1.04)
Burundi	LRI (0.44)	Diarrhoea (0.43)	Malaria (1.33)	NN Preterm (0.54)	Congenital (1.01)	PEM (0.93)	NN Encephalitis (0.62)	HIV (0.90)	TB (1.00)	Meningitis (0.64)
Comoros	LRI (0.50)	Diarrhoea (0.43)	Congenital (0.94)	NN Preterm (0.43)	NN Encephalitis (0.64)	NN Sepsis (1.53)	Iron (0.86)	TB (0.84)	Stroke (0.61)	STD (1.68)
Djibouti	LRI (1.11)	Diarrhoea (1.08)	HIV (1.03)	Congenital (1.26)	Stroke (0.83)	Iron (1.32)	NN Preterm (0.51)	IHD (0.70)	TB (1.54)	PEM (2.98)
Eritrea	Diarrhoea (0.82)	LRI (0.67)	PEM (1.52)	Congenital (0.98)	Iron (1.32)	TB (1.24)	NN Preterm (0.43)	NN Encephalitis (0.56)	Meningitis (0.91)	Stroke (0.97)
Ethiopia	LRI (0.43)	Diarrhoea (0.36)	TB (1.16)	Congenital (0.97)	NN Encephalitis (0.69)	HIV (0.83)	NN Preterm (0.36)	NN Sepsis (1.42)	Meningitis (0.76)	STD (1.74)
Kenya	HIV (3.70)	Diarrhoea (1.08)	LRI (0.75)	NN Encephalitis (0.96)	NN Preterm (0.46)	Congenital (0.86)	Iron (1.30)	Malaria (2.40)	Other NTD (3.37)	Meningitis (1.07)
Madagascar	LRI (0.75)	Diarrhoea (0.78)	NN Preterm (0.52)	PEM (2.24)	Stroke (1.21)	Congenital (0.88)	Iron (1.15)	Malaria (4.00)	NN Sepsis (1.79)	IHD (0.86)
Malawi	HIV (5.07)	Malaria (5.01)	LRI (0.60)	Diarrhoea (0.53)	NN Encephalitis (0.72)	NN Preterm (0.49)	Congenital (1.00)	PEM (1.09)	Meningitis (0.96)	NN Sepsis (1.32)
Mozambique	HIV (7.73)	Malaria (4.06)	LRI (0.31)	Diarrhoea (0.29)	NN Encephalitis (0.72)	Congenital (0.98)	NN Sepsis (1.33)	NN Preterm (0.37)	TB (0.92)	Stroke (0.86)
Rwanda	LRI (0.79)	HIV (1.15)	Congenital (1.23)	Diarrhoea (0.42)	NN Encephalitis (0.50)	NN Sepsis (0.75)	PEM (1.65)	NN Sepsis (2.01)	Road injuries (0.98)	Meningitis (1.07)
Somalia	Diarrhoea (0.68)	LRI (0.64)	PEM (0.83)	TB (1.18)	NN Preterm (0.64)	Iron (1.47)	Malaria (0.32)	Congenital (0.81)	Meningitis (0.59)	War (187.71)
South Sudan	Diarrhoea (0.93)	LRI (0.72)	HIV (1.90)	Malaria (2.01)	PEM (1.38)	Meningitis (1.18)	NN Preterm (0.54)	TB (1.10)	Congenital (0.85)	Iron (1.21)
Tanzania	HIV (2.57)	LRI (0.78)	Congenital (1.38)	Malaria (11.61)	Diarrhoea (0.52)	NN Encephalitis (0.69)	Iron (1.28)	NN Preterm (0.34)	NN Sepsis (2.06)	PEM (1.75)
Uganda	HIV (2.88)	LRI (0.60)	Malaria (9.47)	Diarrhoea (0.50)	NN Encephalitis (0.87)	Congenital (1.12)	NN Preterm (0.46)	Meningitis (1.45)	NN Sepsis (1.87)	Haemog (3.24)
Zambia	HIV (6.46)	LRI (0.96)	Malaria (39.51)	Diarrhoea (1.19)	Congenital (1.05)	NN Encephalitis (0.75)	Meningitis (1.93)	NN Preterm (0.33)	IHD (1.03)	
Central sub-Saharan Africa		Malaria (3.30)	LRI (0.55)	Diarrhoea (0.35)	HIV (1.53)	PEM (1.04)	NN Preterm (0.46)	Congenital (0.89)	NN Sepsis (1.62)	NN Encephalitis (0.64)
Angola	LRI (0.95)	Malaria (25.93)	Malaria (0.66)	Diarrhoea (1.35)	Congenital (1.09)	NN Sepsis (2.85)	PEM (2.45)	NN Preterm (0.38)	Meningitis (1.62)	Road injuries (1.43)
Central African Republic	Malaria (6.71)	HIV (4.23)	LRI (0.95)	Diarrhoea (0.68)	TB (1.82)	NN Preterm (0.73)	Stroke (1.55)	PEM (1.67)	Road injuries (2.01)	NN Sepsis (2.17)
Congo (Brazzaville)	HIV (3.37)	Malaria (109.15)	LRI (0.97)	Congenital (1.04)	NN Sepsis (3.14)	Stroke (0.84)	Diarrhoea (0.84)	NN Encephalitis (0.83)	IHD (0.85)	NN Preterm (0.42)
DR Congo	Malaria (2.57)	LRI (0.46)	Diarrhoea (0.28)	HIV (1.28)	PEM (0.86)	Other NTD (3.84)	NN Encephalitis (0.63)	NN Preterm (0.48)	Congenital (0.81)	NN Sepsis (1.30)
Equatorial Guinea	Malaria (1125.32)	HIV (3.81)	LRI (1.87)	NN Sepsis (6.10)	Congenital (1.39)	NN Preterm (0.73)	Diabetes (1.74)	Road injuries (1.05)	NN Encephalitis (1.22)	Other NN (2.72)
Gabon	HIV (2.84)	Malaria (968.41)	LRI (1.60)	Congenital (1.23)	NN Sepsis (5.65)	Stroke (1.03)	Diabetes (1.51)	Iron (1.90)	Road injuries (0.93)	IHD (0.61)

■ 0-0.63 ■ 0.63-0.79 ■ 0.79-0.90 ■ 0.90-1.00 ■ 1.00-1.09 ■ 1.09-1.24 ■ 1.24-1.56 ■ 1.56-2.49 ■ >2.49

identify more effective preventive and therapy measures.⁵² Similarly, despite broad decreases in age-standardised rates of injury burden, the pace of progress for these causes has been comparatively slow and ultimately has led to minimal changes in the proportion of overall burden due to injuries during the past 25 years. Prevention of injuries requires strong public safety policies,⁵³ but minimising mortality and long-term disability from injuries hinges upon having comprehensive trauma care systems^{54,55} that provide timely, evidence-based care,⁵⁶⁻⁵⁸ including emergency surgical services.^{59,60}

In 2015, sense organ disorders were the second-leading cause of YLDs and resulted in more than 68 million DALYs. Reducing DALYs from vision impairment is achievable through vertically integrated programmes, including the delivery of eyeglasses for refractive error, curative surgery for cataracts, and onchocerciasis and trachoma prevention. Given the availability of cost-effective interventions, greater policy attention is needed for vision loss burden. Although interventions for hearing loss are less clear, the use of timely antibiotics for otitis media and meningitis and the provision of hearing aids for conductive hearing loss are likely to reduce its burden.

Reductions in age-standardised DALY rates due to some NCDs such as cardiovascular diseases, most cancers, chronic respiratory diseases, and many digestive diseases—some of which can be attributed to reductions in risk factors such as tobacco and improvements in cause-specific treatment and event survival—mask the effects of population ageing. This means more people had disease burden from these causes and total DALYs have remained largely unchanged (eg, chronic obstructive pulmonary disease) or significantly increased (eg, cardiovascular diseases, cancers, neurological disorders, diabetes, chronic kidney disease, and musculoskeletal disorders such as osteoarthritis and low back and neck pain) over time. As demographic transitions are widely expected to continue, the burden from NCDs is likely to continue expanding. Widespread efforts must continue to enact societal and environmental policies to reduce risk factor exposure, while national and local health systems must adapt to meet the prevention, screening, and treatment needs of their populations. As we now recognise many of the risk factors related to NCDs, low-SDI and middle-SDI countries could adopt policies to circumvent the mistakes made by other countries as they progressed along the SDI spectrum.

Mental and substance use disorders are a particularly challenging group of conditions with non-trivial levels of disease burden in all geographies. Some countries provide excellent mental health resources, whereas others, particularly LMICs, do not have screening or treatment programmes. Addressing the growing burden and disparity in mental health disorders will be an especially pressing challenge during the SDG era.

A number of emerging and growing health threats also deserve special attention in policy planning, including infectious diseases such as outbreaks of dengue fever, Ebola virus disease, Zika virus, and pandemic influenza, and antimicrobial-resistant pathogens, which represent acute threats to life and highlight health-system deficiencies where they occur; substance abuse disorders, particularly of opioids and cocaine, in eastern Europe, Australia, and North America; and intentional firearm injuries, especially in Latin America, the USA, and South Africa. In the case of dengue fever and potentially other yet to be identified infectious diseases, urbanisation and global environmental change have contributed to an increased incidence and future climate change scenarios depict a rising trend in the coming years.⁶¹ Other emerging infectious diseases, including Zika and chikungunya viruses, have yet to be comprehensively analysed by the GBD.

A major change in the GBD 2015 assessment has been the closer integration of the assessments of mortality and disease sequelae prevalence in modelling. For cancers, HIV/AIDS, and injuries, previous iterations of the GBD modelled mortality, incidence, and prevalence in a coherent manner. For some other diseases, the modelling of disease prevalence and cause-specific mortality rates largely used different data sources and modelling techniques. Independent estimation of prevalence and mortality led in some cases to patterns across locations of excess mortality rates that were not consistent with expected relationships related to health-system access. For GBD 2015, we built the modelling of cause-specific mortality, excess mortality, incidence, and prevalence into nearly every cause. The effect of this approach has led to increases in the number of DALYs from injuries due to YLDs and changes in prevalence for other conditions, particularly those with minimal data for prevalence or incidence. More attention will be paid in future iterations of the GBD study to identifying unpublished data from cohort studies or linkage studies

Figure 7: Leading ten causes of DALYs with the ratio of observed DALYs to DALYs expected on the basis of SDI in 2015, by location

The ratio of observed DALYs to DALYs expected based on SDI is provided in brackets for each cause and cells are colour coded by ratio ranges (calculated to place a roughly equal number of cells into each bin). Shades of blue represent much lower observed DALYs than expected levels based on SDI, whereas red shows observed DALYs that exceed expected levels. IHD=ischaemic heart disease. Back & neck=low back and neck pain. Diabetes=diabetes mellitus. Stroke=Cerebrovascular disease. Lung C=lung, bronchus, and trachea cancers. Sense=sense organ diseases. Depression=depressive disorders. Alzheimer's=Alzheimer's disease and other dementias. Oth MSK=other musculoskeletal disorders. COPD=chronic obstructive pulmonary disease. NN Preterm=neonatal preterm birth complications. Diarrhoea=diarrhoeal diseases. Skin=skin and subcutaneous diseases. NN Enceph=neonatal encephalopathy due to birth asphyxia and trauma. Drugs=drug use disorders. Congenital=congenital anomalies. Liver C=liver cancer. Stomach C=stomach cancer. CKD=chronic kidney disease. Anxiety=anxiety disorders. Iron=iron-deficiency anaemia. HIV=HIV/AIDS. Colorect C=colon and rectum cancer. LRI=lower respiratory infections. Cirr HepC=cirrhosis due to hepatitis C. CMP=cardiomyopathy and myocarditis. Cirr alc=cirrhosis due to alcohol use. Other Cardio=other cardiovascular and circulatory diseases. Alcohol=alcohol use disorders. Violence=interpersonal violence. HTN HD=hypertensive heart disease. Haemog=haemoglobinopathies and haemolytic anaemias. TB=tuberculosis. Prostate C=prostate cancer. War=collective violence and legal intervention. Other Unint=other unintentional injuries. Oth trans=other transport injuries. Other NN=Other neonatal disorders. NN Sepsis=neonatal sepsis and other neonatal infections. PEM=protein-energy malnutrition. STD=sexually transmitted diseases excluding HIV. Ebola=Ebola virus disease. Other NTD=other neglected tropical diseases. GBD=Global Burden of Disease. SDI=Socio-demographic Index. DALYs=disability-adjusted life-years.

on levels of excess mortality by age and sex, especially in LMICs, to further strengthen modelling efforts.

A major development for the GBD 2015 has been adaptation of the GATHER guidelines endorsed by WHO, the Institute for Health Metrics and Evaluation (IHME), and other organisations.^{22,23} GATHER compliance, including the sharing of statistical code for each of the many analytical steps in the GBD, provides a new level of transparency for the overall enterprise. We expect that many researchers will want to investigate, propose improvements, and provide alternative assessments for many components of the GBD. We welcome the debate that will follow on the best way to analyse different components of the GBD. We believe enhanced transparency will lead to healthy debate and exchange and to improved methods, data, and results for many aspects of the GBD. Transparency will not necessarily lead to consensus but it will broaden everyone's understanding of the available evidence on descriptive epidemiology. Adoption by the GBD of the GATHER guidelines will hopefully stimulate other organisations to adopt the guidelines in all aspects of their work as well.

Although the volume of input data to the GBD has continued to increase substantially, major data gaps remain.^{12,20,62,63} Geographical and temporal coverage of all-cause and cause-specific mortality datasets are variable, as is the quality of the data contained in such systems. Development of methods to report overall evidence grades for each outcome–location–year combination would be valuable to help to guide strategies for improving data quality and closing data gaps. Investing to develop and improve cause of death and vital registration systems is crucial to improve the quality of insights from the GBD; incorporation of existing data from existing and new collaborators that are not currently in the GBD is another important aspect of this effort. Several countries have experienced significant recent turmoil, especially armed conflict in Syria, Yemen, and other countries in north Africa and the Middle East. Burden from many conditions is believed to have increased during and following those events, but due to disruption of data-collection systems, the full effect of such events has been difficult to quantify. For non-fatal health outcomes, some of the most notable data gaps pertain to aspects of individual diseases and injuries that are not typically included as part of standard epidemiological reports such as distribution of symptoms for those with chronic conditions at various stages of illness, duration of disability following acute events, or long-term disability after major acute injuries. Therefore, our recommendations regarding data gaps pertaining to non-fatal health loss are twofold. First, reports and scientific journals should strive to include reporting on functional health status including severity, distribution, and duration of symptoms with all epidemiological studies. Second, countries should work to centralise and compile existing non-fatal health data and invest to collect population-level epidemiological data on important causes of YLDs.

The iterative and now annual cycle of the GBD revisions provide opportunities to improve the estimation or scope of the GBD. Due to the high interest in Zika virus, we believe that we should try to quantify the burden related to Zika virus disease in the GBD 2016 analysis. Given the focus in the SDGs on various forms of sexual violence, we believe careful investigation of the evidence base for estimation is warranted. As noted in the GBD papers on mortality and non-fatal outcomes,^{12,20} there are also a number of opportunities to improve data processing and estimation methods that will be explored for the next cycle of estimation. We also expect to include more subnational analyses, particularly for large countries.

Our analysis has several limitations. First and foremost, the calculation of DALYs and HALE reflects the limitations of all the underlying analyses of the GBD, including all-cause mortality, cause-specific mortality, prevalence, incidence, disability-weight derivation, and simulation of comorbidity. Second, as discussed above, data limitations are apparent in a number of facets of our analysis. Third, inherent to the GBD approach is the effort to quantify specific sequelae of each disease and injury. This means that the full disease burden of certain conditions such as heart failure, anaemia, vision and hearing loss, infertility, epilepsy, and intellectual disability are not as readily apparent in high-level review of the GBD results. However, the YLDs for these impairments are reported elsewhere.²⁰ Fourth, our analysis of the relationship between SDI, DALYs, and HALE reflects the average historical relationship between SDI and each measure, so despite often strong correlation with SDI it cannot be interpreted as being causal in nature. In some cases, association of SDI with health indicators could be considered a confounder when the same elements (education, income, and fertility) are used to develop both the index and as a covariate in cause-specific models. SDI utility might be improved in the future through consideration of additional societal elements such as inequality in each component. Other measures that capture the status of women in society, such as the female labour-force participation, could be considered in future revisions. Fifth, we have assumed independence of uncertainty between YLLs and YLDs as well as between YLDs and life expectancy. Empirical evidence to guide alternative assumptions, however, is currently very limited. Sixth, recent events in Syria and Libya and the resulting mass migration have led to considerable health loss, including drownings of many migrants. New migrants have different health problems than the populations of the countries to which they have moved. Both the drownings and the change in health status in countries receiving migrants are not adequately captured in this assessment due to the time-lags in data collection and data capture intrinsic in all health data systems. Seventh, estimates of expected burden from SDI alone are based on the average levels of burden at each level of SDI. For endemic diseases, comparisons of observed rates to expected rates will lead to high observed over expected ratios in endemic countries.

and low ratios in non-endemic countries. Interpretation of the ratios for conditions that are endemic in only some countries needs to take this into account.

WHO has estimated DALYs by cause for the single years of 2000 and 2012.^{13,17} They used published GBD 2010 results used to generate WHO 2012 DALYs for 132 causes with some modifications. First, WHO life tables were used instead of GBD life tables.¹⁴ WHO life tables are different from the UN Population Division life tables and use a set of methods developed by Murray and colleagues in the late 1990s;^{11,64} this approach does not benefit from the many improvements in data processing and estimation methods that have emerged in the last 15 years.^{12,65,66} Second, WHO altered the empirical disability weights, which were derived from an international sample of more than 60 000 respondents from the GBD analysis²⁶ for 32 outcomes using the opinions of 45 respondents.^{13,67} Third, WHO calculated YLLs after changing from the GBD 2010 normative standard life expectancy of 86·0 years to 91·9 years.⁶⁸ Fourth, rather than using GBD results, WHO elected to use alternative estimates for 12 causes of death, including tuberculosis, HIV/AIDS and other sexually transmitted infections, malaria, whooping cough, measles, schistosomiasis, maternal disorders, cancers, alcohol and drug use disorders, epilepsy, conflict and natural disasters, and road traffic accidents.¹³ Finally, WHO substituted prevalence estimates produced internally for vision loss, hearing loss, intellectual disability, infertility, anaemia, back pain, alcohol use disorders, headache, and skin diseases. The final hybrid estimates of DALYs do not provide uncertainty measures and have not been peer-reviewed.

WHO has also produced HALE estimates for three time periods, 2000, 2012, and 2015, using GBD 2010 results as described above for 2000 and 2012, and GBD 2013 results for 2015, also without uncertainty measures.¹⁴ Differences between their estimates and those for HALE from GBD 2015 reflect changes in age-specific YLDs per capita from GBD 2013 to GBD 2015 and differences in WHO life expectancy (appendix p 85). The EC and the OECD also report healthy life expectancy estimates based on self-reported health status from 2004 through 2014, but without specific consideration of prevalence or incidence of disease.^{52,53} Comparison of 2014 estimates from the EC and GBD mostly show lower estimates from EC (appendix pp 86–87). EC estimates also report much wider ranges in HALE across countries in Europe than those estimated through GBD. Further, in a number of countries, EC estimates point to lower HALE among women than for men. These differences, both in terms of absolute estimates and those by sex, are probably due to the inclusion of non-health factors in self-reported assessments of disability.

In conclusion, HALE has increased steadily throughout the world over the MDG era, with a concomitant decrease in age-standardised DALY rates from most conditions. Declines occurred in overall health loss due to many communicable, maternal, neonatal, and nutritional

diseases. Much of the evolution of health is consistent with the expected changes in disease burden with development that have been quantified in this study. Substantial variation in burden compared with levels expected on the basis of SDI suggests wide heterogeneity in the ability of governments and health systems to adequately meet the health needs of their populations. Progress in reducing these gaps will be crucial to achieving the ambitious SDG agenda. Demographic changes leading to increased population size and older average age have offset otherwise important gains in age-specific DALY rates leading to rising burden on health systems for many ailments of ageing. Emerging health threats and causes with lagging progress should be viewed as essential foci for investment in health infrastructure and health data systems to improve the global community's insights into the aggregate quality of care and the overall health of populations.

GBD 2015 DALYs and HALE Collaborators

Nicholas J Kassebaum, Megha Arora, Ryan M Barber, Zulfiqar A Bhutta, Jonathan Brown, Austin Carter, Daniel C Casey, Fiona J Charlson, Matthew M Coates, Megan Coggeshall, Leslie Cornaby, Lalit Dandona, Daniel J Dicker, Holly E Erskine, Alize J Ferrari, Christina Fitzmaurice, Kyle Foreman, Mohammad H Forouzanfar, Nancy Fullman, Peter W Gething, Ellen M Goldberg, Nicholas Graetz, Juanita A Haagsma, Simon I Hay, Catherine O Johnson, Laura Kemmer, Ibrahim A Khalil, Yohannes Kinfu, Michael J Kutz, Hmwe H Kyu, Janni Leung, Xiaofeng Liang, Stephen S Lim, Rafael Lozano, George A Mensah, Joe Mikesell, Ali H Mokdad, Meghan D Mooney, Mohsen Naghavi, Grant Nguyen, Elaine Nsoesie, David M Pigott, Christine Pinho, Zane Rankin, Nikolas Reinig, Joshua A Salomon, Logan Sandar, Alison Smith, Reed J D Sorenson, Jeffrey Stanaway, Caitlyn Steiner, Stephanie Teeple, Chris Troeger, Thomas Truelsen, Amelia VanderZanden, Joseph A Wagner, Valentine Wanga, Harvey A Whiteford, Maigeng Zhou, Leo Zoekler, Amanuel Alemu Ababior*, Kalkidan Hassen Abate*, Cristiana Abbafati*, Kaja M Abbas*, Foad Abd-Allah*, Biju Abraham*, Ibrahim Abubakar*, Laith J Abu-Raddad*, Niveen M E Abu-Rmeileh*, Tom Achoki*, Ilana N Ackerman*, Akinede Olupelumi Adebiyi*, Isaac Akinkunmi Adedeji*, José C Adsuar*, Kossivi Agbelenko Afanvi*, Ashkan Afshin*, Emilie Elisabet Agardh*, Arnav Agarwal*, Sanjay Kumar Agarwal*, Muktar Beshir Ahmed*, Aliasghar Ahmad Kiadaliri*, Hamid Ahmadieh*, Nadia Akseer*, Ziyad Al-Aly*, Khurshid Alam*, Noore K M Alam*, Saleh Fahed Aldhahri*, Miguel Angel Alegretti*, Alicia V Aleman*, Zewdie Aderaw Alemu*, Lily T Alexander*, Raghib Ali*, Ala'a Alkerwi*, François Alla*, Peter Allebeck*, Christine Allen*, Ubai Alsharif*, Khalid A Altirkawi*, Elena Alvarez Martin*, Nelson Alvis-Guzman*, Azmeraw T Amare*, Alemayehu Amberbir*, Adelazza Kofi Amegah*, Heresh Amini*, Walid Ammar*, Stephen Marc Amrock*, Gregory M Anderson*, Benjamin O Anderson*, Carl Abelardo T Antonio*, Palwasha Anvari*, Johan Ärnlöv*, Valentina S Arsic Arsenijevic*, Al Artaman*, Hamid Asayesh*, Rana Jawad Asghar*, Euripide Frinel G Arthur Avokpaho*, Ashish Awasthi*, Beatriz Paulina Ayala Quintanilla*, Peter Azzopardi*, Umar Bacha*, Alaa Badawi*, Kalpana Balakrishnan*, Amitava Banerjee*, Aleksandra Barac*, Suzanne L Barker-Collo*, Till Bärnighausen*, Lars Barregard*, Lope H Barrero*, Sanjay Basu*, Tigist Assefa Bayou*, Justin Beardsley*, Neeraj Bedi*, Ettore Beghi*, Brent Bell*, Michelle L Bell*, Corina Benjet*, Derrick A Bennett*, Isabela M Bensenor*, Adugnaw Berhane*, Eduardo Bernabe*, Balem Demtsu Betsu*, Addisu Shunu Beyene*, Neeraj Bhala*, Anil Bhanzali*, Samir Bhatt*, Sibhata Biadgilign*, Kelly Bienhoff*, Boris Bikbov*, Aref A Bin Abdulhak*, Stan Biryukov*, Donal Bisanzio*, Espen Bjertness*, Jed D Blore*, Rohan Borschmann*, Soufiane Boufous*, Rupert R A Bourne*, Michael Brainin*, Alexandra Brazinova*, Nicholas J K Breitborde*, Traolach S Brugha*,

- Rachelle Buchbinder*, Geoffrey Colin Buckle*, Zahid A Butt*, Bianca Calabria*, Ismael Ricardo Campos-Nonato*, Julio Cesar Campuzano*, Hélène Carabin*, Jonathan R Carapetis*, Rosario Cárdenas*, Juan Jesus Carrero*, Carlos A Castañeda-Orjuela*, Jacqueline Castillo Rivas*, Ferrán Catalá-López*, Fiorella Cavalieri*, Jung-Chen Chang*, Peggy Pei-Chia Chiang*, Mirriam Chibalabala*, Chioma Ezinne Chibueze*, Vesper Hichilombwe Chisumpa*, Jee-Young Jasmine Choi*, Lincoln Choudhury*, Hanne Christensen*, Liliana G Ciobanu*, Valentina Colistro*, Mercedes Colomar*, Samantha M Colquhoun*, Monica Cortinovis*, John A Crump*, Alberto Damasceno*, Rakhi Dandona*, Paul I Dargan*, José das Neves*, Gail Davey*, Adrian C Davis*, Diego De Leo*, Louisa Degenhardt*, Liana C Del Gobbo*, Sarah Derrett*, Don C Des Jarlais*, Gabrielle A deVeber*, Samath D Dharmaratne*, Preet K Dhillon*, Eric L Ding*, Kerrie E Doyle*, Tim R Driscoll*, Leilei Duan*, Manisha Dubey*, Bruce Bartholow Duncan*, Hedyeh Ebrahimi*, Richard G Ellenbogen*, Iqbal Elyazar*, Aman Yesuf Endriess*, Sergey Petrovich Ermakov*, Babak Eshratii*, Alireza Esteghamati*, Kara Estep*, Saman Fahimi*, Talha A Farid*, Carla Sofia e Sa Farinha*, André Faro*, Maryam S Farvid*, Farshad Farzadfar*, Valery L Feigin*, Seyed-Mohammad Fereshtehnejad*, Jefferson G Fernandes*, Joao C Fernandes*, Florian Fischer*, Joseph R A Fitchett*, Nataliya Foigt*, F Gerry R Fowkes*, Richard C Franklin*, Joseph Friedman*, Joseph Frostad*, Thomas Fürst*, Neal D Futran*, Belinda Gabbe*, Fortuné Gbètoho Gankpé*, Alberto L Garcia-Basteiro*, Tsegaye Tewelde Gebrehiwot*, Amanuel Tesfay Gebremedhin*, Johanna M Geleinse*, Katherine B Gibney*, Richard F Gillum*, Ibrahim Abdelmageem Mohamed Ginawi*, Ababi Zergaw Giref*, Maurice Giroud*, Melkamu Dedefo Gishu*, Giorgia Giussani*, William W Godwin*, Hector Gomez-Dantes*, Philimon Gona*, Amador Goodridge*, Sameer Vali Gopalani*, Carolyn C Gotay*, Atsushi Goto*, Hebe N Gouda*, Harish Gugnani*, Yuming Guo*, Rahul Gupta*, Rajeev Gupta*, Vinip Gupta*, Reyna A Gutiérrez*, Nima Hafezi-Nejad*, Demewoz Haile*, Alemanyehu Desalegne Hailu*, Gessessew Bugssa Hailu*, Yara A Halasa*, Randah Ribhi Hamadeh*, Samer Hamidi*, Mouhanad Hammami*, Alexis J Handal*, Graeme J Hankey*, Hilda L Harb*, Sivadasanpillai Harikrishnan*, Josep Maria Haro*, Mohammad Sadegh Hassanvand*, Tahis Ahmed Hassen*, Rasmus Havmoeller*, Roderick J Hay*, Mohammad T Hedayati*, Ileana Beatriz Heredia-Pi*, Pouria Heydarpour*, Hans W Hoek*, Daniel J Hoffman*, Masako Horino*, Nobuyuki Horita*, H Dean Hosgood*, Damian G Hoy*, Mohamed Hsairi*, Hsiang Huang*, John J Huang*, Kim Moesgaard Ibburg*, Bulat T Idrisov*, Kaire Innos*, Manami Inoue*, Kathryn H Jacobsen*, Alejandra Jauregui*, Achala Upendra Jayatileke*, Panniyammakal Jeemon*, Vivekanand Jha*, Guohong Jiang*, Ying Jiang*, Tariku Jibat*, Aida Jimenez-Corona*, Yi Jin*, Jost B Jonas*, Zubair Kabir*, Dan K Kajungu*, Yogeshwar Kalkonde*, Ritul Kamal*, Haidong Kan*, Amit Kandel*, André Karch*, Corine Kakizi Karema*, Chante Karimkhani*, Amir Kaseaeian*, Marzieh Katibeh*, Anil Kaul*, Norito Kawakami*, Dhruv S Kazi*, Peter Njenga Keiyoro*, Andrew Haddon Kemp*, Andre Pascal Kengne*, Andre Keren*, Chandrasekharan Nair Kesavachandran*, Yousef Saleh Khader*, Abdur Rahman Khan*, Ejaz Ahmad Khan*, Young-Ho Khang*, Tawfiq Ahmed Muthafee Khoja*, Jagdish Khubchandani*, Christian Kieling*, Cho-il Kim*, Daniel Kim*, Yun Jin Kim*, Niranjan Kissoon*, Miia Kivipelto*, Luke D Knibbs*, Ann Kristin Knudsen*, Yoshihiro Kokubo*, Dhaval Kolte*, Jacek A Kopec*, Parvaiz A Koul*, Ai Koyanagi*, Barthelemy Kuate Defo*, Ricardo S Kuchenbecker*, Burcu Kucuk Bicer*, Ernst J Kuipers*, G Anil Kumar*, Gene F Kwan*, Ratilal Laloo*, Tea Lallukka*, Anders Larsson*, Asma Abdul Latif*, Pablo M Lavados*, Alicia Elena Beatriz Lawrynowicz*, Janet L Leasher*, James Leigh*, Ricky Leung*, Yichong Li*, Yongmei Li*, Steven E Lipshultz*, Patrick Y Liu*, Yang Liu*, Belinda K Lloyd*, Giancarlo Logroscino*, Katharine J Looker*, Paulo A Lotufo*, Robyn M Lucas*, Raimundas Lunevicius*, Ronan A Lyons*, Hassan Magdy Abd El Razek*, Mahdi Mahdavi*, Marek Majdan*, Azeem Majeed*, Reza Malekzadeh*, Deborah Carvalho Malta*, Wagner Marcenes*, Jose Martinez-Raga*, Felix Masiye*, Amanda J Mason-Jones*, Richard Matzopoulos*, Bongani M Mayosi*, John J McGrath*, Martin McKee*, Peter A Meaney*, Alem Mehari*, Yohannes Adama Melaku*, Peter Memiah*, Ziad A Memish*, Walter Mendoza*, Atte Meretoja*, Tuomo J Meretoja*, Yonatan Moges Mesfin*, Francis Apolinario Mhimbira*, Anoushka Millear*, Ted R Miller*, Edward J Mills*, Mojde Mirarefin*, Erkin M Mirrakhimov*, Philip B Mitchell*, Charles N Mock*, Karzan Abdulmuhsin Mohammad*, Alireza Mohammadji*, Shafiu Mohammed*, Lorenzo Monasta*, Julio Cesar Montañez Hernandez*, Marcella Montico*, Maziar Moradi-Lakeh*, Rintaro Mori*, Ulrich O Mueller*, John Everett Mumford*, Michele E Murdoch*, Gudlavalleti Venkata Satyanarayana Murthy*, Jean B Nachega*, Aliya Naheed*, Luigi Naldi*, Vinay Nangia*, John N Newton*, Marie Ng*, Frida Namnyam Ngalesoni*, Quyen Le Nguyen*, Muhammad Imran Nisar*, Patrick Martial Nkamedje Pete*, Joan M Nolla*, Ole F Norheim*, Rosana E Norman*, Bo Norrvig*, Carla Makhlouf Obermeyer*, Felix Akpajene Ogbo*, In-Hwan Oh*, Olanrewaju Oladimeji*, Pedro R Olivares*, Bolajoko Olubukunola Olusanya*, Jacob Olusegun Olusanya*, Eyal Oren*, Alberto Ortiz*, Erika Ota*, Abayomi Samuel Oyekale*, Mahesh PA*, Eun-Kee Park*, Mahboubeh Parsaeian*, Scott B Patten*, George C Patton*, João Mário Pedro*, David M Pereira*, Norberto Perico*, Konrad Pesudovs*, Max Petzold*, Michael Robert Phillips*, Frédéric B Piel*, Julian David Pillay*, Farhad Pisghar*, Dietrich Plass*, Suzanne Polinder*, Svetlana Popova*, Richie G Poulton*, Farshad Pourmalek*, Noela M Prasad*, Mostafa Qorbani*, Rynaz H S Rabiee*, Amir Radfar*, Anwar Rafay*, Kazem Rahimi*, Vafa Rahimi-Movaghhar*, Mahfuzar Rahman*, Mohammad Hifz Ur Rahman*, Sajjad Ur Rahman*, Dheeraj Rai*, Rajesh Kumar Rai*, Sasa Rajsic*, Murugesan Raju*, Usha Ram*, Kavitha Ranganathan*, Amany H Refaat*, Marissa B Reitsma*, Giuseppe Remuzzi*, Serge Resnikoff*, Alex Reynolds*, Antonio L Ribeiro*, Stefano Ricci*, Hirbo Shore Roba*, David Rojas-Rueda*, Luca Ronfani*, Gholamreza Roshandel*, Gregory A Roth*, Ambuj Roy*, Ben Benasco Sackey*, Rajesh Sagari*, Juan R Sanabria*, Maria Dolores Sanchez-Niño*, Itamar S Santos*, João Vasco Santos*, Rodrigo Sarmiento-Suarez*, Benn Sartori*, Maheswar Satpathy*, Miljoje Savic*, Monika Sawhney*, Maria Inês Schmidt*, Ione J C Schneider*, Aletta E Schutte*, David C Schwebel*, Soraya Seedat*, Sadaf G Sepanlou*, Edson E Servan-Mori*, Saeid Shahraz*, Masood Ali Shaikh*, Rajesh Sharma*, Jun She*, Sara Sheikhbahaei*, Jiabin Shen*, Kevin N Sheth*, Kenji Shibuya*, Mika Shigematsu*, Min-Jeong Shin*, Rahman Shiri*, Inga Dora Sigfusdottir*, Diego Augusto Santos Silva*, Jonathan I Silverberg*, Edgar P Simard*, Abhishek Singh*, Jasvinder A Singh*, Prashant Kumar Singh*, Vegard Skirbekk*, Jens Christoffer Skogen*, Michael Soljak*, Kjetil Søredeide*, Reed J D Sorensen*, Chandrashekhar T Sreeramaredy*, Vasiliki Stathopoulou*, Nicholas Steel*, Dan J Stein*, Murray B Stein*, Timothy J Steiner*, Lars Jacob Stovner*, Saverio Stranges*, Konstantinos Stroumpoulis*, Bruno F Sunguya*, Patrick J Sur*, Soumya Swaminathan*, Bryan L Sykes*, Cassandra E I Szoete*, Rafael Tabarés-Seisdedos*, Nikhil Tandon*, David Tanne*, Mohammad Tavakkoli*, Bineyam Taye*, Hugh R Taylor*, Braden J Te Ao*, Teketo Kassaw Tegegne*, Dejen Yemane Tekle*, Abdullah Suliman Terkawi*, Gizachew Assefa Tessema*, J S Thakur*, Alan J Thomson*, Andrew L Thorne-Lyman*, Amanda G Thrift*, George D Thurston*, Ruoyan Tobe-Gai*, Marcello Tonelli*, Roman Topor-Madry*, Fotis Topouzis*, Bach Xuan Tran*, Thomas Truelsen*, Zacharie Tsala Dimbuene*, Miltiadis Tsilimbaris*, Abera Kenay Tura*, Emin Murat Tuzcu*, Stefanos Tyrovolas*, Kingsley N Ukwaja*, Eduardo A Undurraga*, Chigozie Jesse Umeke*, Olalekan A Uthman*, Coen H van Gool*, Jim van Os*, Tommi Vasankari*, Ana Maria Nogales Vasconcelos*, Narayanaswamy Venketasubramanian*, Francesco S Violante*, Vasiliy Victorovich Vlassov*, Stein Emil Vollset*, Gregory R Wagner*, Mitchell T Wallin*, Linhong Wang*, Scott Weinchenthal*, Elisabete Weiderpass*, Robert G Weintraub*, Andrea Werdecker*, Ronny Westerman*, Tissa Wijeratne*, James D Wilkinson*,

Hywel C Williams*, Charles Shey Wiysonge*, Solomon Meseret Woldeyohannes*, Charles D A Wolfe*, Sungho Won*, Gelin Xu*, Ajit Kumar Yadav*, Bereket Yakob*, Lijing L Yan*, Yuichiro Yano*, Mehdi Yaseri*, Pengpeng Ye*, Paul Yip*, Naohiro Yonemoto*, Seok-Jun Yoon*, Mustafa Z Younis*, Chuhanhua Yu*, Zoubida Zaidi*, Maysaa El Sayed Zaki*, Hajo Zeeb*, Sanjay Zodpey*, David Zoniess*, Liesl Joanna Zuhlik*, Theo Vos†, Alan D Lopez†, Christopher J L Murray†.

*Authors listed alphabetically. †Joint senior authors.

Affiliations

Institute for Health Metrics and Evaluation (N J Kassebaum MD, M Arora BSA, R M Barber BS, J Brown MAIS, A Carter BS, D C Casev BA, F J Charlson PhD, M M Coates MPH, M Coggleshall BA, L Cornaby BS, Prof L Dandona MD, D J Dicker BS, H E Erskine PhD, A J Ferrari PhD, C Fitzmaurice MD, K Foreman PhD, M H Forouzanfar MD, N Fullman MPH, E M Goldberg BS, N Graetz MPH, J A Haagsma PhD, Prof S I Hay DSc, C O Johnson PhD, L Kemmer PhD, I A Khalil MD, Y Kinfu PhD, M J Kutz BS, H H Kyu PhD, J Leung PhD, Prof S S Lim PhD, Prof R Lozano PhD, J Mikesell BS, Prof A H Mokdad PhD, M D Mooney BS, Prof M Naghavi MD PhD, G Nguyen MPH, E Nsoesie PhD, D M Pigott DPhil, C Pinho BA, Z Rankin BS, N Reinig BS, L Sandar BS, A Smith BA, R J D Sorensen MPH, J Stanaway PhD, C Steiner MPH, S Teeple BA, B A Thomas MD, C Troeger MPH, A VanderZanden MSc, J A Wagner BS, V Wanga MS, Prof H A Whiteford PhD, Prof M Zhou PhD, L Zoeckler BA, T Achoki MD, A Afshin MD, L T Alexander BA, C Allen BA, G M Anderson MSE, B Bell MLIS, K Bienhoff MA, S Biryukov BS, J D Blore PhD, K Estep MPA, J Friedman BA, J Frostad MPH, W W Godwin BS, P Y Liu BA, F Masiye PhD, A Millear BA, M Mirarefin MPH, M Moradi-Lakeh MD, J E Mumford BA, M Ng PhD, M B Reitsma BS, A Reynolds BA, G A Roth MD, P J Sur BA, Prof S E Vollset DrPH, Prof T Vos PhD, Prof A D Lopez PhD, Prof C J L Murray DPhil), Harborview/UW Medicine (R G Ellenbogen MD), Harborview Injury Prevention and Research Center (C N Mock PhD), University of Washington, Seattle, WA, USA (Prof B O Anderson MD, N D Futran MD); Centre of Excellence in Women and Child Health (Prof Z A Bhutta PhD), Aga Khan University, Karachi, Pakistan (M I Nisar MSc); Centre for Global Child Health (Prof Z A Bhutta PhD, N Akseer MSc), The Hospital for Sick Children, Toronto, ON, Canada (G A deVeber MD); School of Public Health (F J Charlson PhD, H E Erskine PhD, J Leung PhD, Prof H A Whiteford PhD, A A Abajobir MPH, A J Ferrari PhD, L D Knibbs PhD), School of Dentistry (Prof R Laloo PhD), University of Queensland, Brisbane, QLD, Australia (N K M Alam MPH, H N Gouda PhD, Y Guo PhD, Prof J J McGrath MD); Queensland Centre for Mental Health Research, Brisbane, QLD, Australia (F J Charlson PhD, H E Erskine PhD, J Leung PhD, Prof H A Whiteford PhD, A J Ferrari PhD); Centre for Control of Chronic Conditions (P Jeemon PhD), Public Health Foundation of India, New Delhi, India (Prof L Dandona MD, R Dandona PhD, G A Kumar PhD); Department of Zoology (P W Gething PhD), Oxford Big Data Institute, Li Ka Shing Centre for Health Information and Discovery, University of Oxford, Oxford, UK (Prof S I Hay DSc), and Nuffield Department of Medicine (D Bisanzio PhD), University of Oxford, Oxford, UK (R Ali FRCP, D A Bennett PhD, Prof V Jha DM, K Rahimi DM); Department of Public Health, Erasmus MC, University Medical Center Rotterdam, Netherlands (J A Haagsma PhD); Centre for Research & Action in Public Health, Faculty of Health, University of Canberra, Canberra, ACT, Australia (Y Kinfu PhD); National Center for Chronic and Noncommunicable Disease Control and Prevention (L Duan MD, Y Jin MS, Y Li MPH, Prof L Wang MD, P Ye MPH, Prof M Zhou PhD), Chinese Center for Disease Control and Prevention, Beijing, China (Prof X Liang MD); Center for Translation Research and Implementation Science, National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, MD, USA (G A Mensah MD); Department of Global Health and Population (Prof J A Salomon PhD), Department of Nutrition (A L Thorne-Lyman ScD), Harvard T H Chan School of Public Health (Prof T Bärnighausen MD, I R Campos-Nonato PhD, E L Ding ScD, M S Farvid PhD, G R Wagner MD), Harvard University, Boston, MA, USA (J R A Fitchett MD); Jimma University, Jimma,

Ethiopia (K H Abate MS, M B Ahmed MPH, T T Gebrehiwot MPH, A T Gebremedhin MPH); La Sapienza, University of Rome, Rome, Italy (C Abbafati PhD); Virginia Tech, Blacksburg, VA, USA (Prof K M Abbas PhD); Department of Neurology, Cairo University, Cairo, Egypt (Prof F Abd-Allah MD); NMSSM Government College Kalpetta, Kerala, India (Prof B Abraham MPhil); Institute for Global Health (Prof I Abubakar PhD), Farr Institute of Health Informatics Research (A Banerjee DPhil), University College London, London, UK; Infectious Disease Epidemiology Group, Weill Cornell Medical College in Qatar, Doha, Qatar (L J Abu-Raddad PhD); Institute of Community and Public Health, Birzeit University, Ramallah, Palestine (N M Abu-Rmeileh PhD); Department of Epidemiology and Preventive Medicine, School of Public Health and Preventive Medicine (I N Ackerman PhD, Prof R Buchbinder PhD, Prof B Gabbe PhD), Department of Medicine, School of Clinical Sciences at Monash Health, (Prof A G Thrift PhD), Eastern Health Clinical School (B K Lloyd PhD), Monash University, Melbourne, VIC, Australia; College of Medicine, University of Ibadan, Ibadan, Nigeria (A O Adebiyi MD); University College Hospital, Ibadan, Nigeria (A O Adebiyi MD); Olabisi Onabanjo University, Ago-Iwoye, Nigeria (I A Adedeji MS); University of Extremadura, Cáceres, Spain (Prof J C Adsuar PhD); Direction du District Sanitaire de Haho, Notse, Togo (K A Afanvi MD); Faculté des Sciences de Santé, Université de Lomé, Lomé, Togo (K A Afanvi MD); Institution of Public Health Sciences, Stockholm, Sweden (E E Agardh PhD); Dalla Lana School of Public Health (N Akseer MSc), Department of Nutritional Sciences, Faculty of Medicine (A Badawi PhD), Centre for Addiction and Mental Health (S Popova PhD), University of Toronto, Toronto, ON, Canada (A Agarwal BHSc); McMaster University, Hamilton, ON, Canada (A Agarwal BHSc); All India Institute of Medical Sciences, New Delhi, India (Prof S K Agarwal MD, A Roy DM, R Sagar MD, M Satpathy PhD, Prof N Tandon PhD); Department of Clinical Sciences Lund, Orthopedics, Clinical Epidemiology Unit (A Ahmad Kiadaliri PhD), Skane University Hospital, Department of Clinical Sciences Lund (Prof B Norrvig PhD), Lund University, Lund, Sweden; Health Services Management Research Center, Institute for Futures Studies in Health, Kerman University of Medical Sciences, Kerman, Iran (A Ahmad Kiadaliri PhD); Ophthalmic Research Center (H Ahmadih MD, M Yaseri PhD), Ophthalmic Epidemiology Research Center (M Katibeh MD), Shahid Beheshti University of Medical Sciences, Tehran, Iran; Department of Ophthalmology, Labbafinejad Medical Center, Tehran, Iran (H Ahmadih MD); Washington University in Saint Louis, St Louis, MO, USA (Z Al-Aly MD); Murdoch Childrens Research Institute (K Alam PhD, P Azzopardi MEpi, R Borschmann PhD, S M Colquhoun PhD, Prof G C Patton MD, R G Weintraub MBBS), Department of Paediatrics (P Azzopardi MEpi), Department of Medicine (A Meretoja PhD), Institute of Health and Ageing (Prof C E I Szoete PhD), Melbourne School of Population and Global Health (Prof A D Lopez PhD), University of Melbourne, Melbourne, VIC, Australia (K Alam PhD, R Borschmann PhD, Prof H R Taylor AC, Prof T Wijeratne MD); Sydney School of Public Health (Prof T R Driscoll PhD), University of Sydney, Sydney, NSW, Australia (K Alam PhD, J Leigh PhD, Prof A H Kemp PhD); Queensland Health, Herston, QLD, Australia (N K M Alam MPH); King Saud University, Riyadh, Saudi Arabia (S F Aldhahri MD, K A Altirkawi MD); Department of Anesthesiology (A S Terkawi MD), King Fahad Medical City, Riyadh, Saudi Arabia (S F Aldhahri MD); Department of Preventive and Social Medicine, Faculty of Medicine (M A Alegretti MD), School of Medicine (A V Aleman MD), Faculty of Medicine (F Cavalleri BS), University of the Republic, Montevideo, Uruguay (V Colistro MSc); Debre Markos University, Debre Markos, Ethiopia (Z A Alemu MPH, T K Tegegne MPH); Luxembourg Institute of Health (LIH), Strassen, Luxembourg (A Alkerwi PhD); School of Public Health, University of Lorraine, Nancy, France (Prof F Alla PhD); Department of Public Health Sciences (P Allebeck PhD, R H S Rabiee MPH), Department of Clinical Science, Intervention and Technology (Prof J J Carrero PhD), Department of Neurobiology, Care Sciences and Society (NVS) (S M Fereshtehnejad PhD), Aging Research Center (Prof M Kivipelto PhD), Department of Medical Epidemiology and Biostatistics (E Weiderpass PhD), Karolinska Institutet, Stockholm, Sweden (R Havmoeller PhD, Shireen Sindi PhD); Charité

Universitätsmedizin, Berlin, Germany (U Alsharif MPH); Spanish Observatory on Drugs, Government Delegation for the National Plan on Drugs, Ministry of Health, Social Policy and Equality, Madrid, Spain (E Alvarez Martin PhD); Universidad de Cartagena, Cartagena de Indias, Colombia (Prof N Alvis-Guzman PhD); School of Medicine (A T Amare MPH, Y A Melaku MPH), University of Adelaide, Adelaide, SA, Australia (L G Ciobanu MS, G A Tessema MPH); College of Medicine and Health Sciences, Bahir Dar University, Bahir Dar, Ethiopia (A T Amare MPH); Dignitas International, Zomba, Malawi (A Amberbir PhD); University of Cape Coast, Cape Coast, Ghana (A K Amegah PhD); National Hospital, Abuja, Nigeria (Prof E A Ameh MBBS); Environmental Health Research Center, Kurdistan University of Medical Sciences, Sanandaj, Iran (H Amini MSPH); Department of Epidemiology and Public Health (H Amini MSPH, T Fürst PhD), Swiss Tropical and Public Health Institute, Basel, Switzerland (C K Karema MSc); University of Basel, Basel, Switzerland (H Amini MSPH, T Fürst PhD); Ministry of Public Health, Beirut, Lebanon (W Ammar PhD, H L Harb MPH); Oregon Health and Science University, Portland, OR, USA (S M Amrock PhD, D Zonies MD); Department of Health Policy and Administration, College of Public Health, University of the Philippines Manila, Manila, Philippines (C A T Antonio MD); Self-employed, Kabul, Afghanistan (P Anvari MD); Department of Medical Sciences, Uppsala University, Uppsala, Sweden (Prof J Ärnöö PhD, Prof A Larsson PhD); Dalarna University, Falun, Sweden (Prof J Ärnöö PhD); School of Medicine, Institute of Microbiology and Immunology (Prof V S Arsic Arsenijevic PhD), Faculty of Medicine (A Barac PhD), University of Belgrade, Belgrade, Serbia; University Children Hospital, Belgrade, Serbia (Prof V S Arsic Arsenijevic PhD); University of Manitoba, Winnipeg, MB, Canada (A Artaman PhD); Department of Medical Emergency, School of Paramedic, Qom University of Medical Sciences, Qom, Iran (H Asayesh PhD); South Asian Public Health Forum, Islamabad, Pakistan (R J Asghar MD); Institut de Recherche Clinique du Bénin, Cotonou, Benin (E F G A Avokpaho MPH); Laboratoire d'Etudes et de Recherche-Action en Santé (LERAS Afrique), Parakou, Benin (E F G A Avokpaho MPH, F G Gankpé MD); Sanjay Gandhi Postgraduate Institute of Medical Sciences, Lucknow, India (A Awasthi MSc); The Judith Lumley Centre for Mother, Infant and Family Health Research, La Trobe University, Melbourne, VIC, Australia (B P Ayala Quintanilla PhD); Peruvian National Institute of Health, Lima, Peru (B P Ayala Quintanilla PhD); Wardliparingga Aboriginal Research Unit, South Australian Health and Medical Research Institute, Adelaide, SA, Australia (P Azzopardi MEpi); School of Health Sciences, University of Management and Technology, Lahore, Pakistan (U Bacha PhD); Public Health Agency of Canada, Toronto, ON, Canada (A Badawi PhD); Department of Environmental Health Engineering, Sri Ramachandra University, Chennai, India (K Balakrishnan PhD); School of Psychology, University of Auckland, Auckland, New Zealand (S L Barker-Collo PhD); Africa Health Research Institute, Mtubatuba, South Africa (Prof T Bärnighausen MD); Institute of Public Health, Heidelberg University, Heidelberg, Germany (Prof T Bärnighausen MD, S Mohammed PhD); Department of Occupational and Environmental Health (Prof L Barregard MD), Health Metrics Unit (Prof M Petzold PhD), University of Gothenburg, Gothenburg, Sweden; Department of Industrial Engineering, School of Engineering, Pontificia Universidad Javeriana, Bogotá, Colombia (L H Barrero ScD); Stanford University, Stanford, CA, USA (S Basu PhD, L C Del gobbo PhD); School of Public Health (Y A Melaku MPH), Mekelle University, Mekelle, Ethiopia (T A Bayou BS, B D Betsu MPH, G B Hailu MSc, D Y Tekle MS); Oxford University, Ho Chi Minh City, Vietnam (J Beardsley MChB); College of Public Health and Tropical Medicine, Jazan, Saudi Arabia (N Bedi MD); IRCCS—Istituto di Ricerche Farmacologiche Mario Negri, Milan, Italy (E Beghi MD); School of Medicine (K N Sheth MD), Yale University, New Haven, CT, USA (Prof M L Bell PhD, J J Huang MD); National Institute of Psychiatry Ramon de la Fuente, Mexico City, Mexico (C Benjet PhD, R A Gutiérrez PhD); Internal Medicine Department (Prof I S Santos PhD), University of São Paulo, São Paulo, Brazil (I M Bensenor PhD, Prof P A Lotufo DrPH); Debre Berhane University, Debre Berhan, Ethiopia (A Berhane PhD); Division of Health and Social Care Research (Prof C D Wolfe MD), King's College London, London, UK (E Bermabé PhD, Prof R J Hay DM); College of Health and Medical Sciences (H S Roba MPH), Haramaya University, Harar, Ethiopia (A S Beyene MPH, T A Hassen MS, Y M Mesfin MPH); Queen Elizabeth Hospital Birmingham, Birmingham, UK (N Bhala DPhil); University of Otago Medical School, Wellington, New Zealand (N Bhala DPhil); Postgraduate Institute of Medical Education and Research, Chandigarh, India (A Bhansali DM); Department of Infectious Disease Epidemiology (T Fürst PhD), Department of Epidemiology and Biostatistics (F B Piel PhD), Division of Brain Sciences (Prof T J Steiner PhD), Imperial College London, London, UK (S Bhatt DPhil, Prof A Majeed MD, M Soljak PhD); Independent Public Health Consultants, Addis Ababa, Ethiopia (S Biadgilign MPH); Department of Nephrology Issues of Transplanted Kidney, Academician V I Shumakov Federal Research Center of Transplantology and Artificial Organs, Moscow, Russia (B Bikbov MD); University of Iowa Hospitals and Clinics, Iowa City, IA, USA (A A Bin Abdulhak MD); Department of Community Medicine, University of Oslo, Oslo, Norway (Prof E Bjertness PhD); Vision & Eye Research Unit, Anglia Ruskin University, Cambridge, UK (Prof R R A Bourne FRCophth); Danube-University Krems, Krems, Austria (Prof M Brainin PhD); Faculty of Health Sciences and Social Work, Department of Public Health, Trnava University, Trnava, Slovakia (A Brazinova PhD, M Majdani PhD); International Neurotrauma Research Organization, Vienna, Austria (A Brazinova PhD); College of Medicine (J Shen PhD), Ohio State University, Columbus, OH, USA (Prof N J K Breitborde PhD); German Cancer Research Center, Heidelberg, Germany (Prof H Brenner MD); University of Leicester, Leicester, UK (Prof T S Brugha MD); Monash Department of Clinical Epidemiology, Cabrini Institute, Melbourne, VIC, Australia (Prof R Buchbinder PhD); University of California, San Francisco, San Francisco, CA, USA (G C Buckle MD); Al Shifa Trust Eye Hospital, Rawalpindi, Pakistan (Z A Butt PhD); National Centre for Epidemiology and Population Health, The Australian National University, Canberra, ACT, Australia (B Calabria PhD, Prof R M Lucas PhD); Transport and Road Safety (TARS) Research (S Boufous PhD), National Drug and Alcohol Research Centre (Prof L Degenhardt PhD), Brien Holden Vision Institute (Prof S Resnikoff MD), University of New South Wales, Kensington, NSW, Australia (B Calabria PhD, Prof P B Mitchell MD); National Institute of Public Health, Cuernavaca, Mexico (I R Campos-Nonato PhD, J C Campuzano PhD, H Gomez-Dantes MSc, I B Heredia-Pi PhD, A Jauregui MSc, J C Montañez Hernandez MSc, Prof E Servan-Mori MSc); Department of Biostatistics and Epidemiology, University of Oklahoma Health Sciences Center, Oklahoma City, OK, USA (H Carabin PhD); Telethon Kids Institute, the University of Western Australia, Princess Margaret Hospital for Children, Subiaco, WA, Australia (Prof J R Carapetis PhD); Metropolitan Autonomous University, Mexico City, Mexico (R Cárdenas ScD); Colombian National Health Observatory, Instituto Nacional de Salud, Bogotá, Colombia (C A Castañeda-Orjuela MSc); Epidemiology and Public Health Evaluation Group, Public Health Department, Universidad Nacional de Colombia, Bogotá, Colombia (C A Castañeda-Orjuela MSc); Caja Costarricense de Seguro Social, San Jose, Costa Rica (Prof J Castillo Rivas MPH); Universidad de Costa Rica, San Pedro, Montes de Oca, Costa Rica (Prof J Castillo Rivas MPH); Department of Medicine, University of Valencia/INCLIVA Health Research Institute and CIBERSAM, Valencia, Spain (F Catalá-López PhD); Clinical Epidemiology Program, Ottawa Hospital Research Institute, Ottawa, ON, Canada (F Catalá-López PhD); School of Nursing, College of Medicine, National Taiwan University, Taipei, Taiwan (Prof J Chang PhD); Clinical Governance Unit, Gold Coast Health, Southport, QLD, Australia (P P Chiang PhD); Crowd Watch Africa, Lusaka, Zambia (M Chibalabala BS); National Center for Child Health and Development, Setagaya, Japan (C E Chibueze PhD, R Mori PhD); University of Zambia, Lusaka, Zambia (V H Chisumpa MPhil, F Masiye PhD); University of Witwatersrand, Johannesburg, South Africa (V H Chisumpa MPhil); Seoul National University Medical Library, Seoul, South Korea (J J Choi PhD); World Bank, New Delhi, India (L Choudhury MPH); Bispebjerg University Hospital, Copenhagen, Denmark (Prof H Christensen DMSci); Ministerio de Salud Pública, Montevideo, Uruguay (V Colistro MSc); UNICEM, Montevideo, Uruguay (M Colomar MSc); Centre for

International Health, Dunedin School of Medicine (Prof J A Crump MD), Injury Prevention Research Unit, Department of Preventive and Social Medicine, Dunedin School of Medicine (Prof S Derrett PhD), University of Otago, Dunedin, New Zealand (Prof R G Poulton PhD); IRCCS—Istituto di Ricerche Farmacologiche Mario Negri, Bergamo, Italy (G Giussani Biol D, M Cortinovis Biotech D, N Perico MD, Prof G Remuzzi MD); Faculty of Medicine, Eduardo Mondlane University, Maputo, Mozambique (Prof A Damasceno PhD); Guy's and St Thomas' NHS Foundation Trust, London, UK (Prof P I Dargan FRCP); i3S—Instituto de Investigação e Inovação em Saúde and INEB—Instituto de Engenharia Biomédica (J das Neves PhD), EPIUnit—Institute of Public Health (J M Pedro MS), University of Porto, Porto, Portugal (J V Santos BHlthSc); Wellcome Trust Brighton & Sussex Centre for Global Health Research, Brighton, UK (Prof G Davey MD); Public Health England, London, UK (Prof A C Davis PhD, Prof J N Newton FRCP, Prof N Steel PhD); Griffith University, Brisbane, QLD, Australia (Prof D De Leo DSc); Mount Sinai Beth Israel, New York, NY, USA (Prof D C Des Jarlais PhD); Icahn School of Medicine at Mount Sinai, New York, NY, USA (Prof D C Des Jarlais PhD); Department of Community Medicine, Faculty of Medicine, University of Peradeniya, Peradeniya, Sri Lanka (S D Dharmaratne MD); Centre for Control of Chronic Conditions (P Jeemon PhD), Public Health Foundation of India, Gurgaon, India (P K Dhillon PhD, Prof S Zodpey PhD); RMIT University, Bundoora, VIC, Australia (Prof K E Doyle PhD); International Institute for Population Sciences, Mumbai, India (M Dubey MPhil, M H U Rahman MPhil, Prof U Ram PhD, A Singh PhD, A K Yadav MPhil); Federal University of Rio Grande do Sul, Porto Alegre, Brazil (B B Duncan PhD, C Kieling MD, Prof M I Schmidt MD); University of North Carolina, Chapel Hill, NC, USA (B B Duncan PhD); Non-Communicable Diseases Research Center, Endocrinology and Metabolism Population Sciences Institute (H Ebrahimi MD, Prof A Esteghamati MD, F Farzaadfar MD, N Hafezi-Nejad MD, A Kasaeian PhD, M Parsaeian PhD, F Pishgar MD, S Sheikbahaei MD), Liver and Pancreaticobiliary Diseases Research Center, Digestive Disease Research Institute, Shariati Hospital (H Ebrahimi MD), Digestive Diseases Research Institute (S Fahimi PhD, Prof R Malekzadeh MD, G Roshandel PhD, S G Sepanlou PhD), Center for Air Pollution Research, Institute for Environmental Research (M S Hassanzand PhD), Multiple Sclerosis Research Center, Neuroscience Institute (P Heydarpour MD), Hematology-Oncology and Stem Cell Transplantation Research Center (A Kasaeian PhD), Department of Epidemiology and Biostatistics, School of Public Health (M Parsaeian PhD), Uro-Oncology Research Center (F Pishgar MD), Sina Trauma and Surgery Research Center (Prof V Rahimi-Movaghar MD), Tehran University of Medical Sciences, Tehran, Iran (M Yaseri PhD); Eijkman-Oxford Clinical Research Unit, Jakarta, Indonesia (I Elyazar PhD); Arba Minch University, Arba Minch, Ethiopia (A Y Endries MPH); The Institute of Social and Economic Studies of Population, Russian Academy of Sciences, Moscow, Russia (Prof S P Ermakov DSc); Federal Research Institute for Health Organization and Informatics, Ministry of Health of the Russian Federation, Moscow, Russia (Prof S P Ermakov DSc); Ministry of Health and Medical Education, Tehran, Iran (B Eshrati PhD); Arak University of Medical Sciences, Arak, Iran (B Eshrati PhD); University of Louisville, Louisville, KY, USA (T A Farid MD, A R Khan MD); DGS Directorate General of Health, Lisbon, Portugal (C S E S Farinha MSc); Universidade Aberta, Lisbon, Portugal (C S E S Farinha MSc); Federal University of Sergipe, Aracaju, Brazil (Prof A Faro PhD); Harvard/MGH Center on Genomics, Vulnerable Populations, and Health Disparities, Mongan Institute for Health Policy, Massachusetts General Hospital, Boston, MA, USA (M S Farvid PhD); National Institute for Stroke and Applied Neurosciences (V L Feigin PhD), Auckland University of Technology, Auckland, New Zealand (B J Te Ao MPH); Institute of Education and Sciences, German Hospital Oswaldo Cruz, São Paulo, Brazil (Prof J G Fernandes PhD); Centre for Experimental Medicine & Rheumatology, William Harvey Research Institute, Barts and The London School of Medicine & Dentistry, Queen Mary University of London, London, UK (J C Fernandes PhD); Bielefeld University, Bielefeld, Germany (F Fischer MPH); Institute of Gerontology, Academy of Medical Science, Kiev, Ukraine (N Foigt PhD); University of

Edinburgh, Edinburgh, UK (Prof F G R Fowkes PhD); James Cook University, Townsville, QLD, Australia (R C Franklin PhD); CHU Hassan II, Fès, Morocco (F G Gankpé MD); Manhiça Health Research Center, Manhiça, Mozambique (A L Garcia-Basteiro MSc Epidemiology); Barcelona Institute for Global Health, Barcelona, Spain (A L Garcia-Basteiro MSc Epidemiology); Ludwig Maximilians University, Munich, Germany (A T Gelbremedhin MPH); Division of Human Nutrition (J M Geleijnse PhD), Wageningen University, Wageningen, Netherlands (T Jibat MS); The Peter Doherty Institute for Infection and Immunity, University of Melbourne & The Royal Melbourne Hospital, Melbourne, VIC, Australia (K B Gibney FRACP); The Royal Melbourne Hospital, Melbourne, VIC, Australia (K B Gibney FRACP); College of Medicine, Howard University, Washington, DC, USA (R F Gillum MD, A Mehari MD); College of Medicine, University of Hail, Hail, Saudi Arabia (I A Ginawi MD); School of Public Health (A D Hailu MPH), Addis Ababa University, Addis Ababa, Ethiopia (A Z Giref PhD, D Haile MPH, T Jibat MS); University Hospital of Dijon, Dijon, France (Prof M Giroud MD); College of Health and Medical Sciences (H S Roba MPH), Haramaya University, Dire Dawa, Ethiopia (M D Gishu MS A K Tura MPH); Kersa Health and Demographic Surveillance System, Harar, Ethiopia (M D Gishu MS); University of Massachusetts Boston, Boston, MA, USA (Prof P Gona PhD); Instituto de Investigaciones Científicas y Servicios de Alta Tecnología—INDICASAT-AIP, Ciudad del Saber, Panamá (A Goodridge PhD); Department of Health and Social Affairs, Government of the Federated States of Micronesia, Palikir, Federated States of Micronesia (S V Gopalani MPH); University of British Columbia, Vancouver, BC, Canada (C C Gotay PhD, Prof N Kisssoon MD, J A Kopec PhD, F Pourmalek PhD); Division of Epidemiology, Center for Public Health Sciences (A Goto PhD), National Cancer Center, Tokyo, Japan (M Inoue MD); Departments of Microbiology and Epidemiology & Biostatistics, Saint James School of Medicine, The Quarter, Anguilla (Prof H Gugnani PhD); West Virginia Bureau for Public Health, Charleston, WV, USA (R Gupta MD); Eternal Heart Care Centre and Research Institute, Jaipur, India (R Gupta PhD); Department of Anthropology, University of Delhi, Delhi, India (V Gupta PhD); Department of Global Public Health and Primary Care (A K Knudsen PhD, Prof S E Vollset DrPH), University of Bergen, Bergen, Norway (A D Hailu MPH, Prof O F Norheim PhD); Kilte Awlaelo Health and Demographic Surveillance System, Mekelle, Ethiopia (G B Hailu MSc); Brandeis University, Waltham, MA, USA (Y A Halasa MS, E A Undurraga PhD); Arabian Gulf University, Manama, Bahrain (Prof R R Hamadeh DPhil); Hamdan Bin Mohammed Smart University, Dubai, United Arab Emirates (S Hamidi PhD); Wayne County Department of Health and Human Services, Detroit, MI, USA (M Hammami MD); University of New Mexico, Albuquerque, NM, USA (A J Handal PhD); School of Medicine and Pharmacology, University of Western Australia, Perth, WA, Australia (Prof G J Hankey MD); Harry Perkins Institute of Medical Research, Nedlands, WA, Australia (Prof G J Hankey MD); Western Australian Neuroscience Research Institute, Nedlands, WA, Australia (Prof G J Hankey MD); Sree Chitra Tirunal Institute for Medical Sciences and Technology, Trivandrum, India (S Harikrishnan DM); Parc Sanitari Sant Joan de Déu—CIBERSAM, Sant Boi de Llobregat (Barcelona), Spain (J M Haro MD); Universitat de Barcelona, Barcelona, Spain (J M Haro MD); International Foundation for Dermatology, London, UK (Prof R J Hay DM); Department of Medical Mycology and Parasitology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran (Prof M T Hedayati PhD); Department of Psychiatry, University Medical Center Groningen (Prof H W Hoek MD), University of Groningen, Groningen, Netherlands (A K Tura MPH); Department of Epidemiology, Mailman School of Public Health (Prof H W Hoek MD), Columbia University, New York, NY, USA (Prof V Skirbekki PhD); Rutgers University, New Brunswick, NJ, USA (D J Hoffman PhD); Nevada Division of Public and Behavioral Health, Department of Health and Human Services, Carson City, NV, USA (M Horino MPH); Department of Pulmonology, Yokohama City University Graduate School of Medicine, Yokohama, Japan (N Horita MD); Albert Einstein College of Medicine, Bronx, NY, USA (Prof H D Hosgood PhD); Public Health Division, The Pacific Community, Noumea, New Caledonia (D G Hoy PhD); Department of Epidemiology, Salah Azaiz Institute,

Tunis, Tunisia (Prof M Hsairi MD); Cambridge Health Alliance, Cambridge, MA, USA (H Huang MD); Aarhus University, Aarhus, Denmark (K M Iburg PhD); Boston Medical Center (B T Idrisov MD), School of Medicine (G F Kwan MD), Boston University, Boston, MA, USA; National Institute for Health Development, Tallinn, Estonia (K Innos PhD); Graduate School of Medicine (M Inoue MD), School of Public Health (Prof N Kawakami MD), University of Tokyo, Tokyo, Japan (K Shibuya MD); Department of Global and Community Health, George Mason University, Fairfax, VA, USA (K H Jacobsen PhD); Postgraduate Institute of Medicine, Colombo, Sri Lanka (A U Jayatilleke PhD); Institute of Violence and Injury Prevention, Colombo, Sri Lanka (A U Jayatilleke PhD); Centre for Chronic Disease Control, New Delhi, India (P Jeemon PhD); George Institute for Global Health India, New Delhi, India (Prof V Jha DM); Tianjin Centers for Disease Control and Prevention, Tianjin, China (G Jiang MD); Department of Health Development, Institute of Industrial Ecological Sciences, University of Occupational and Environmental Health, Kitakyushu, Japan (Y Jiang PhD); Department of Ocular Epidemiology and Visual Health, Institute of Ophthalmology Conde de Valencia, Mexico City, Mexico (A Jimenez-Corona PhD); General Directorate of Epidemiology, Ministry of Health, Mexico City, Mexico (A Jimenez-Corona PhD); Department of Ophthalmology, Medical Faculty Mannheim, Ruprecht-Karls-University Heidelberg, Mannheim, Germany (Prof J B Jonas MD); University College Cork, Cork, Ireland (Z Kabir PhD); Sante Stat & Analytical Research Institute, Kampala, Uganda (D K Kajungu MS); Mildmay Uganda, Kampala, Uganda (D K Kajungu MS); Society for Education, Action and Research in Community Health, Gadchiroli, India (Y Kalkonde MD); CSIR—Indian Institute of Toxicology Research, Lucknow, India (R Kamal MSc, C N Kesavachandran PhD); Department of Pulmonary Medicine, Zhongshan Hospital (J She MD), Fudan University, Shanghai, China (H Kan MD); University at Buffalo, Buffalo, NY, USA (A Kandel MBBS); Epidemiological and Statistical Methods Research Group, Helmholtz Centre for Infection Research, Braunschweig, Germany (A Karch MD); Hannover-Braunschweig Site, German Center for Infection Research, Braunschweig, Germany (A Karch MD); Quality and Equity Health Care, Kigali, Rwanda (C K Karema MSc); Case Western University Hospitals, Cleveland, OH, USA (C Karimkhani MD); Oklahoma State University, Tulsa, OK, USA (A Kaul MD); University of California San Francisco, San Francisco, CA, USA (D S Kazi MD); Institute of Tropical and Infectious Diseases, Nairobi, Kenya (P N Keiyoro PhD); School of Continuing and Distance Education, Nairobi, Kenya (P N Keiyoro PhD); Farr Institute (Prof R A Lyons MD), Swansea University, Swansea, UK (Prof A H Kemp PhD); South African Medical Research Council, Cape Town, South Africa (A P Kengne PhD, R Matzopoulos PhD, Prof C S Wiysonge PhD); School of Public Health and Family Medicine (R Matzopoulos PhD), Department of Psychiatry (Prof D J Stein PhD), University of Cape Town, Cape Town, South Africa (A P Kengne PhD, Prof B M Mayosi DPhil); Assuta Hospitals, Assuta Hashalom, Tel Aviv, Israel (Prof A Keren MD); Jordan University of Science and Technology, Irbid, Jordan (Prof Y S Khader ScD); Health Services Academy, Islamabad, Pakistan (E A Khan MPH); College of Medicine (Prof Y H Khan MD), Graduate School of Public Health (Prof S Won PhD), Seoul National University, Seoul, South Korea; Executive Board of the Health Ministers' Council for Cooperation Council States, Riyadh, Saudi Arabia (Prof T A M Khoja FRCP); Ball State University, Muncie, IN, USA (J Khubchandani PhD); Hospital de Clínicas de Porto Alegre, Porto Alegre, Brazil (C Kieling MD); Korea Health Industry Development Institute, Cheongju-si, South Korea (C Kim PhD); Department of Health Sciences, Northeastern University, Boston, MA, USA (Prof D Kim DrPH); Southern University College, Skudai, Malaysia (Y J Kim PhD); Centre for Disease Burden (A K Knudsen PhD, Prof S E Vollset DrPH), Department of Health Promotion (J C Skogen PhD), Norwegian Institute of Public Health, Oslo, Norway (M Savic PhD, Prof V Skirbekk PhD); Department of Preventive Cardiology, National Cerebral and Cardiovascular Center, Suita, Japan (Y Kokubo PhD); Division of Cardiology, Brown University, Providence, RI, USA (D Kolte MD); Sher-i-Kashmir Institute of Medical Sciences, Srinagar, India (Prof P A Koul MD); Research and Development Unit, Parc Sanitari Sant Joan de Deu (CIBERSAM), Barcelona, Spain (A Koyanagi MD); Department of Demography and

Public Health Research Institute (Prof B Kuade Defo PhD), Department of Social and Preventive Medicine, School of Public Health (Prof B Kuade Defo PhD), University of Montreal, Montreal, QC, Canada; Graduation Studies in Epidemiology, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil (Prof R S Kuchenbecker PhD); Institute of Public Health, Hacettepe University, Ankara, Turkey (B Kucuk Bicer PhD); Erasmus MC, University Medical Center Rotterdam, Rotterdam, Netherlands (Prof E J Kuipers PhD); Work Organizations, Work Disability Prevention, The Finnish Institute of Occupational Health, Helsinki, Finland (T Lallukka PhD, R Shiri PhD); Department of Public Health, Faculty of Medicine (T Lallukka PhD), University of Helsinki, Helsinki, Finland (T J Meretoja PhD); Department of Zoology, Lahore College for Women University, Lahore, Pakistan (A A Latif PhD); Servicio de Neurologia, Clinica Alemana, Universidad del Desarrollo, Santiago, Chile (P M Lavados MD); Instituto Nacional de Epidemiología "Dr Juan H Jara," Mar del Plata, Argentina (A E B Lawrynowicz MPH); College of Optometry, Nova Southeastern University, Fort Lauderdale, FL, USA (J L Leasher OD); State University of New York, Albany, Rensselaer, NY, USA (R Leung PhD); San Francisco VA Medical Center, San Francisco, CA, USA (Y Li PhD); School of Medicine, Wayne State University, Detroit, MI, USA (Prof S E Lipshultz MD, Prof J D Wilkinson MD); Children's Hospital of Michigan, Detroit, MI, USA (Prof S E Lipshultz MD); Rollins School of Public Health (E P Simard PhD), Emory University, Atlanta, GA, USA (Prof Y Liu PhD, Prof M R Phillips MD); Turning Point, Eastern Health, Melbourne, VIC, Australia (B K Lloyd PhD); University of Bari, Bari, Italy (Prof G Logroscino PhD); University of Bristol, Bristol, UK (K J Looker PhD, D Rai PhD); Aintree University Hospital National Health Service Foundation Trust, Liverpool, UK (Prof R Lunevicius PhD); School of Medicine, University of Liverpool, Liverpool, UK (Prof R Lunevicius PhD); Mansoura Faculty of Medicine, Mansoura, Egypt (H Magdy Abd El Razek MBBCH); Social Security Organization Research Institute, Tehran, Iran (M Mahdavi PhD); Institute of Health Policy and Management, Erasmus University Rotterdam, Rotterdam, Netherlands (M Mahdavi PhD); Universidade Federal de Minas Gerais, Belo Horizonte, Brazil (Prof D C Malta PhD); Division of Population and Patient Health, King's College London Dental Institute, London, UK (Prof W Marcenes PhD); University Hospital Doctor Peset, University of Valencia, Valencia, Spain (J Martinez-Raga PhD); CEU Cardenal Herrera University, Moncada (Valencia), Spain (J Martinez-Raga PhD); Department of Health Sciences, University of York, York, UK (A J Mason-Jones PhD); London School of Hygiene & Tropical Medicine, London, UK (Prof M McKee DSc, Prof G V S Murthy MD); Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA (P A Meaney MD); Children's Hospital of Philadelphia, Philadelphia, PA, USA (P A Meaney MD); University of West Florida, Pensacola, FL, USA (P Memiah PhD); Saudi Ministry of Health, Riyadh, Saudi Arabia (Prof Z A Memish MD); College of Medicine, Alfaisal University, Riyadh, Saudi Arabia (Prof Z A Memish MD); United Nations Population Fund, Lima, Peru (W Mendoza MD); Department of Neurology, Helsinki University Hospital, Helsinki, Finland (A Meretoja PhD); Helsinki University Hospital, Comprehensive Cancer Center, Breast Surgery Unit, Helsinki, Finland (T J Meretoja PhD); Ifakara Health Institute, Bagamoyo, Tanzania (F A Mhimbira MS); Pacific Institute for Research & Evaluation, Calverton, MD, USA (T R Miller PhD); Centre for Population Health, Curtin University, Perth, WA, Australia (T R Miller PhD); University of Ottawa, Ottawa, ON, Canada (E J Mills PhD); Kyrgyz State Medical Academy, Bishkek, Kyrgyzstan (Prof E M Mirrakhimov PhD); National Center of Cardiology and Internal Disease, Bishkek, Kyrgyzstan (Prof E M Mirrakhimov PhD); University of Salahaddin, Erbil, Iraq (K A Mohammad PhD); Neuroscience Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran (A Mohammadi PhD); Health Systems and Policy Research Unit, Ahmadu Bello University, Zaria, Nigeria (S Mohammed PhD); Institute for Maternal and Child Health, IRCCS Burlo Garofolo, Trieste, Italy (L Monasta DSc, M Montico MSc, L Ronfani PhD); Department of Community Medicine, Gastrointestinal and Liver Disease Research Center, Preventive Medicine and Public Health Research Center, Iran University of Medical Sciences, Tehran, Iran (M Moradi-Lakeh MD); Competence Center Mortality-Follow-Up of

the German National Cohort (A Werdecker PhD), Federal Institute for Population Research, Wiesbaden, Germany (Prof U O Mueller PhD, R Westerman PhD); West Herts Hospitals NHS Trust, Watford, UK (M E Murdoch FRCP); Indian Institute of Public Health (Prof G V S Murthy MD), Public Health Foundation of India, Gurgaon, India (P K Dhillon PhD, Prof S Zodpey PhD); Graduate School of Public Health, University of Pittsburgh, Pittsburgh, PA, USA (Prof J B Nachege PhD); Stellenbosch University, Cape Town, South Africa (Prof J B Nachege PhD, Prof S Seedat PhD, Prof C S Wiysonge PhD); Bloomberg School of Public Health (Prof J B Nachege PhD), Johns Hopkins University, Baltimore, MD, USA (B X Tran PhD); International Centre for Diarrhoeal Disease Research, Bangladesh (icddr,b), Dhaka, Bangladesh (A Naheed PhD); Azienda Ospedaliera Papa Giovanni XXIII, Bergamo, Italy (Prof L Naldi MD, Prof G Remuzzi MD); Suraj Eye Institute, Nagpur, India (V Nangia MD); Ministry of Health and Social Welfare, Dar es Salaam, Tanzania (F N Ngalesoni MSc); Institute for Global Health Innovations, Duy Tan University, Da Nang, Vietnam (Q L Nguyen MD); Institute For Research, Socio-Economic Development and Communication, Yaoundé, Cameroon (P M Nkamedje Pete MS); Hospital Universitari de Bellvitge, L'Hospitalet, Spain (J M Nolla PhD); Institute of Health and Biomedical Innovation, Queensland University of Technology, Brisbane, QLD, Australia (R E Norman PhD); Center for Research on Population and Health, Faculty of Health Sciences, American University of Beirut, Beirut, Lebanon (Prof C M Obermeyer DSc); Centre for Health Research, Western Sydney University, Sydney, NSW, Australia (F A Ogbo MPH); Department of Preventive Medicine, School of Medicine, Kyung Hee University, Seoul, South Korea (Prof I Oh PhD); Human Sciences Research Council, Durban, South Africa (O Oladimeji MD); University of KwaZulu-Natal, Durban, South Africa (O Oladimeji MD, Prof B Sartorius PhD); Universidad Autónoma de Chile, Talca, Chile (Prof P R Olivares PhD); Center for Healthy Start Initiative, Lagos, Nigeria (B O Olusanya PhD, J O Olusanya MBA); University of Arizona, Tucson, AZ, USA (Prof E Oren PhD); IIS-Fundacion Jimenez Diaz-UAM, Madrid, Spain (Prof A Ortiz PhD); St Luke's International University, Tokyo, Japan (E Ota PhD); Hypertension in Africa Research Team (HART) (Prof A E Schutte PhD), North-West University, Mafikeng, South Africa (Prof A S Oyekale PhD); JSS Medical College, JSS University, Mysore, India (Prof M PA DNB); Department of Medical Humanities and Social Medicine, College of Medicine, Kosin University, Busan, South Korea (E Park PhD); Department of Community Health Sciences (Prof S B Patten PhD), University of Calgary, Calgary, AB, Canada (Prof M Tonelli MD); Health Research Centre of Angola, Caxito, Angola (J M Pedro MS); REQUIMTE/LAQV, Laboratório de Farmacognosia, Departamento de Química, Faculdade de Farmácia, Universidade do Porto, Porto, Portugal (Prof D M Pereira PhD); Flinders University, Adelaide, SA, Australia (Prof K Pesudovs PhD); University of the Witwatersrand, Johannesburg, South Africa (Prof M Petzold PhD); Shanghai Jiao Tong University School of Medicine, Shanghai, China (Prof M R Phillips MD); Durban University of Technology, Durban, South Africa (J D Pillay PhD); Exposure Assessment and Environmental Health Indicators, German Environment Agency, Berlin, Germany (D Plass DrPH); Department of Public Health, Erasmus University Medical Center, Rotterdam, Netherlands (S Polinder PhD); The Fred Hollows Foundation, Sydney, NSW, Australia (N M Prasad DO); Centre for Eye Research Australia, Melbourne, VIC, Australia (N M Prasad DO); Department of Community Medicine, School of Medicine, Alborz University of Medical Sciences, Karaj, Iran (M Qorbani PhD); A T Still University, Kirksville, MO, USA (A Radfar MD); Comtech School of Public Health, Lahore, Pakistan (A Rafay MS); Research and Evaluation Division, BRAC, Dhaka, Bangladesh (M Rahman PhD); Hamad Medical Corporation, Doha, Qatar (S U Rahman FCPS); Society for Health and Demographic Surveillance, Suri, India (R K Rai MPH); ERAWEB Program, UMIT, Hall in Tirol, Austria (S Rajsic MD); University of Missouri, Columbia, MO, USA (M Raji PhD); University of Michigan Health Systems, Ann Arbor, MI, USA (K Ranganathan MD); Walden University, Minneapolis, MN, USA (Prof A H Refaat PhD); Suez Canal University, Ismailia, Egypt (Prof A H Refaat PhD); Department of Biomedical and Clinical Sciences L Sacco, University of Milan, Milan, Italy (Prof G Remuzzi MD); Hospital das Clínicas da Universidade Federal de Minas Gerais, Belo

Horizonte, Brazil (Prof A L Ribeiro MD); UO Neurologia USL Umbria 1, Città di Castello, Italy (S Ricci FRCPEd); (ISGlobal) Instituto de Salud Global de Barcelona, Barcelona, Spain (D Rojas-Rueda PhD); Golestan Research Center of Gastroenterology and Hepatology, Golestan University of Medical Sciences, Gorgan, Iran (G Roshandel PhD); World Health Organization, Accra, Ghana (B B Sackey MS); Marshall University J Edwards School of Medicine, Huntington, WV, USA (J R Sanabria MD); Case Western Reserve University, Cleveland, OH, USA (J R Sanabria MD); IIS-Fundacion Jimenez Diaz, Madrid, Spain (M D Sanchez-Niño PhD); Universidad Ciencias Aplicadas y Ambientales, Bogotá, Colombia (R Sarmento-Suarez MPH); Marshall University, Huntington, WV, USA (M Sawhney PhD); Federal University of Santa Catarina, Florianópolis, Brazil (I J C Schneider PhD, D A S Silva PhD); South African Medical Research Council, Potchefstroom, South Africa (Prof A E Schutte PhD); University of Alabama at Birmingham, Birmingham, AL, USA (D C Schwebel PhD, J A Singh MD); Tufts Medical Center, Boston, MA, USA (S Shahraz PhD); Independent Consultant, Karachi, Pakistan (M A Shaikh MD); Indian Institute of Technology Ropar, Rupnagar, India (R Sharma MA); Research Institute at Nationwide Children's Hospital, Columbus, OH, USA (J Shen PhD); National Institute of Infectious Diseases, Tokyo, Japan (M Shigematsu PhD); Sandia National Laboratories, Albuquerque, NM, USA (M Shigematsu PhD); Department of Public Health Science, Graduate School (Prof M Shin PhD), Department of Preventive Medicine, College of Medicine (S Yoon PhD), Korea University, Seoul, South Korea; Reykjavik University, Reykjavik, Iceland (I D Sigfusdotir PhD); Feinberg School of Medicine (J I Silverberg MD), Department of Preventive Medicine (Y Yano MD), Northwestern University, Chicago, IL, USA; Institute for Human Development, New Delhi, India (P K Singh PhD); Alcohol and Drug Research Western Norway (J C Skogen PhD), Stavanger University Hospital, Stavanger, Norway (K Søreide PhD); Department of Community Medicine, International Medical University, Kuala Lumpur, Malaysia (C T Sreeramareddy MD); Attikon University Hospital, Athens, Greece (V Stathopoulou PhD); University of East Anglia, Norwich, UK (Prof N Steel PhD); South African Medical Research Council Unit on Anxiety & Stress Disorders, Cape Town, South Africa (Prof D J Stein PhD); University of California, San Diego, La Jolla, CA, USA (M B Stein MD); Department of Neuroscience, Norwegian University of Science and Technology, Trondheim, Norway (Prof T J Steiner PhD, Prof L J Stovner PhD); Norwegian Advisory Unit on Headache, St Olavs Hospital, Trondheim, Norway (Prof L J Stovner PhD); Luxembourg Institute of Health, Strassen, Luxembourg (S Stranges PhD); Alexandra General Hospital of Athens, Athens, Greece (K Stroumpoulis PhD); Centre Hospitalier Public du Cotentin, Cherbourg, France (K Stroumpoulis PhD); Muhimbili University of Health and Allied Sciences, Dar es Salaam, Tanzania (B F Sunguya PhD); Indian Council of Medical Research, New Delhi, India (S Swaminathan MD); Departments of Criminology, Law & Society, Sociology, and Public Health, University of California, Irvine, Irvine, CA, USA (Prof B L Sykes PhD); Department of Medicine, University of Valencia, INCLIVA Health Research Institute and CIBERSAM, Valencia, Spain (Prof R Tabarés-Seisdedos PhD); Chaim Sheba Medical Center, Tel Hashomer, Israel (Prof D Tanne MD); Tel Aviv University, Tel Aviv, Israel (Prof D Tanne MD); New York Medical College, Valhalla, NY, USA (M Tavakkoli MD); Department of Biology, Colgate University, Hamilton, NY, USA (B Taye PhD); Department of Anesthesiology, University of Virginia, Charlottesville, VA, USA (A S Terkawi MD); Outcomes Research Consortium (A S Terkawi MD), Cleveland Clinic, Cleveland, OH, USA (Prof E M Tuzcu MD); University Of Gondar, Gondar, Ethiopia (G A Tessema MPH); School of Public Health, Post Graduate Institute of Medical Education and Research, Chandigarh, India (Prof J Thakur MD); Adaptive Knowledge Management, Victoria, BC, Canada (A J Thomson PhD); WorldFish, Penang, Malaysia (A L Thorne-Lyman ScD); Nelson Institute of Environmental Medicine, School of Medicine, New York University, Tuxedo, NY, USA (Prof G D Thurston ScD); National Center for Child Health and Development, Tokyo, Japan (R Tobe-Gai PhD); Institute of Public Health, Faculty of Health Sciences, Jagiellonian University Medical College, Kraków, Poland (R Topor-Madry PhD); Faculty of Health Sciences, Wrocław Medical University, Wrocław, Poland (R Topor-

Madry PhD); Aristotle University of Thessaloniki, Thessaloniki, Greece (Prof F Topouzis PhD); Hanoi Medical University, Hanoi, Vietnam (B X Tran PhD); Department of Neurology, Rigshospitalet, University of Copenhagen, Copenhagen, Denmark (T Truelsen DMSc); Department of Population Sciences and Development, Faculty of Economics and Management, University of Kinshasa, Kinshasa, Democratic Republic of the Congo (Z Tsala Dimbuene PhD); African Population and Health Research Center, Nairobi, Kenya (Z Tsala Dimbuene PhD); Department of Medicine, University of Crete, Heraklion, Greece (Prof M Tsilimbaris PhD); Parc Sanitari Sant Joan de Déu, Fundació Sant Joan de Déu, Universitat de Barcelona, CIBERSAM, Barcelona, Spain (S Tyrovolas PhD); Department of Internal Medicine, Federal Teaching Hospital, Abakaliki, Nigeria (K N Ukwaja MD); Ebonyi State University, Abakaliki, Nigeria (C J Uneke PhD); Warwick Medical School, University of Warwick, Coventry, UK (O A Uthman PhD); National Institute for Public Health and the Environment, Bilthoven, Netherlands (C H van Gool PhD); Maastricht University Medical Centre, Maastricht, Netherlands (Prof J van Os PhD); UKK Institute for Health Promotion Research, Tampere, Finland (Prof T Vasankari PhD); University of Brasilia, Brasília, Brazil (Prof A M N Vasconcelos PhD); Raffles Neuroscience Centre, Raffles Hospital, Singapore, Singapore (N Venketasubramanian FRCP); University of Bologna, Bologna, Italy (Prof F S Violante MD); National Research University Higher School of Economics, Moscow, Russia (Prof V V Vlassov MD); National Institute for Occupational Safety and Health, Washington, DC, USA (G R Wagner MD); VA Medical Center, Washington, DC, USA (M T Wallin MD); Neurology Department, Georgetown University, Washington, DC, USA (M T Wallin MD); McGill University, Montreal, QC, Canada (S Weichenthal PhD); Department of Research, Cancer Registry of Norway, Institute of Population-Based Cancer Research, Oslo, Norway (E Weiderpass PhD); Department of Community Medicine, Faculty of Health Sciences, University of Tromsø, The Arctic University of Norway, Tromsø, Norway (E Weiderpass PhD); Genetic Epidemiology Group, Folkhälsan Research Center, Helsinki, Finland (E Weiderpass PhD); Royal Children's Hospital, Melbourne, VIC, Australia (R G Weintraub MBBS); German National Cohort Consortium, Heidelberg, Germany (R Westerman PhD); Western Health, Footscray, VIC, Australia (Prof T Wijeratne MD); Children's Hospital of Michigan, Detroit, MI, USA (Prof J D Wilkinson MD); Centre of Evidence-based Dermatology, University of Nottingham, Nottingham, UK (Prof H C Williams DSc); Department of Epidemiology and Biostatistics, Institute of Public Health, University of Gondar, Gondar, Ethiopia (S M Woldeyohannes MPH); National Institute for Health Research Comprehensive Biomedical Research Centre, Guy's & St Thomas' NHS Foundation Trust and King's College London, London, UK (Prof C D Wolfe MD); Department of Neurology, Jinling Hospital, Nanjing University School of Medicine, Nanjing, China (Prof G Xu PhD); Discipline of Public Health Medicine, School of Nursing and Public Health, University of KwaZulu Natal, Durban, South Africa (B Yakob PhD); Global Health Research Center, Duke Kunshan University, Kunshan, China (Prof L L Yan PhD); Social Work and Social Administration Department and The Hong Kong Jockey Club Centre for Suicide Research and Prevention, University of Hong Kong, Hong Kong, China (Prof P Yip PhD); Department of Biostatistics, School of Public Health, Kyoto University, Kyoto, Japan (N Yonemoto MPH); Jackson State University, Jackson, MS, USA (Prof M Z Younis DrPH); Department of Epidemiology and Biostatistics, School of Public Health (Prof C Yu PhD); Global Health Institute (Prof C Yu PhD), Wuhan University, Wuhan, China; University Hospital, Setif, Algeria (Prof Z Zaidi PhD); Faculty of Medicine, Mansoura University, Mansoura, Egypt (Prof M E Zaki PhD); Leibniz Institute for Prevention Research and Epidemiology, Bremen, Germany (Prof H Zeeb PhD); Red Cross War Memorial Children's Hospital, Cape Town, South Africa (L J Zuhlk PhD)

Contributors

CJLM, NK, and NF prepared the first draft. CJLM and ADL conceived of the study and provided overall guidance. All other authors provided data, developed models, reviewed results, initiated modelling infrastructure, and/or reviewed and contributed to the report.

Declaration of interests

Benjamin O Anderson is supported by the Susan G Komen Leadership Grant Research Project, award number SAC160001. Itamar S Santos reports grants from FAPESP (Brazilian Public Agency), outside the submitted work. Carl Abelardo T Antonio reports grants, personal fees, and non-financial support from Johnson & Johnson (Philippines), outside the submitted work. Rafael Tabarés-Seisdedos and Ferrán Catalá-López are supported in part by grant PROMETEOII/2015/021 from Generalitat Valenciana, and Rafael Tabarés-Seisdedos is supported by the national grant PI14/00894 from ISCIII-FEDER. Pablo M Lavados reports grants, personal fees, and non-financial support from Bayer, non-financial support from Boehringer Ingelheim, grants and personal fees from AstraZeneca, grants from CONICYT, and grants from The George Institute for Global Health, outside the submitted work. Jasvinder A Singh serves as the principal investigator for an investigator-initiated study funded by Horizon pharmaceuticals through a grant to DINORA, a 501c3 entity, and is on the steering committee of OMERACT, an international organization that develops measures for clinical trials and receives arms-length funding from 36 pharmaceutical companies. Ai Koyanagi's work is supported by the Miguel Servet contract financed by the CP13/00150 and PI15/00862 projects, integrated into the National R+D+I and funded by the ISCIII General Branch Evaluation and Promotion of Health Research and the European Regional Development Fund (ERDF-FEDER). Aletta F Schutte is funded by the Medical Research Council of South Africa, and the South African Research Chair Initiative by the National Research Foundation. Ana Maria Nogales Vasconcelos and her team in Brazil has received funding from Ministry of Health (process number: 25000192049/2014-14). Mahesh PA acknowledges the Indian Council of Medical Research for financial support (ICMR Grant number 5/8/4-4 (Env)/2003-NCD-I). Katharine J Looker has received funding from the World Health Organization for the HSV-2 seroprevalence review which informs this work; during the study, KJL also received separate funding from the World Health Organization, USAID/PATH, Sexual Health 24, and the National Institute for Health Research (NIHR) Health Protection Research Unit (HPRU) in Evaluation of Interventions at the University of Bristol. Donal Bisanzio is supported by Bill & Melinda Gates Foundation (#OPP1068048). Thomas Fürst received financial support from the Swiss National Science Foundation (SNSF; project no P300P3-154634). Jost B Jonas reports personal fees from Consultant for Mundipharma (Cambridge, UK); from Patent holder with Biocompatibles UK (Fransham, Surrey, UK); Treatment of eye diseases using encapsulated cells encoding and secreting neuroprotective factor and/or anti-angiogenic factor; Patent number: 20120263794, patent application with University of Heidelberg (Heidelberg, Germany; Agents for use in the therapeutic or prophylactic treatment of myopia or hyperopia; Europäische Patentanmeldung 15 000 771.4), outside the submitted work. Rodrigo Sarmiento-Suarez receives institutional support from Universidad de Ciencias Aplicadas y Ambientales, UDCA, Bogotá, Colombia. Stefanos Tyrovolas's work is supported by the Foundation for Education and European Culture (IPEP), the Sara Borrell postdoctoral programme (reference no CD15/00019 from the Instituto de Salud Carlos III (ISCIII—Spain) and the Fondos Europeo de Desarrollo Regional (FEDER). Beatriz Paulina Ayala Quintanilla acknowledges institutional support of PRONABEC (National Program of Scholarship and Educational Loan), provided by the Peruvian Government, while studying for her doctoral course at the Judith Lumley Centre of La Trobe University funded by PRONABEC. Manami Inoue is the beneficiary of a financial contribution from the AXA Research fund as chair holder of the AXA Department of Health and Human Security, Graduate School of Medicine, The University of Tokyo from Nov 1, 2012; the AXA Research Fund has no role in this work. Laith J Abu-Raddad acknowledges the support of Qatar National Research Fund (NPRP 9-040-3-008) who provided the main funding for generating the data provided to the GBD-IHME effort. Yogeshwar Kalkonde is a Wellcome Trust/ DBT India Alliance Intermediate Fellow in Public Health. Sarah Derrett reports grants and fees from the Euroqol Foundation, outside the submitted work. Dan J Stein reports personal fees from Lundbeck, Novartis, AMBRF, Biocodex, Sevier, SUN, and CIPLA, and grants from

NRGF and MRC, outside the submitted work. Tea Lallukka reports funding from The Academy of Finland, grant 287488, as an Academy Research Fellow. Charles D A Wolfe's research was funded/supported by the National Institute for Health Research (NIHR) Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust and King's College London. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, the UK Department of Health, Public Health England, or the World Health Organization. All other authors declare no competing interests.

Acknowledgments

We would like to thank the countless individuals who have contributed to the Global Burden of Disease Study 2015 in various capacities. The data reported here have been supplied by the US Renal Data System (USRDS). Data for this research was provided by MEASURE Evaluation, funded by the United States Agency for International Development (USAID). Collection of these data was made possible by the US Agency for International Development (USAID) under the terms of cooperative agreement GPO-A-00-08-000-D3-00. Views expressed do not necessarily reflect those of USAID, the US Government, or MEASURE Evaluation. Parts of this material are based on data and information provided by the Canadian institute for Health Information. However, the analyses, conclusions, opinions and statements expressed herein are those of the author and not those of the Canadian Institute for Health information. The Palestinian Central Bureau of Statistics granted the researchers access to relevant data in accordance with license no SLN2014-3-170, after subjecting data to processing aiming to preserve the confidentiality of individual data in accordance with the General Statistics Law, 2000. The researchers are solely responsible for the conclusions and inferences drawn upon available data. This paper uses data from SHARE Waves 1, 2, 3 (SHARELIFE), 4 and 5 (DOIs: 10.6103/SHARE.w1.500, 10.6103/SHARE.w2.500, 10.6103/SHARE.w3.500, 10.6103/SHARE.w4.500, 10.6103/SHARE.w5.500), see Börsch-Supan and colleagues, 2013, for methodological details. The SHARE data collection has been primarily funded by the European Commission through FP5 (QLK6-CT-2001-00360), FP6 (SHARE-I3: RII-CT-2006-062193, COMPARE: CIT5-CT-2005-028857, SHARELIFE: CIT4-CT-2006-028812) and FP7 (SHARE-PREP: number 211909, SHARE-LEAP: number 227822, SHARE M4: number 261982). Additional funding from the German Ministry of Education and Research, the US National Institute on Aging (U01_AG09740-13S2, P01_AG005842, P01_AG08291, P30_AG12815, R21_AG025169, Y1-AG-4553-01, IAG_BSR06-11, and OGHA_04-064) and from various national funding sources is gratefully acknowledged. This study has been realised using the data collected by the Swiss Household Panel (SHP), which is based at the Swiss Centre of Expertise in the Social Sciences FORS. The project is financed by the Swiss National Science Foundation. The following individuals would like to acknowledge various forms of institutional support: Simon I Hay is funded by a Senior Research Fellowship from the Wellcome Trust (#095066), and grants from the Bill & Melinda Gates Foundation (OPP119467, OPP1093011, OPP1106023 and OPP1132415). Amanda G Thrift is supported by a fellowship from the National Health and Medical Research Council (GNT1042600). Panniyammakal Jeemon is supported by the Wellcome Trust-DBT India Alliance, Clinical and Public Health, Intermediate Fellowship (2015–2020). Boris Bikbov, Norberto Percio, and Giuseppe Remuzzi acknowledge that work related to this paper has been done on the behalf of the GBD Genitourinary Disease Expert Group supported by the International Society of Nephrology (ISN). Amador Goodridge acknowledges funding from Sistema Nacional de Investigadores de Panamá-SNI. José das Neves was supported in his contribution to this work by a Fellowship from Fundação para a Ciência e a Tecnologia, Portugal (SFRH/BPD/92934/2013). Lijing L Yan is supported by the National Natural Sciences Foundation of China grants (71233001 and 71490732). Olanrewaju Oladimeji is an African Research Fellow at Human Sciences Research Council (HSRC) and Doctoral Candidate at the University of KwaZulu-Natal (UKZN), South Africa, and would like to acknowledge the institutional support by leveraging on the existing organisational research infrastructure at HSRC and UKZN. Nicholas Steel received funding from Public Health England as a Visiting Scholar in the Institute for Health Metrics and Evaluation in 2016. No individuals acknowledged received additional compensation for their efforts.

References

- UN. Sustainable development goals. Oct 29, 2015. <http://www.un.org/sustainabledevelopment/sustainable-development-goals/> (accessed Oct 29, 2015).
- Murray CJL. Choosing indicators for the health-related SDG targets. *Lancet* 2015; **386**: 1314–17.
- Murray CJL. Shifting to Sustainable Development Goals—Implications for global health. *N Engl J Med* 2015; **373**: 1390–93.
- Nilsson M, Griggs D, Visbeck M. Policy: Map the interactions between Sustainable Development Goals. *Nature News* 2016; **534**: 320.
- UN. United Nations Sustainable Development. <http://www.un.org/sustainabledevelopment/health/> (accessed June 26, 2016).
- Murray CJL, Salomon JA, Mathers CD, Lopez AD. Summary Measures in Population Health: Concepts, Ethics, Measurement, and Applications. Geneva: World Health Organization, 2002.
- Sullivan DF. A single index of mortality and morbidity. *HSMHA Health Rep* 1971; **86**: 347–54.
- Mathers CD, Sadana R, Salomon JA, Murray CJ, Lopez AD. Healthy life expectancy in 191 countries, 1999. *Lancet* 2001; **357**: 1685–91.
- Murray CJ, Vos T, Lozano R, et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012; **380**: 2197–223.
- Murray CJ, Barber RM, Foreman KJ, et al. Global, regional, and national disability-adjusted life years (DALYs) for 306 diseases and injuries and healthy life expectancy (HALE) for 188 countries, 1990–2013: quantifying the epidemiological transition. *Lancet* 2015; **386**: 2145–91.
- Murray CJ. Quantifying the burden of disease: the technical basis for disability-adjusted life years. *Bull World Health Organ* 1994; **72**: 429–45.
- GBD 2015 Mortality and Causes of Death Collaborators. Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 2016; **388**: 1459–544.
- WHO. WHO methods and data sources for global burden of disease estimates 2000–2011. Geneva: World Health Organization, 2013.
- WHO. WHO methods and data sources for life tables 1990–2015. Geneva: World Health Organization, 2016.
- EHLEIS. Bibliography on Health Expectancy in Europe, April 2015. http://www.eurohex.eu/pdf/Reports_2015/2015_TR4%204_Bibliography.pdf (accessed June 7, 2016).
- WHO. WHO methods and data sources for country-level causes of death 2000–2012. Geneva: World Health Organization, 2014.
- WHO. Estimates for 2000–2012. http://www.who.int/healthinfo/global_burden_disease/estimates/en/ (accessed June 11, 2016).
- European Commission. Healthy Life Years (HLY). http://ec.europa.eu/health/indicators/healthy_life_years/hly_en.htm#fragment2 (accessed June 7, 2016).
- OECD. Health at a Glance: Europe 2012. Paris: Organisation for Economic Co-operation and Development, 2012. <http://www.oecd-ilibrary.org/content/book/9789264183896-en> (accessed June 7, 2016).
- GBD 2015 Risk Factors Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries during 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 2016; **388**: 1545–604.
- Murray CJ, Ezzati M, Flaxman AD, et al. GBD 2010: design, definitions, and metrics. *Lancet* 2012; **380**: 2063–66.
- Stevens GA, Alkema L, Black RE, et al. Guidelines for Accurate and Transparent Health Estimates Reporting: the GATHER statement. *Lancet* 2016; published online June 28. [http://dx.doi.org/10.1016/S0140-6736\(16\)30388-9](http://dx.doi.org/10.1016/S0140-6736(16)30388-9).
- Stevens GA, Alkema L, Black RE, et al. Guidelines for Accurate and Transparent Health Estimates Reporting: the GATHER statement. *PLoS Med* 2016; **13**: e1002056.
- Stanaway JD, Shepard DS, Undurraga EA, et al. The global burden of dengue: an analysis from the Global Burden of Disease Study 2013. *Lancet Infect Dis* 2016; **16**: 712–23.
- Salomon JA, Vos T, Hogan DR, et al. Common values in assessing health outcomes from disease and injury: disability weights measurement study for the Global Burden of Disease Study 2010. *Lancet* 2012; **380**: 2129–43.

For funding for SHARE see
<http://www.share-project.org>

- 26 Salomon JA, Haagsma JA, Davis A, et al. Disability weights for the Global Burden of Disease 2013 study. *Lancet Glob Health* 2015; 3: e712–23.
- 27 Salomon JA, Wang H, Freeman MK, et al. Healthy life expectancy for 187 countries, 1990–2010: a systematic analysis for the Global Burden Disease Study 2010. *Lancet* 2012; 380: 2144–62.
- 28 Calculating the Indices. Human Development Reports. <http://hdr.undp.org/en/content/calculating-indices> (accessed April 27, 2016).
- 29 WHO Ebola Response Team. West African Ebola epidemic after one year—slowing but not yet under control. *N Engl J Med* 2015; 372: 584–87.
- 30 Fries JF. The compression of morbidity. *Milbank Q* 2005; 83: 801–23.
- 31 Langa KM, Larson EB, Karlawish JH, et al. Trends in the prevalence and mortality of cognitive impairment in the United States: is there evidence of a compression of cognitive morbidity? *Alzheimers Dement* 2008; 4: 134–44.
- 32 Bardenheier BH, Lin J, Zhuo X, et al. Compression of disability between two birth cohorts of US adults with diabetes, 1992–2012: a prospective longitudinal analysis. *Lancet Diabetes Endocrinol* 2016; 4: 686–94.
- 33 Beltran-Sanchez H, Preston S, Canudas-Romo V. An integrated approach to cause-of-death analysis: cause-deleted life tables and decompositions of life expectancy. *Demogr Res* 2008; 19: 1323–50.
- 34 Jagger C, Matthews R, Matthews F, et al. The burden of diseases on disability-free life expectancy in later life. *J Gerontol A Biol Sci Med Sci* 2007; 62: 408–14.
- 35 van Gool CH, Picavet HSJ, Deeg DJ, et al. Trends in activity limitations: the Dutch older population between 1990 and 2007. *Int J Epidemiol* 2011; 40: 1056–67.
- 36 King G, Murray CJL, Salomon JA, Tandon A. Enhancing the validity and cross-cultural comparability of measurement in survey research. *Am Polit Sci Rev* 2004; 98: 191–207.
- 37 Salomon JA, Tandon A, Murray CJL. Comparability of self rated health: cross sectional multi-country survey using anchoring vignettes. *BMJ* 2004; 328: 258.
- 38 Tandon A, Murray CJL, Salomon JA, King G. Statistical Models for Enhancing Cross-Population Comparability. Geneva: World Health Organization, 2002.
- 39 Bowling A. Commentary: Trends in activity limitation. *Int J Epidemiol* 2011; 40: 1068–70.
- 40 Jamison DT, Summers LH, Alleyne G, et al. Global health 2035: a world converging within a generation. *Lancet* 2013; 382: 1898–955.
- 41 GBD 2015 SDG Collaborators. Measuring the health-related Sustainable Development Goals in 188 countries: a baseline analysis from the Global Burden of Disease Study 2015. *Lancet* 2016; 388: 1815–52.
- 42 Engineer MH, Roy N, Fink S. ‘Healthy’ Human Development Indices. *Soc Indic Res* 2009; 99: 61–80.
- 43 National Institutes of Health (NIH). NIH unveils FY2016–2020 Strategic Plan. Dec 15, 2015. <http://www.nih.gov/news-events/news-releases/nih-unveils-fy2016–2020-strategic-plan> (accessed March 26, 2016).
- 44 Gillum LA, Gouveia C, Dorsey ER, et al. NIH disease funding levels and burden of disease. *PLoS One* 2011; 6: e16837.
- 45 Emdin CA, Odutayo A, Hsiao AJ, et al. Association between randomised trial evidence and global burden of disease: cross sectional study (Epidemiological Study of Randomized Trials—ESORT). *BMJ* 2015; 350: h117.
- 46 Catalá-López Ferrán, García-Altés Anna, Álvarez-Martín E, Génova-Máleras R, Morant-Ginestar C. Does the development of new medicinal products in the European Union address global and regional health concerns? *Popul Health Metr* 2010; 8: 34.
- 47 WHO. Investing in health research and development. <http://www.who.int/tdr/publications/tdr-research-publications/investing-in-health/en/> (accessed June 11, 2016).
- 48 Chalmers I, Bracken MB, Djulbegovic B, et al. How to increase value and reduce waste when research priorities are set. *Lancet* 2014; 383: 156–65.
- 49 GBD 2015 Risk Factors Collaborators. Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 2016; 388: 1661–736.
- 50 Leigh JP, Sheetz RM. Prevalence of back pain among fulltime United States workers. *Br J Ind Med* 1989; 46: 651–57.
- 51 Leino-Arjas P, Hänninen K, Puska P. Socioeconomic variation in back and joint pain in Finland. *Eur J Epidemiol* 1998; 14: 79–87.
- 52 Waddell G, Burton AK. Occupational health guidelines for the management of low back pain at work: evidence review. *Occup Med (Lond)* 2001; 51: 124–35.
- 53 Haddon W. Advances in the epidemiology of injuries as a basis for public policy. *Public Health Rep* 1980; 95: 411–21.
- 54 Moore L, Evans D, Hameed SM, et al. Mortality in Canadian trauma systems: a multicenter cohort Study. *Ann Surg* 2016; published online Jan 7 DOI:10.1097/SLA.0000000000001614.
- 55 Uthkarsh PS, Gururaj G, Reddy SS, Rajanna MS. Assessment and availability of trauma care services in a district hospital of south India: a field observational study. *Bull Emerg Trauma* 2016; 4: 93–100.
- 56 Ghajar J. Traumatic brain injury. *Lancet* 2000; 356: 923–29.
- 57 Cheatham ML, Malbrain ML, Kirkpatrick A, et al. Results from the International Conference of Experts on Intra-abdominal Hypertension and Abdominal Compartment Syndrome. II. Recommendations. *Intensive Care Med* 2007; 33: 951–62.
- 58 Stuke LE, Pons PT, Guy JS, Chapleau WP, Butler FK, McSwain NE. Prehospital spine immobilization for penetrating trauma—review and recommendations from the Prehospital Trauma Life Support Executive Committee. *J Trauma* 2011; 71: 763–69.
- 59 Rickard JL, Ntakiriyura G, Chu KM. Associations with perioperative mortality rate at a major referral hospital in Rwanda. *World J Surg* 2015; 40: 784–90.
- 60 Odhiambo FO, Beynon CM, Ogwang S, et al. Trauma-related mortality among adults in rural western Kenya: characterising deaths using data from a Health and Demographic Surveillance System. *PLoS One* 2013; 8: e79840.
- 61 Naish S, Dale P, Mackenzie JS, McBride J, Mengersen K, Tong S. Climate change and dengue: a critical and systematic review of quantitative modelling approaches. *BMC Infect Dis* 2014; 14: 167.
- 62 GBD 2015 Child Mortality Collaborators. Global, regional, national, and selected subnational levels of stillbirths, neonatal, infant, and under-5 mortality, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 2016; 388: 1727–76.
- 63 GBD 2015 Maternal Mortality Collaborators. Maternal mortality 1990 to 2015: a systematic analysis of the Global Burden of Disease 2015 Study. *Lancet* 2016; 388: 1777–814.
- 64 Murray CJL, Ferguson BD, Lopez AD, Guillot M, Salomon JA, Ahmad O. Modified logit life table system: principles, empirical validation, and application. *Popul Stud* 2003; 57: 165–82.
- 65 Wang H, Dwyer-Lindgren L, Lofgren KT, et al. Age-specific and sex-specific mortality in 187 countries, 1970–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012; 380: 2071–94.
- 66 GBD 2013 Mortality and Causes of Death Collaborators. Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2015; 385: 117–71.
- 67 Stouthard ME, Essink-Bot M, Bonsel G, Barendregt J, Kramers P. Disability weights for diseases in the Netherlands. Rotterdam: Department of Public Health, Erasmus University, 1997.
- 68 UN, Department of Economic and Social Affairs, Population Division. World Population Prospects: The 2012 Revision, Key Findings and Advance Tables. Working Paper number ESA/P/WP.227. New York: United Nations, 2013.