

Dimethyl 6-bromo-2-methyl-1,2-dihydroquinoline-2,4-dicarboxylate

Zeynep GÜLTEKİN,^a Michael Bolte^b and Tuncer HöKELEK^{c*}

^aDepartment of Chemistry, Çankırı Karatekin University, TR-18100, Çankırı, Turkey,

^bInstitut für Anorganische Chemie, J. W. Goethe-Universität Frankfurt, Max-von-Laue-Strasse 7, D-60438 Frankfurt/Main, Germany, and ^cDepartment of Physics, Hacettepe University, 06800 Beytepe, Ankara, Turkey

Correspondence e-mail: merzifon@hacettepe.edu.tr

Received 6 February 2012; accepted 8 February 2012

Key indicators: single-crystal X-ray study; $T = 173\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.007\text{ \AA}$; R factor = 0.053; wR factor = 0.153; data-to-parameter ratio = 14.2.

In the title compound, $\text{C}_{14}\text{H}_{14}\text{BrNO}_4$, the dihydropyridine ring adopts a screw-boat conformation. In the crystal, pairs of $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into inversion $R_2^2(10)$ dimers.

Related literature

For the synthesis of 1,2-dihydroquinolines, see: Hu *et al.* (2011); Yadav *et al.* (2007, 2008); Waldmann *et al.* (2008); Zhang & Ji (2011). For the biological activity of dihydroquinolines, see: Craig & Pearson (1971); Muren & Weissman (1971); Hamann *et al.* (1998); He *et al.* (2003); LaMontagne *et al.* (1989). For related structures, see: Gültekin *et al.* (2010, 2011a,b, 2012). For hydrogen-bond motifs, see: Bernstein *et al.* (1995). For ring-puckering parameters, see: Cremer & Pople (1975).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{14}\text{BrNO}_4$

$M_r = 340.16$

Triclinic, $P\bar{1}$

$a = 7.8273(9)\text{ \AA}$

$b = 10.4827(11)\text{ \AA}$

$c = 10.5029(12)\text{ \AA}$

$\alpha = 115.837(9)^\circ$

$\beta = 105.655(9)^\circ$

$\gamma = 96.889(8)^\circ$

$V = 718.25(17)\text{ \AA}^3$

$Z = 2$

Mo $K\alpha$ radiation

$\mu = 2.87\text{ mm}^{-1}$

$T = 173\text{ K}$

$0.33 \times 0.32 \times 0.22\text{ mm}$

Data collection

Stoe IPDS II two-circle diffractometer

Absorption correction: multi-scan (*MULABS*; Spek, 2009; Blessing, 1995)

$T_{\min} = 0.451$, $T_{\max} = 0.571$

7423 measured reflections
2692 independent reflections
2267 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.045$

$R_{\text{min}} = 0.451$, $T_{\max} = 0.571$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.053$

$wR(F^2) = 0.153$

$S = 1.21$

2692 reflections

189 parameters

1 restraint

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.96\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.46\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1 \cdots O1 ⁱ	0.90 (8)	2.12 (8)	3.013 (6)	176 (7)

Symmetry code: (i) $-x + 1, -y, -z$.

Data collection: *X-AREA* (Stoe & Cie, 2001); cell refinement: *X-AREA*; data reduction: *X-RED32* (Stoe & Cie, 2001); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON* (Spek, 2009).

The title compound was synthesized at RWTH Aachen University. The authors thank Professor Magnus Rueping of RWTH Aachen University, Germany, for helpful discussions.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5465).

References

- Bernstein, J., Davies, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Blessing, R. H. (1995). *Acta Cryst. A* **51**, 33–38.
- Craig, J. C. & Pearson, P. E. (1971). *J. Med. Chem.* **14**, 1221–1222.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Gültekin, Z., Bolte, M. & Hökelek, T. (2012). *Acta Cryst. E* **68**, o606.
- Gültekin, Z., Frey, W. & Hökelek, T. (2011b). *Acta Cryst. E* **67**, o576.
- Gültekin, Z., Frey, W., Tercan, B. & Hökelek, T. (2010). *Acta Cryst. E* **66**, o2891–o2892.
- Gültekin, Z., Frey, W., Tercan, B. & Hökelek, T. (2011a). *Acta Cryst. E* **67**, o672–o673.
- Hamann, L. G., Higuchi, R. I., Zhi, L., Edwards, J. P., Wang, X. N., Marschke, K. B., Kong, J. W., Farmer, L. J. & Jones, T. K. (1998). *J. Med. Chem.* **41**, 623–639.
- He, L., Chang, H.-X., Chou, T.-C., Savaraj, N. & Cheng, C.-C. (2003). *Eur. J. Med. Chem.* **38**, 101–107.
- Hu, X.-Y., Zhang, J.-C., Wei, W. & Ji, J.-X. (2011). *Tetrahedron Lett.* **52**, 2903–2905.
- LaMontagne, M. P., Blumbergs, B. & Smith, D. C. (1989). *J. Med. Chem.* **32**, 1728–1732.
- Muren, J. F. & Weissman, A. (1971). *J. Med. Chem.* **14**, 49–53.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Stoe & Cie (2001). *X-AREA* and *X-RED32*. Stoe & Cie, Darmstadt, Germany.

- Waldmann, H., Karunakar, G. V. & Kumar, K. (2008). *Org. Lett.* **10**, 2159–2162.
Yadav, J. S., Reddy, B. V. S., Premalatha, K. & Murty, M. S. R. (2007). *J. Mol. Catal. A Chem.* **271**, 161–163.
Yadav, J. S., Reddy, B. V. S., Yadav, N. N., Gupta, M. K. & Sridhar, B. (2008). *J. Org. Chem.* **73**, 6857–6859.
Zhang, J.-C. & Ji, J.-X. (2011). *ACS Catalysis*, **1**, 1360–1363.

supplementary materials

Acta Cryst. (2012). E68, o710–o711 [doi:10.1107/S1600536812005600]

Dimethyl 6-bromo-2-methyl-1,2-dihydroquinoline-2,4-dicarboxylate

Zeynep Gültekin, Michael Bolte and Tuncer Hökelek

Comment

1,2-Dihydroquinoline derivatives have been considerably important for the preparation of biologically important compounds (Craig & Pearson, 1971; Muren & Weissman, 1971). Many methods have been reported in the literature for the preparation of 1,2-dihydroquinolines (Yadav *et al.*, 2007, 2008). The most convenient method is the condensation of aromatic amines with ketones using a catalytic amount of a Lewis acid or Brønsted acid (Hu *et al.*, 2011; Waldmann *et al.*, 2008; Zhang & Ji, 2011). Dihydroquinolines are also powerful intermediates for the preparation of quinolines and many quinolines display biological effects (Hamann *et al.*, 1998; LaMontagne *et al.*, 1989; He *et al.*, 2003).

The structures of some 1,2-dihydroquinoline derivatives, $C_{16}H_{19}NO_4$ (Gültekin *et al.*, 2010), $C_{14}H_{15}NO_4$ (Gültekin *et al.*, 2011a), $C_{17}H_{21}NO_7$ (Gültekin *et al.*, 2011b) and $C_{16}H_{17}NO_5$ (Gültekin *et al.*, 2012) have also been determined.

In the title compound, (I), (Fig. 1), the ring A (C2–C4/C9/C10/N1) is not planar, but adopting a screw-boat conformation with puckering parameters (Cremer & Pople, 1975) $Q_T = 0.339(5)\text{\AA}$, $\varphi = -162.6(1.5)^\circ$ and $\theta = 129.2(1.1)^\circ$.

In the crystal structure, intermolecular N—H \cdots O hydrogen bonds (Table 1) link the molecules into centrosymmetric $R_{2}^{2}(10)$ dimers (Bernstein *et al.*, 1995) (Fig. 2).

Experimental


The title compound was synthesized by the literature method (Waldmann *et al.*, 2008). p-bromo aniline (100 mg, 1 eq) was dissolved in acetonitrile (1.5 ml), and then $Bi(OTf)_3$ (5 mol%, 0.05 eq) and methyl pyruvate (2.2 eq) were added to the mixture. The mixture was heated by microwave irradiation for 7 h until the starting material was completely consumed as monitored by TLC. The resultant residue was directly purified by flash chromatography on silica (EtOAc:Cyclohexane 1:2). Recrystallization over pentane and ethyl acetate (70:30) gave a yellow crystalline solid (yield: 81%), R_f 0.5 (2:1 Cyclohexane/EtOAc) m.p. 379–381 K.

Refinement


Amino H atom was located in a difference map and refined isotropically, the C-bound H atoms were positioned geometrically with C—H = 0.93 and 0.96 Å, and constrained to ride on their parent atoms, with $U_{iso}(H) = k \times U_{eq}(C)$, where $k = 1.2$ for aromatic and $k = 1.5$ for methyl H atoms, respectively.

Computing details

Data collection: *X-AREA* (Stoe & Cie, 2001); cell refinement: *X-AREA* (Stoe & Cie, 2001); data reduction: *X-AREA* (Stoe & Cie, 2001); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON* (Spek, 2009).

**Figure 1**

The molecular structure of the title compound with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

A partial packing diagram. Hydrogen bonds are shown as dashed lines.

Dimethyl 6-bromo-2-methyl-1,2-dihydroquinoline-2,4-dicarboxylate

Crystal data

C₁₄H₁₄BrNO₄

M_r = 340.16

Triclinic, P¹

Hall symbol: -P 1

a = 7.8273 (9) Å

b = 10.4827 (11) Å

c = 10.5029 (12) Å

α = 115.837 (9)°

β = 105.655 (9)°

γ = 96.889 (8)°

V = 718.25 (17) Å³

Z = 2

F(000) = 344

D_x = 1.573 Mg m⁻³

Mo Kα radiation, λ = 0.71073 Å

Cell parameters from 6851 reflections

θ = 3.5–25.9°

μ = 2.87 mm⁻¹

$T = 173\text{ K}$

Block, yellow

Data collection

Stoe IPDS II two-circle
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 ω scans
Absorption correction: multi-scan
(*MULABS*; Spek, 2009; Blessing, 1995)
 $T_{\min} = 0.451$, $T_{\max} = 0.571$

 $0.33 \times 0.32 \times 0.22\text{ mm}$

7423 measured reflections
2692 independent reflections
2267 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.045$
 $\theta_{\max} = 25.6^\circ$, $\theta_{\min} = 3.5^\circ$
 $h = -9 \rightarrow 9$
 $k = -12 \rightarrow 12$
 $l = -12 \rightarrow 12$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.153$
 $S = 1.21$
2692 reflections
189 parameters
1 restraint
Primary atom site location: structure-invariant
direct methods
Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0648P)^2 + 1.4243P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.96\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.46\text{ e \AA}^{-3}$
Extinction correction: *SHELXL97* (Sheldrick,
2008), $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{1/4}$
Extinction coefficient: 0.038 (5)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.98733 (9)	0.36331 (7)	0.80285 (6)	0.0521 (3)
O1	0.3080 (4)	0.0758 (4)	0.0556 (4)	0.0340 (8)
O2	0.2146 (4)	0.2722 (4)	0.0660 (4)	0.0309 (7)
O3	0.7138 (6)	0.6908 (4)	0.6183 (4)	0.0453 (9)
O4	0.6368 (5)	0.7295 (4)	0.4208 (4)	0.0400 (8)
N1	0.6748 (5)	0.2278 (4)	0.1501 (4)	0.0280 (8)
H1	0.678 (13)	0.138 (5)	0.085 (9)	0.11 (3)*
C2	0.5298 (6)	0.2890 (5)	0.1017 (5)	0.0262 (9)
C3	0.5671 (6)	0.4450 (5)	0.2227 (5)	0.0256 (9)
H3	0.5245	0.5116	0.1950	0.031*
C4	0.6599 (6)	0.4919 (5)	0.3699 (5)	0.0249 (9)
C5	0.8118 (6)	0.4208 (5)	0.5664 (5)	0.0308 (10)
H5	0.8093	0.5073	0.6451	0.037*

C6	0.8886 (6)	0.3213 (6)	0.5976 (5)	0.0326 (10)
C7	0.8965 (7)	0.1930 (6)	0.4853 (6)	0.0368 (11)
H7	0.9516	0.1288	0.5096	0.044*
C8	0.8212 (6)	0.1596 (5)	0.3345 (5)	0.0305 (10)
H8	0.8243	0.0720	0.2577	0.037*
C9	0.7412 (5)	0.2573 (5)	0.2986 (5)	0.0239 (9)
C10	0.7378 (6)	0.3919 (5)	0.4170 (5)	0.0237 (9)
C11	0.5237 (7)	0.2779 (6)	-0.0504 (5)	0.0374 (11)
H11A	0.5022	0.1767	-0.1236	0.056*
H11B	0.6393	0.3348	-0.0369	0.056*
H11C	0.4260	0.3153	-0.0856	0.056*
C12	0.3391 (6)	0.1982 (5)	0.0735 (5)	0.0250 (9)
C13	0.0293 (6)	0.1986 (6)	0.0410 (6)	0.0375 (11)
H13A	-0.0441	0.2672	0.0560	0.056*
H13B	0.0354	0.1623	0.1113	0.056*
H13C	-0.0256	0.1177	-0.0608	0.056*
C14	0.6766 (6)	0.6451 (5)	0.4849 (5)	0.0296 (10)
C15	0.6353 (10)	0.8756 (7)	0.5235 (8)	0.0572 (16)
H15A	0.6072	0.9287	0.4692	0.086*
H15B	0.7543	0.9266	0.6029	0.086*
H15C	0.5434	0.8686	0.5668	0.086*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0640 (4)	0.0637 (5)	0.0362 (3)	0.0183 (3)	0.0110 (3)	0.0345 (3)
O1	0.0334 (17)	0.0248 (17)	0.0434 (19)	0.0079 (14)	0.0182 (14)	0.0139 (15)
O2	0.0225 (15)	0.0373 (18)	0.0417 (18)	0.0120 (13)	0.0114 (13)	0.0257 (15)
O3	0.060 (2)	0.035 (2)	0.0338 (19)	0.0175 (18)	0.0171 (17)	0.0102 (16)
O4	0.054 (2)	0.0290 (18)	0.050 (2)	0.0216 (16)	0.0250 (18)	0.0240 (16)
N1	0.0270 (19)	0.032 (2)	0.0267 (18)	0.0135 (16)	0.0107 (15)	0.0142 (17)
C2	0.024 (2)	0.033 (2)	0.026 (2)	0.0108 (18)	0.0100 (17)	0.0169 (19)
C3	0.022 (2)	0.029 (2)	0.033 (2)	0.0087 (17)	0.0116 (17)	0.0196 (19)
C4	0.024 (2)	0.027 (2)	0.031 (2)	0.0083 (17)	0.0130 (17)	0.0186 (19)
C5	0.033 (2)	0.034 (3)	0.029 (2)	0.010 (2)	0.0124 (18)	0.018 (2)
C6	0.033 (2)	0.041 (3)	0.031 (2)	0.009 (2)	0.0105 (19)	0.025 (2)
C7	0.032 (2)	0.046 (3)	0.048 (3)	0.016 (2)	0.015 (2)	0.035 (3)
C8	0.029 (2)	0.030 (2)	0.036 (2)	0.0144 (19)	0.0129 (19)	0.016 (2)
C9	0.0187 (19)	0.030 (2)	0.029 (2)	0.0077 (17)	0.0115 (16)	0.0171 (18)
C10	0.023 (2)	0.025 (2)	0.028 (2)	0.0072 (17)	0.0108 (16)	0.0157 (18)
C11	0.037 (3)	0.050 (3)	0.032 (2)	0.014 (2)	0.015 (2)	0.024 (2)
C12	0.025 (2)	0.030 (2)	0.0205 (19)	0.0074 (18)	0.0099 (16)	0.0120 (17)
C13	0.022 (2)	0.047 (3)	0.050 (3)	0.007 (2)	0.011 (2)	0.030 (3)
C14	0.026 (2)	0.030 (2)	0.038 (3)	0.0093 (18)	0.0155 (19)	0.019 (2)
C15	0.073 (4)	0.034 (3)	0.079 (4)	0.031 (3)	0.040 (4)	0.029 (3)

Geometric parameters (\AA , $^\circ$)

Br1—C6	1.905 (5)	C8—H8	0.9300
O2—C13	1.455 (5)	C9—C8	1.395 (6)

O4—C15	1.442 (7)	C10—C5	1.393 (6)
N1—C2	1.452 (6)	C10—C9	1.427 (6)
N1—C9	1.380 (6)	C11—H11A	0.9600
N1—H1	0.899 (10)	C11—H11B	0.9600
C2—C3	1.499 (6)	C11—H11C	0.9600
C2—C11	1.537 (6)	C12—O1	1.197 (6)
C2—C12	1.548 (6)	C12—O2	1.322 (5)
C3—C4	1.344 (6)	C13—H13A	0.9600
C3—H3	0.9300	C13—H13B	0.9600
C4—C10	1.469 (6)	C13—H13C	0.9600
C4—C14	1.493 (6)	C14—O3	1.200 (6)
C5—H5	0.9300	C14—O4	1.344 (6)
C6—C5	1.378 (7)	C15—H15A	0.9600
C7—C6	1.371 (8)	C15—H15B	0.9600
C7—H7	0.9300	C15—H15C	0.9600
C8—C7	1.393 (7)		
C12—O2—C13	116.0 (4)	N1—C9—C10	120.0 (4)
C14—O4—C15	114.8 (4)	C8—C9—C10	119.7 (4)
C2—N1—H1	119 (6)	C5—C10—C4	125.1 (4)
C9—N1—C2	120.0 (4)	C5—C10—C9	118.7 (4)
C9—N1—H1	112 (6)	C9—C10—C4	116.2 (4)
N1—C2—C3	109.0 (3)	C2—C11—H11A	109.5
N1—C2—C11	108.4 (4)	C2—C11—H11B	109.5
N1—C2—C12	110.7 (4)	C2—C11—H11C	109.5
C3—C2—C11	112.4 (4)	H11A—C11—H11B	109.5
C3—C2—C12	109.3 (4)	H11A—C11—H11C	109.5
C11—C2—C12	107.0 (3)	H11B—C11—H11C	109.5
C2—C3—H3	118.8	O1—C12—O2	124.9 (4)
C4—C3—C2	122.3 (4)	O1—C12—C2	124.3 (4)
C4—C3—H3	118.8	O2—C12—C2	110.7 (4)
C3—C4—C10	120.3 (4)	O2—C13—H13A	109.5
C3—C4—C14	119.0 (4)	O2—C13—H13B	109.5
C10—C4—C14	120.7 (4)	O2—C13—H13C	109.5
C6—C5—C10	120.0 (4)	H13A—C13—H13B	109.5
C6—C5—H5	120.0	H13A—C13—H13C	109.5
C10—C5—H5	120.0	H13B—C13—H13C	109.5
C5—C6—Br1	119.1 (4)	O3—C14—O4	121.9 (5)
C7—C6—Br1	119.0 (4)	O3—C14—C4	126.2 (4)
C7—C6—C5	122.0 (4)	O4—C14—C4	111.8 (4)
C6—C7—C8	119.5 (4)	O4—C15—H15A	109.5
C6—C7—H7	120.2	O4—C15—H15B	109.5
C8—C7—H7	120.2	O4—C15—H15C	109.5
C7—C8—C9	120.1 (4)	H15A—C15—H15B	109.5
C7—C8—H8	120.0	H15A—C15—H15C	109.5
C9—C8—H8	120.0	H15B—C15—H15C	109.5
N1—C9—C8	120.1 (4)		
C9—N1—C2—C3	40.9 (5)	C3—C4—C14—O4	-16.6 (6)

C9—N1—C2—C11	163.5 (4)	C10—C4—C14—O3	-17.1 (7)
C9—N1—C2—C12	-79.4 (5)	C10—C4—C14—O4	165.6 (4)
C2—N1—C9—C8	154.6 (4)	Br1—C6—C5—C10	-179.6 (3)
C2—N1—C9—C10	-29.0 (6)	C7—C6—C5—C10	0.5 (7)
N1—C2—C3—C4	-28.2 (6)	C8—C7—C6—Br1	178.6 (4)
C11—C2—C3—C4	-148.4 (4)	C8—C7—C6—C5	-1.5 (7)
C12—C2—C3—C4	92.9 (5)	C9—C8—C7—C6	1.0 (7)
N1—C2—C12—O1	-17.8 (6)	N1—C9—C8—C7	176.8 (4)
N1—C2—C12—O2	164.7 (3)	C10—C9—C8—C7	0.4 (7)
C3—C2—C12—O1	-137.9 (4)	C4—C10—C5—C6	-177.1 (4)
C3—C2—C12—O2	44.6 (4)	C9—C10—C5—C6	0.9 (6)
C11—C2—C12—O1	100.1 (5)	C4—C10—C9—N1	0.4 (6)
C11—C2—C12—O2	-77.4 (4)	C4—C10—C9—C8	176.9 (4)
C2—C3—C4—C10	3.3 (6)	C5—C10—C9—N1	-177.8 (4)
C2—C3—C4—C14	-174.5 (4)	C5—C10—C9—C8	-1.3 (6)
C3—C4—C10—C5	-169.9 (4)	O1—C12—O2—C13	3.0 (6)
C3—C4—C10—C9	12.0 (6)	C2—C12—O2—C13	-179.5 (4)
C14—C4—C10—C5	7.9 (7)	O3—C14—O4—C15	-2.4 (7)
C14—C4—C10—C9	-170.2 (4)	C4—C14—O4—C15	175.1 (4)
C3—C4—C14—O3	160.7 (5)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1···O1 ⁱ	0.90 (8)	2.12 (8)	3.013 (6)	176 (7)

Symmetry code: (i) $-x+1, -y, -z$.