

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

## *trans*-Tetraaquabis(isonicotinamide- $\kappa$ N<sup>1</sup>)nickel(II) bis(3-hydroxybenzoate) tetrahydrate

Ibrahim Göker Zaman,<sup>a</sup> Nagihan Çaylak Delibaş,<sup>b</sup> Hacali Necefoğlu<sup>a</sup> and Tuncer Hökelek<sup>c\*</sup>

<sup>a</sup>Department of Chemistry, Kafkas University, 36100 Kars, Turkey, <sup>b</sup>Department of Physics, Sakarya University, 54187 Esentepe, Sakarya, Turkey, and <sup>c</sup>Department of Physics, Hacettepe University, 06800 Beytepe, Ankara, Turkey  
Correspondence e-mail: merzifon@hacettepe.edu.tr


Received 16 January 2012; accepted 18 January 2012

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.051;  $wR$  factor = 0.157; data-to-parameter ratio = 14.5.

The asymmetric unit of the title compound,  $[\text{Ni}(\text{C}_6\text{H}_6\text{N}_2\text{O})_2(\text{H}_2\text{O})_4](\text{C}_7\text{H}_5\text{O}_3)_2 \cdot 4\text{H}_2\text{O}$ , contains one-half of the complex cation with the Ni<sup>II</sup> ion located on an inversion center, a 3-hydroxybenzoate counter-anion and two uncoordinated water molecules. Four water O atoms in the equatorial plane around the Ni<sup>II</sup> ion [Ni—O = 2.052 (2) and 2.079 (2) Å] form a slightly distorted square-planar arrangement, which is completed up to a distorted octahedron by the two N atoms [Ni—N = 2.075 (3) Å] from two isonicotinamide ligands. In the anion, the carboxylate group is twisted from the attached benzene ring by 8.8 (3)°. In the crystal, a three-dimensional hydrogen-bonding network, formed by classical O—H...O and N—H...O hydrogen bonds, consolidates the crystal packing, which also exhibits  $\pi$ – $\pi$  interactions between the benzene and pyridine rings, with centroid–centroid distances of 3.455 (2) and 3.621 (2) Å, respectively.

### Related literature

For general background, see: Bigoli *et al.* (1972); Krishnamachari (1974). For related structures, see: Hökelek *et al.* (2009*a,b,c,d,e*); Sertçelik *et al.* (2009*a,b*).



### Experimental

#### Crystal data

$[\text{Ni}(\text{C}_6\text{H}_6\text{N}_2\text{O})_2(\text{H}_2\text{O})_4](\text{C}_7\text{H}_5\text{O}_3)_2 \cdot 4\text{H}_2\text{O}$   
 $M_r = 721.29$   
Monoclinic,  $P2_1/n$   
 $a = 6.6884$  (3) Å  
 $b = 16.9271$  (5) Å  
 $c = 13.5543$  (4) Å

$\beta = 100.186$  (3)°  
 $V = 1510.37$  (9) Å<sup>3</sup>  
 $Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 0.73$  mm<sup>-1</sup>  
 $T = 100$  K  
 $0.46 \times 0.33 \times 0.18$  mm

#### Data collection

Bruker Kappa APEXII CCD area-detector diffractometer  
Absorption correction: multi-scan (SADABS; Bruker, 2005)  
 $T_{\min} = 0.750$ ,  $T_{\max} = 0.877$

13495 measured reflections  
3723 independent reflections  
3365 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.029$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.051$   
 $wR(F^2) = 0.157$   
 $S = 1.26$   
3723 reflections  
256 parameters  
12 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.86$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.69$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2—H21...O1 <sup>i</sup>	0.83 (5)	2.20 (5)	3.018 (4)	169 (4)
N2—H22...O8 <sup>ii</sup>	0.84 (5)	2.21 (5)	3.012 (4)	159 (5)
O3—H31...O7	0.79 (5)	1.91 (5)	2.696 (4)	175 (5)
O5—H51...O3 <sup>iii</sup>	0.85 (4)	1.87 (4)	2.716 (3)	177 (5)
O5—H52...O1 <sup>iii</sup>	0.84 (5)	1.99 (6)	2.795 (4)	160 (6)
O6—H61...O4 <sup>ii</sup>	0.85 (3)	1.86 (3)	2.693 (3)	169 (4)
O6—H62...O1 <sup>iv</sup>	0.85 (5)	1.87 (5)	2.685 (4)	159 (5)
O7—H71...O7 <sup>v</sup>	0.78 (4)	2.03 (3)	2.795 (4)	167 (6)
O7—H72...O2 <sup>vi</sup>	0.85 (5)	1.88 (5)	2.731 (4)	177 (5)
O8—H81...O7 <sup>vii</sup>	0.77 (4)	2.10 (4)	2.802 (4)	152 (6)
O8—H82...O2	0.83 (4)	1.93 (5)	2.752 (4)	171 (4)

Symmetry codes: (i)  $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$ ; (ii)  $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$ ; (iii)  $-x, -y, -z + 1$ ; (iv)  $-x + 1, -y, -z + 1$ ; (v)  $-x, -y + 1, -z + 1$ ; (vi)  $-x - \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$ ; (vii)  $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$ .

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve

structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON* (Spek, 2009).

The authors are indebted to Anadolu University and the Medicinal Plants and Medicine Research Centre of Anadolu University, Eskişehir, Turkey, for the use of X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV5237).

## References

- Bigoli, F., Braibanti, A., Pellinghelli, M. A. & Tiripicchio, A. (1972). *Acta Cryst.* **B28**, 962–966.
- Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Hökelek, T., Dal, H., Tercan, B., Aybirdi, Ö. & Necefoğlu, H. (2009e). *Acta Cryst.* **E65**, m627–m628.
- Hökelek, T., Dal, H., Tercan, B., Özbek, F. E. & Necefoğlu, H. (2009a). *Acta Cryst.* **E65**, m1330–m1331.
- Hökelek, T., Yılmaz, F., Tercan, B., Gürgen, F. & Necefoğlu, H. (2009b). *Acta Cryst.* **E65**, m1101–m1102.
- Hökelek, T., Yılmaz, F., Tercan, B., Özbek, F. E. & Necefoğlu, H. (2009c). *Acta Cryst.* **E65**, m768–m769.
- Hökelek, T., Yılmaz, F., Tercan, B., Sertçelik, M. & Necefoğlu, H. (2009d). *Acta Cryst.* **E65**, m1130–m1131.
- Krishnamachari, K. A. V. R. (1974). *Am. J. Clin. Nutr.* **27**, 108–111.
- Sertçelik, M., Tercan, B., Şahin, E., Necefoğlu, H. & Hökelek, T. (2009a). *Acta Cryst.* **E65**, m326–m327.
- Sertçelik, M., Tercan, B., Şahin, E., Necefoğlu, H. & Hökelek, T. (2009b). *Acta Cryst.* **E65**, m389–m390.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

## supporting information

*Acta Cryst.* (2012). E68, m200–m201 [doi:10.1107/S1600536812002218]

## ***trans*-Tetraaquabis(isonicotinamide- $\kappa N^1$ )nickel(II) bis(3-hydroxybenzoate) tetrahydrate**

**Ibrahim Göker Zaman, Nağihan Çaylak Delibaş, Hacali Necefoğlu and Tuncer Hökelek**

### **S1. Comment**

As a part of our ongoing investigation on transition metal complexes of nicotinamide (NA), one form of niacin (Krishnamachari, 1974), and/or the nicotinic acid derivative *N,N*-diethylnicotinamide (DENA), an important respiratory stimulant (Bigoli *et al.*, 1972), the title compound (I) was synthesized and its crystal structure is reported herein.

The asymmetric unit of (I) (Fig. 1) contains one Ni<sup>II</sup> ion on a centre of symmetry, one isonicotinamide (INA) ligand, one 3-hydroxybenzoate (HB) molecule, two coordinated and two uncoordinated water molecules, respectively. The structures of some DENA and/or NA complexes of Ni<sup>II</sup> and Co(II) ions, [Ni(C<sub>8</sub>H<sub>5</sub>O<sub>3</sub>)<sub>2</sub>(C<sub>10</sub>H<sub>14</sub>N<sub>2</sub>O)<sub>2</sub>(H<sub>2</sub>O)<sub>2</sub>] (Sertçelik *et al.*, 2009a), [Ni(C<sub>6</sub>H<sub>6</sub>N<sub>2</sub>O)<sub>2</sub>(H<sub>2</sub>O)<sub>4</sub>](C<sub>7</sub>H<sub>4</sub>FO<sub>2</sub>)<sub>2</sub> (Hökelek *et al.*, 2009a), [Ni(C<sub>6</sub>H<sub>6</sub>N<sub>2</sub>O)<sub>2</sub>(H<sub>2</sub>O)<sub>4</sub>](C<sub>8</sub>H<sub>5</sub>O<sub>3</sub>)<sub>2</sub>.2(H<sub>2</sub>O) (Hökelek *et al.*, 2009b), [Ni(C<sub>7</sub>H<sub>4</sub>BrO<sub>2</sub>)<sub>2</sub>(C<sub>6</sub>H<sub>6</sub>N<sub>2</sub>O)<sub>2</sub>(H<sub>2</sub>O)<sub>2</sub>] (Hökelek *et al.*, 2009c), [Co(C<sub>6</sub>H<sub>6</sub>N<sub>2</sub>O)<sub>2</sub>(H<sub>2</sub>O)<sub>4</sub>](C<sub>8</sub>H<sub>5</sub>O<sub>3</sub>)<sub>2</sub>.2(H<sub>2</sub>O) (Hökelek *et al.*, 2009d), [Co(C<sub>6</sub>H<sub>6</sub>N<sub>2</sub>O)(C<sub>9</sub>H<sub>10</sub>NO<sub>2</sub>)<sub>2</sub>(H<sub>2</sub>O)<sub>2</sub>] (Hökelek *et al.*, 2009e) and [Co(C<sub>8</sub>H<sub>5</sub>O<sub>3</sub>)<sub>2</sub>(C<sub>10</sub>H<sub>14</sub>N<sub>2</sub>O)<sub>2</sub>(H<sub>2</sub>O)<sub>2</sub>] (Sertçelik *et al.*, 2009b) have also been determined. In (I), INA ligands are monodentate. The four O atoms (O5, O6, and the symmetry-related atoms, O5', O6') in the equatorial plane around the Ni atom form a slightly distorted square-planar arrangement, while the slightly distorted octahedral coordination is completed by the two pyridine N atoms (N1, N1') of the INA ligands at 2.075 (3) Å from the Ni atom in the axial positions (Fig. 1). The average Ni—O bond length is 2.066 (2) Å. The intramolecular O—H...O hydrogen bonds (Table 1) link the uncoordinated water molecules to the HB anion. The dihedral angle between the planar carboxylate group (O1/O2/C1) and the benzene ring A (C2—C7) is 8.77 (27)°, while that between rings A and B (N1/C8—C12) is 1.53 (11)°.


In the crystal structure, intermolecular O—H...O and N—H...O hydrogen bonds (Table 1) link the molecules into a three-dimensional network, in which they may be effective in the stabilization of the structure.  $\pi$ – $\pi$  Contacts between the benzene and phenyl rings, Cg1...Cg2 and Cg1...Cg2<sup>i</sup>, [symmetry code: (i) -1 + x, y, z, where Cg1 and Cg2 are centroids of the rings A (C2—C7) and B (N1/C8—C12), respectively] may further stabilize the structure, with centroid-centroid distances of 3.621 (2) and 3.455 (2) Å, respectively.

### **S2. Experimental**

The title compound was prepared by the reaction of NiSO<sub>4</sub>·6H<sub>2</sub>O (1.314 g, 5 mmol) in H<sub>2</sub>O (100 ml) and INA (1.220 g, 10 mmol) in H<sub>2</sub>O (50 ml) with sodium 3-hydroxybenzoate (1.601 g, 10 mmol) in H<sub>2</sub>O (100 ml). The mixture was filtered and set aside to crystallize at ambient temperature for four weeks, giving blue single crystals.

### **S3. Refinement**

Atoms H51, H52, H61, H62, H71, H72, H81 and H82 (for H<sub>2</sub>O), H21 and H22 (for NH<sub>2</sub>) and H31 (for OH) were located in difference Fourier map and were refined by applying restraints. C-bound H-atoms were positioned geometrically (C—H = 0.93 Å) and constrained to ride on their parent atoms, with  $U_{\text{iso}}(\text{H}) = 1.2 \times U_{\text{eq}}(\text{C})$ .



**Figure 1**

The molecular structure of (I) with the atom-numbering scheme [symmetry code: (') -  $x$ , -  $y$ , -  $z$ ]. Displacement ellipsoids are drawn at the 50% probability level. Hydrogen bonds are shown as dashed lines.

***trans*-Tetraaquabis(isonicotinamide- $\kappa$ N')nickel(II) bis(3-hydroxybenzoate) tetrahydrate**

*Crystal data*

$[\text{Ni}(\text{C}_6\text{H}_6\text{N}_2\text{O})_2(\text{H}_2\text{O})_4](\text{C}_7\text{H}_5\text{O}_3)_2 \cdot 4\text{H}_2\text{O}$

$M_r = 721.29$

Monoclinic,  $P2_1/n$

Hall symbol: -P 2yn

$a = 6.6884$  (3) Å

$b = 16.9271$  (5) Å

$c = 13.5543$  (4) Å

$\beta = 100.186$  (3)°

$V = 1510.37$  (9) Å<sup>3</sup>

$Z = 2$

$F(000) = 756$

$D_x = 1.586$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 7580 reflections

$\theta = 2.4$ – $28.4$ °

$\mu = 0.73$  mm<sup>-1</sup>

$T = 100$  K

Rod-shaped, blue

$0.46 \times 0.33 \times 0.18$  mm

*Data collection*

Bruker Kappa APEXII CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2005)

$T_{\min} = 0.750$ ,  $T_{\max} = 0.877$

13495 measured reflections

3723 independent reflections

3365 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.029$

$\theta_{\max} = 28.6$ °,  $\theta_{\min} = 1.9$ °

$h = -8 \rightarrow 8$

$k = -22 \rightarrow 21$

$l = -17 \rightarrow 15$

Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.051$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.157$	$w = 1/[\sigma^2(F_o^2) + (0.0492P)^2 + 5.1781P]$
$S = 1.26$	where $P = (F_o^2 + 2F_c^2)/3$
3723 reflections	$(\Delta/\sigma)_{\max} < 0.001$
256 parameters	$\Delta\rho_{\max} = 0.86 \text{ e } \text{\AA}^{-3}$
12 restraints	$\Delta\rho_{\min} = -0.69 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.5000	0.0000	0.5000	0.00840 (16)
O1	-0.0786 (4)	0.04089 (13)	0.29703 (18)	0.0135 (5)
O2	-0.1778 (4)	0.11616 (14)	0.16276 (17)	0.0145 (5)
O3	-0.0150 (4)	0.39248 (14)	0.27901 (19)	0.0151 (5)
H31	-0.011 (8)	0.428 (3)	0.317 (4)	0.027 (13)*
O4	0.3941 (4)	0.32984 (14)	0.18014 (18)	0.0160 (5)
O5	0.3225 (4)	0.04752 (14)	0.59662 (18)	0.0128 (5)
H51	0.372 (9)	0.068 (3)	0.653 (3)	0.051 (17)*
H52	0.245 (8)	0.014 (3)	0.616 (5)	0.054 (18)*
O6	0.7696 (4)	0.03386 (14)	0.58654 (19)	0.0148 (5)
H61	0.803 (7)	0.0796 (16)	0.609 (3)	0.026 (12)*
H62	0.851 (7)	0.000 (3)	0.618 (4)	0.041 (16)*
O7	0.0025 (4)	0.51961 (15)	0.40009 (19)	0.0168 (5)
H71	-0.018 (9)	0.508 (4)	0.453 (2)	0.050*
H72	-0.101 (6)	0.548 (3)	0.380 (4)	0.047 (17)*
O8	0.0926 (4)	0.06780 (16)	0.04380 (19)	0.0192 (5)
H81	0.196 (5)	0.060 (4)	0.077 (4)	0.050*
H82	0.021 (7)	0.082 (3)	0.085 (3)	0.047 (17)*
N1	0.4858 (4)	0.10763 (15)	0.4259 (2)	0.0098 (5)
N2	0.5148 (5)	0.39569 (17)	0.3228 (2)	0.0140 (6)
H21	0.525 (7)	0.439 (3)	0.296 (3)	0.017 (11)*
H22	0.559 (8)	0.396 (3)	0.385 (4)	0.030 (13)*
C1	-0.1092 (5)	0.10742 (18)	0.2549 (2)	0.0112 (6)
C2	-0.0587 (5)	0.18065 (18)	0.3173 (2)	0.0106 (6)

C3	-0.0642 (5)	0.25416 (18)	0.2706 (2)	0.0110 (6)
H3	-0.1005	0.2580	0.2013	0.013*
C4	-0.0156 (5)	0.32139 (18)	0.3277 (3)	0.0120 (6)
C5	0.0358 (5)	0.31639 (19)	0.4314 (3)	0.0130 (6)
H5	0.0684	0.3619	0.4694	0.016*
C6	0.0386 (5)	0.2433 (2)	0.4783 (3)	0.0137 (6)
H6	0.0707	0.2399	0.5478	0.016*
C7	-0.0068 (5)	0.17514 (19)	0.4214 (2)	0.0121 (6)
H7	-0.0026	0.1261	0.4526	0.015*
C8	0.4370 (5)	0.11290 (18)	0.3253 (2)	0.0113 (6)
H8	0.4102	0.0666	0.2884	0.014*
C9	0.4250 (5)	0.18391 (18)	0.2748 (2)	0.0119 (6)
H9	0.3897	0.1851	0.2053	0.014*
C10	0.4660 (5)	0.25372 (18)	0.3287 (2)	0.0106 (6)
C11	0.5176 (5)	0.24858 (18)	0.4322 (2)	0.0124 (6)
H11	0.5467	0.2940	0.4706	0.015*
C12	0.5253 (5)	0.17508 (19)	0.4776 (2)	0.0118 (6)
H12	0.5594	0.1724	0.5471	0.014*
C13	0.4552 (5)	0.33050 (18)	0.2717 (2)	0.0115 (6)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Ni1	0.0102 (3)	0.0075 (3)	0.0072 (3)	-0.00025 (19)	0.00071 (19)	0.00018 (19)
O1	0.0146 (11)	0.0103 (10)	0.0151 (12)	0.0006 (8)	0.0015 (9)	0.0001 (8)
O2	0.0171 (12)	0.0151 (11)	0.0105 (11)	-0.0004 (9)	0.0007 (9)	-0.0006 (9)
O3	0.0218 (13)	0.0101 (11)	0.0132 (12)	0.0002 (9)	0.0022 (10)	0.0006 (9)
O4	0.0246 (13)	0.0122 (11)	0.0107 (11)	0.0019 (9)	0.0016 (10)	0.0017 (8)
O5	0.0145 (12)	0.0136 (11)	0.0113 (11)	-0.0003 (8)	0.0049 (9)	-0.0006 (9)
O6	0.0155 (12)	0.0098 (11)	0.0162 (12)	0.0000 (9)	-0.0052 (9)	-0.0013 (9)
O7	0.0199 (13)	0.0139 (11)	0.0155 (12)	0.0031 (9)	0.0004 (10)	0.0000 (9)
O8	0.0202 (13)	0.0214 (13)	0.0173 (13)	0.0031 (10)	0.0064 (10)	0.0001 (10)
N1	0.0084 (12)	0.0117 (12)	0.0096 (12)	0.0005 (9)	0.0025 (10)	0.0005 (9)
N2	0.0199 (15)	0.0109 (13)	0.0105 (14)	-0.0009 (10)	0.0007 (11)	0.0027 (10)
C1	0.0072 (13)	0.0127 (14)	0.0142 (15)	0.0003 (10)	0.0030 (11)	-0.0011 (11)
C2	0.0079 (14)	0.0115 (14)	0.0126 (15)	0.0008 (10)	0.0022 (11)	-0.0012 (11)
C3	0.0095 (14)	0.0147 (14)	0.0088 (14)	0.0014 (11)	0.0015 (11)	0.0008 (11)
C4	0.0094 (14)	0.0108 (14)	0.0160 (16)	0.0009 (11)	0.0027 (12)	0.0020 (12)
C5	0.0117 (14)	0.0129 (14)	0.0141 (16)	0.0003 (11)	0.0019 (12)	-0.0030 (11)
C6	0.0133 (15)	0.0172 (15)	0.0101 (15)	0.0010 (11)	0.0010 (12)	0.0001 (12)
C7	0.0129 (15)	0.0116 (14)	0.0124 (15)	0.0017 (11)	0.0034 (12)	0.0020 (11)
C8	0.0107 (14)	0.0113 (14)	0.0118 (15)	-0.0002 (11)	0.0019 (11)	-0.0009 (11)
C9	0.0120 (14)	0.0132 (14)	0.0105 (15)	0.0002 (11)	0.0016 (12)	0.0006 (11)
C10	0.0092 (14)	0.0102 (13)	0.0125 (15)	0.0011 (10)	0.0024 (11)	0.0018 (11)
C11	0.0143 (15)	0.0100 (14)	0.0129 (15)	0.0003 (11)	0.0026 (12)	-0.0011 (11)
C12	0.0107 (14)	0.0135 (14)	0.0115 (15)	0.0007 (11)	0.0028 (11)	-0.0003 (11)
C13	0.0100 (14)	0.0119 (14)	0.0133 (15)	0.0022 (11)	0.0038 (12)	0.0020 (11)

*Geometric parameters (Å, °)*

Ni1—O5	2.079 (2)	N2—H21	0.83 (5)
Ni1—O5 <sup>i</sup>	2.079 (2)	N2—H22	0.84 (5)
Ni1—O6	2.052 (2)	C2—C1	1.505 (4)
Ni1—O6 <sup>i</sup>	2.052 (2)	C2—C3	1.394 (4)
Ni1—N1	2.075 (3)	C2—C7	1.395 (4)
Ni1—N1 <sup>i</sup>	2.075 (3)	C3—H3	0.9300
O1—C1	1.263 (4)	C4—C3	1.383 (4)
O2—C1	1.260 (4)	C5—C4	1.389 (5)
O3—C4	1.373 (4)	C5—C6	1.390 (5)
O3—H31	0.79 (5)	C5—H5	0.9300
O4—C13	1.236 (4)	C6—H6	0.9300
O5—H51	0.85 (2)	C7—C6	1.391 (5)
O5—H52	0.85 (2)	C7—H7	0.9300
O6—H61	0.85 (2)	C8—H8	0.9300
O6—H62	0.85 (2)	C9—C8	1.378 (4)
O7—H71	0.784 (18)	C9—C10	1.391 (4)
O7—H72	0.85 (2)	C9—H9	0.9300
O8—H81	0.769 (18)	C10—C11	1.386 (5)
O8—H82	0.83 (2)	C10—C13	1.507 (4)
N1—C8	1.347 (4)	C11—H11	0.9300
N1—C12	1.341 (4)	C12—C11	1.385 (4)
N2—C13	1.326 (4)	C12—H12	0.9300
O5—Ni1—O5 <sup>i</sup>	180.00 (9)	C7—C2—C1	120.3 (3)
O6—Ni1—O5	94.23 (10)	C2—C3—H3	120.1
O6 <sup>i</sup> —Ni1—O5	85.77 (10)	C4—C3—C2	119.7 (3)
O6—Ni1—O5 <sup>i</sup>	85.77 (10)	C4—C3—H3	120.1
O6 <sup>i</sup> —Ni1—O5 <sup>i</sup>	94.23 (10)	O3—C4—C3	118.2 (3)
O6 <sup>i</sup> —Ni1—O6	180.0	O3—C4—C5	121.2 (3)
O6—Ni1—N1	89.53 (10)	C3—C4—C5	120.5 (3)
O6 <sup>i</sup> —Ni1—N1	90.47 (10)	C4—C5—C6	119.9 (3)
O6—Ni1—N1 <sup>i</sup>	90.47 (10)	C4—C5—H5	120.1
O6 <sup>i</sup> —Ni1—N1 <sup>i</sup>	89.53 (10)	C6—C5—H5	120.1
N1—Ni1—O5	89.02 (10)	C5—C6—C7	120.1 (3)
N1 <sup>i</sup> —Ni1—O5	90.98 (10)	C5—C6—H6	120.0
N1—Ni1—O5 <sup>i</sup>	90.98 (10)	C7—C6—H6	120.0
N1 <sup>i</sup> —Ni1—O5 <sup>i</sup>	89.02 (10)	C2—C7—H7	120.1
N1 <sup>i</sup> —Ni1—N1	180.0	C6—C7—C2	119.8 (3)
C4—O3—H31	111 (4)	C6—C7—H7	120.1
Ni1—O5—H51	123 (4)	N1—C8—C9	122.8 (3)
Ni1—O5—H52	113 (4)	N1—C8—H8	118.6
H52—O5—H51	99 (6)	C9—C8—H8	118.6
Ni1—O6—H61	128 (3)	C8—C9—C10	119.4 (3)
Ni1—O6—H62	121 (4)	C8—C9—H9	120.3
H61—O6—H62	110 (5)	C10—C9—H9	120.3
H72—O7—H71	100 (4)	C9—C10—C13	118.5 (3)

H81—O8—H82	103 (4)	C11—C10—C9	117.9 (3)
C8—N1—Ni1	122.0 (2)	C11—C10—C13	123.6 (3)
C12—N1—Ni1	120.4 (2)	C10—C11—H11	120.4
C12—N1—C8	117.6 (3)	C12—C11—C10	119.3 (3)
C13—N2—H21	123 (3)	C12—C11—H11	120.4
C13—N2—H22	123 (3)	N1—C12—C11	122.9 (3)
H21—N2—H22	113 (4)	N1—C12—H12	118.5
O1—C1—C2	118.5 (3)	C11—C12—H12	118.5
O2—C1—O1	123.7 (3)	O4—C13—N2	123.2 (3)
O2—C1—C2	117.8 (3)	O4—C13—C10	119.0 (3)
C3—C2—C1	119.6 (3)	N2—C13—C10	117.9 (3)
C3—C2—C7	120.1 (3)		
O5—Ni1—N1—C8	-131.1 (3)	C1—C2—C7—C6	-179.7 (3)
O5 <sup>i</sup> —Ni1—N1—C8	48.9 (3)	C3—C2—C7—C6	0.2 (5)
O5—Ni1—N1—C12	48.7 (2)	O3—C4—C3—C2	177.5 (3)
O5 <sup>i</sup> —Ni1—N1—C12	-131.3 (2)	C5—C4—C3—C2	-0.9 (5)
O6—Ni1—N1—C8	134.6 (3)	C6—C5—C4—O3	-178.4 (3)
O6 <sup>i</sup> —Ni1—N1—C8	-45.4 (3)	C6—C5—C4—C3	0.0 (5)
O6—Ni1—N1—C12	-45.5 (2)	C4—C5—C6—C7	1.1 (5)
O6 <sup>i</sup> —Ni1—N1—C12	134.5 (2)	C2—C7—C6—C5	-1.2 (5)
Ni1—N1—C8—C9	179.4 (2)	C10—C9—C8—N1	0.5 (5)
C12—N1—C8—C9	-0.5 (5)	C8—C9—C10—C11	-0.1 (5)
Ni1—N1—C12—C11	-179.8 (2)	C8—C9—C10—C13	179.0 (3)
C8—N1—C12—C11	0.1 (5)	C9—C10—C11—C12	-0.3 (5)
C3—C2—C1—O1	171.2 (3)	C13—C10—C11—C12	-179.4 (3)
C3—C2—C1—O2	-8.0 (4)	C9—C10—C13—O4	6.0 (5)
C7—C2—C1—O1	-8.9 (5)	C9—C10—C13—N2	-173.2 (3)
C7—C2—C1—O2	171.9 (3)	C11—C10—C13—O4	-175.0 (3)
C1—C2—C3—C4	-179.3 (3)	C11—C10—C13—N2	5.8 (5)
C7—C2—C3—C4	0.8 (5)	N1—C12—C11—C10	0.3 (5)

Symmetry code: (i)  $-x+1, -y, -z+1$ .

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
N2—H21 $\cdots$ O1 <sup>ii</sup>	0.83 (5)	2.20 (5)	3.018 (4)	169 (4)
N2—H22 $\cdots$ O8 <sup>iii</sup>	0.84 (5)	2.21 (5)	3.012 (4)	159 (5)
O3—H31 $\cdots$ O7	0.79 (5)	1.91 (5)	2.696 (4)	175 (5)
O5—H51 $\cdots$ O3 <sup>iii</sup>	0.85 (4)	1.87 (4)	2.716 (3)	177 (5)
O5—H52 $\cdots$ O1 <sup>iv</sup>	0.84 (5)	1.99 (6)	2.795 (4)	160 (6)
O6—H61 $\cdots$ O4 <sup>iii</sup>	0.85 (3)	1.86 (3)	2.693 (3)	169 (4)
O6—H62 $\cdots$ O1 <sup>i</sup>	0.85 (5)	1.87 (5)	2.685 (4)	159 (5)
O7—H71 $\cdots$ O7 <sup>v</sup>	0.78 (4)	2.03 (3)	2.795 (4)	167 (6)
O7—H72 $\cdots$ O2 <sup>vi</sup>	0.85 (5)	1.88 (5)	2.731 (4)	177 (5)



---

O8—H81···O7 <sup>vii</sup>	0.77 (4)	2.10 (4)	2.802 (4)	152 (6)
O8—H82···O2	0.83 (4)	1.93 (5)	2.752 (4)	171 (4)

---

Symmetry codes: (i)  $-x+1, -y, -z+1$ ; (ii)  $-x+1/2, y+1/2, -z+1/2$ ; (iii)  $x+1/2, -y+1/2, z+1/2$ ; (iv)  $-x, -y, -z+1$ ; (v)  $-x, -y+1, -z+1$ ; (vi)  $-x-1/2, y+1/2, -z+1/2$ ; (vii)  $-x+1/2, y-1/2, -z+1/2$ .