

Macrophyllsaponin E: A Novel Compound from the Roots of *Astragalus oleifolius*

Erdal BEDİR,^{a,b} Ihsan CALIS,^a and Ikhlas A. KHAN^{*,b,c}

Hacettepe University, Faculty of Pharmacy, Department of Pharmacognosy,^a TR-06100 Ankara, Turkey, National Center for Natural Products Research, Research Institute of Pharmaceutical Sciences,^b and Department of Pharmacognosy, School of Pharmacy,^c The University of Mississippi, University, MS 38677, U.S.A.

Received December 17, 1999; accepted March 17, 2000

Macrophyllsaponin E, a novel cycloartane-type triterpene, has been isolated from the roots of *Astragalus oleifolius*. The structure elucidation of the compound was achieved by a combination of one- and two-dimensional NMR techniques [¹H–¹H-correlation spectroscopy (COSY), ¹H–¹³C-heteronuclear multiple quantum correlation spectroscopy (HMQC), and ¹H–¹³C-heteronuclear multiple-bond correlation spectroscopy (HMBC)] and high resolution electrospray ionization Fourier transformation mass spectrometry (HR-ESI-FT-MS).

Key words *Astragalus oleifolius*; cycloartane-type triterpene; macrophyllsaponin E, Leguminosae

Astragalus L., the largest genus in the family Leguminosae, is represented by 380 species in the flora of Turkey.¹⁾ The roots of *Astragalus* species represent a very old and well-known drug in traditional medicine for its usage as an antiperspirant, diuretic and tonic drug. It has also been used in the treatment of diabetes mellitus, nephritis, leukemia and uterine cancer.²⁾ In the district of Anatolia, located in South Eastern Turkey, an aqueous extract of the roots of *Astragalus* is traditionally used against leukemia and for its wound-healing properties. Known biologically active constituents of *Astragalus* roots represent two major classes of compounds, polysaccharides and saponins.²⁾ *Astragalus* polysaccharides are already known to have anticancer and immune enhancing properties in both *in vitro* and *in vivo* experiments.^{3–5)} Chemical studies on *Astragalus* saponins have reported the presence of cycloartane-type triterpenoid glycosides which were found to exert biological activities (*e.g.* anti-inflammatory, analgesic, diuretic, hypotensive and sedative effects).⁶⁾

Our earlier investigations performed on *Astragalus* species resulted in the isolation of a series of cycloartane-type triterpenic saponins,^{7–13)} as well as the compounds which were evaluated in lymphocyte stimulation tests showed immunomodulatory activity. The activity is relatively potent and thereby deserves further attention.⁸⁾

We previously reported the isolation and the structure determination of macrophyllsaponins A–D from *Astragalus oleifolius*.⁷⁾ In our continuing search, we have isolated a novel cycloartane-type triterpene glycoside, named as macrophyllsaponin E (**1**). This paper deals with the isolation and the structural elucidation of **1**.

The high resolution electrospray ionization Fourier transformation mass spectrometry (HR-ESI-FT-MS) spectra of **1** exhibited ion peak for [M+Na]⁺ at *m/z* 839.4721, which is compatible with the molecular formulae C₄₂H₇₂O₁₅.

Taking into account the results of our comprehensive ¹H- and ¹³C-NMR studies and previous knowledge derived from metabolites isolated from the genus *Astragalus*,^{7–13)} the main features of a cycloartane-type triterpene were evident: characteristic signals due to cyclopropane-methylene protons as an AX system (δ 0.44, 0.79, J_{AX} =4.5 Hz, H₂-19), six tertiary methyl groups (δ 0.89, 1.05, 1.07, 1.09, 1.2, 1.27; respectively, H₃-29, H₃-18, H₃-30, H₃-28, H₃-27, H₃-26) and a sec-

ondary methyl group (0.95 d, J =6.3; H₃-21). Additionally, the resonances for two anomeric protons were observed at δ 4.54 (d, J =7.8 Hz) and 4.36 (d, J =7.8 Hz), indicative of the presence of two β -linked sugar units. Thus, **1** was considered to be a cycloartane-type triterpene diglycoside. This observation was supported by the ¹³C-NMR spectral data of **1** (Table 1). The ¹H- and ¹³C-NMR data (Table 1) supported the assignment of the sugar moieties in **1** as two β -D-glucopyranose.¹⁴⁾ The remaining carbon and proton resonances were consistent with C₃₀H₅₂O₅ for the aglycon moiety. This implied five saturated ring systems because there were no olefinic protons.

Among the sugar signals, the remaining signals showed correlations with the resonances at highfield, indicating the presence of four protons geminal to oxygenated carbons of the sapogenol moiety (δ 3.47, dd, J =5.8, 11.6 Hz, H-24; δ 3.54, dd, J =3.6, 2.7 Hz, H-1; δ 3.57 ddd, 10.5, 8.3, 5.0 Hz, H-7; δ 3.73, dd, J =4.3, 12.0 Hz, H-3). The resonances for the oxygenated carbons also indicated the presence of four oxymethine carbons (δ 70.2, d, C-7; δ 73.1, d, C-1; δ 77.0, d, C-24; δ 84.1, d, C-3) and an oxygenated quaternary carbon (δ 80.5, s, C-25).

To complete the assignment of the chemical shifts of the triterpene skeleton and its substitution pattern had to be determined for which task the ¹H–¹H-correlation spectroscopy (COSY) spectrum proved to be most useful. Detailed examination of this spectrum indicated the presence of seven spin systems (Chart 2). In order to establish the interfragment relationship, a heteronuclear multiple-bond correlation experiment (HMBC) was performed (Chart 2). Thus, the anomeric proton at δ 4.36 (d, J =7.8 Hz) showed a long range correlation with the carbon resonance at δ 84.10 (d, C-3) while the anomeric proton at δ 4.54 (d, J =7.8 Hz) exhibited a long range correlation with the carbon resonance at δ 80.50 (s, C-25), indicating the bidesmosidic structure of **1**.



¹³C-NMR data for C-24 is comparable to those reported for analogous compounds having a 24(*S*) configuration.^{15,16)} It has to be mentioned that the ¹³C-NMR data can be regarded as characteristic parameters in the determination of absolute configurations of C-24. In the case of 24(*R*) configuration δ_{C-24} gives resonance at 80.0–80.5 ppm,^{17,18)} while for the 24(*S*) configuration δ_{C-24} gives resonance at 77.0–77.2

* To whom correspondence should be addressed. e-mail: rikhan@olemiss.edu

Table 1. ^1H - and ^{13}C -Assignments of **1**, (in CD_3OD ; at 500 and 125 MHz, Respectively)

C/H	δ (ppm), J (in Hz)	δ (ppm)	C/H	δ (ppm), J (in Hz)	δ (ppm)
1	3.54 dd (2.7, 3.6)	73.1 d	23	1.45 m	28.0 t
2	2.17 m, 1.86 ddd (11.8, 2.7, 1.6)	36.4 t	24	3.47 dd (5.8, 11.6)	77.0 d
3	3.73 dd (12.0, 4.3)	84.1 d	25	—	80.5 s
4	—	40.7 s	26	1.27 s	22.0 q
5	2.16 t (12.0, 4.8)	39.3 d	27	1.23 s	21.6 q
6	1.05 m, 1.75 m	31.0 t	28	1.09 s	24.8 q
7	3.57 ddd (10.5, 8.3, 5.0)	70.2 d	29	0.89 s	13.5 q
8	1.56 m	55.1 d	30	1.07 s	18.2 q
9	—	21.0 s			
10	—	30.4 s	1'	4.36 d (7.8)	105.7 d
11	2.27 m, 1.36 m	26.0 t	2'	3.22 ^{a)}	74.3 d
12	1.70 m	33.0 t	3'	3.37 ^{a)}	77.2 ^{b)} d
13	—	45.9 s	4'	3.30 ^{a)}	70.8 d
14	—	48.9 s	5'	3.28 ^{a)}	76.8 ^{c)} d
15	1.58 m	37.5 t	6'	3.84 dd (10.3, 2.0)	61.8 t
16	1.36 m, 1.96 m	28.6 t		3.65 ^{a)}	
17	1.58 m	52.2 d	1''	4.54 d (7.8)	97.0 d
18	1.05 s	17.4 q	2''	3.20 ^{a)}	74.7 d
19	0.44 d (4.5), 0.79 d (4.5)	28.4 t	3''	3.37 ^{a)}	77.3 ^{b)} d
20	1.45 m	36.2 d	4''	3.30 ^{a,b)}	70.6 d
21	0.95 d (6.3)	17.9 q	5''	3.28 ^{a)}	77.0 ^{c)} d
22	1.56 m, 1.35 m	33.7 t	6''	3.91 dd (10.1, 2.0)	62.1 t
				3.65 ^{a)}	

Assignments confirmed by COSY, HMQC and HMBC experiments. a) Signal pattern was unclear due to overlapping. b, c) Assignments may be interchangeable.

Chart 1. Structure of **1**Chart 2. Partial Structures Deduced from 2D-NMR Measurements (HMQC, COSY) and Key HMBC of **1**

ppm.^{15,16} Additionally, the relative configurations of the oxygenated carbon atoms were determined from the magnitude of the vicinal proton–proton coupling constants to be: C-1 (α -OH; δ 3.54 dd, $J=2.7, 3.6$ Hz, $H_{\text{eq}}-1$), C-3 (β -OH; δ 3.73 dd, $J=12.0, 4.3$ Hz, $H_{\text{ax}}-3$), C-7 (β -OH; δ 3.57 ddd, $J=10.5, 8.3, 5.0$ Hz, $H_{\text{ax}}-7$), and C-24 (δ 3.47 dd, $J=11.6, 5.8$ Hz, H-24).

Consequently, the structure of **1** was established as 3,25-di- O - β -D-glucopyranosyl-1 α ,3 β ,7 β ,24(S),25-pentahydroxy-

cycloartane.

Experimental

General The 1D- and 2D-NMR spectra were obtained on a Bruker® Avance DRX 500 FT spectrometer operating at 500 and 125 MHz, respectively. The chemical shift values are reported as parts per million (ppm) units relative to tetramethylsilane (TMS) for ^1H - and ^{13}C -; and the coupling constants are in Hz (in parentheses). For the ^{13}C -NMR spectra, multiplicities were determined by a distortionless enhancement by polarization transfer (DEPT) experiment. HR-ESI-FT-MS were obtained using a Bruker BioApex

FT-MS in ESI mode.

Chromatographic Conditions TLC: precoated Si 250F plates (Baker); developing system: CHCl₃-MeOH-H₂O (70:30:3); visualization: 30% H₂SO₄. Column chromatography: silica gel 230-400 mesh (Merck).

Plant Material *Astragalus oleifolius* DC. was collected from Ahlatlibel, Ankara, Central Anatolia, Turkey in May 1994. Voucher specimens (94-004) have been deposited at the Herbarium of the Department of Pharmacognosy, Faculty of Pharmacy, Hacettepe University, Ankara, Turkey.

Extraction and Isolation Air-dried and powdered roots of *A. oleifolius* (250 g) were kept in EtOH-H₂O (4:1; 2:1) overnight, then refluxed for 2 h and filtered. The filtrate was concentrated to dryness *in vacuo* (65.0 g, yield 26%). An aliquot of the extract (30.0 g) was partitioned in H₂O and subjected to vacuum liquid chromatography (VLC) using reversed phase material (Sephalyte 40 μm; 250 g), employing H₂O, H₂O-MeOH (90:10→10:90) and MeOH as the eluents. Fractions rich in saponins were combined (7.0 g). This fraction further subjected to column chromatography (silica gel, 150 g) eluted with CH₂Cl₂-MeOH (85:15) and CH₂Cl₂-MeOH-H₂O mixtures (80:20:1; 80:20:2; 70:30:3) to give six main fractions (frs. A-F). Fr. E (166 mg) was subjected to VLC using reversed phase material as stationary phase (Sephalyte 40 μm; 15 g). Elution with increasing amount of MeOH in H₂O (40:60→70:30) yielded compound **1** (11.5 mg).

Compound **1** 3,25-di-*O*-β-D-glucopyranosyl-1α,3β,7β,24(S),25-pentahydroxy-cycloartane: ¹H- and ¹³C-NMR: see Table 1. HR-ESI-FT-MS: [M+Na]⁺ at *m/z* 839.4721.

Acknowledgments We are grateful to Dr. Chuck Dunbar for conducting the HR-ESI-MS analysis. We also thank Mr. Frank Wiggers for his assistance in obtaining the 2D-NMR spectra of **1**. This work was supported in part by the United States Dept. of Agricultural, ARS Specific Cooperative Agreement No. 58-6408-7-012.

References

- 1) Davis P. H., "Flora of Turkey and East Aegean Islands," Vol. 4, University Press, Edinburgh, 1970, pp. 49-254.
- 2) Tang W., Eisenbrand G., "Chinese Drugs of Plant Origin," Springer-Verlag, Berlin, 1992.
- 3) Smee D. F., Verbiscar A. J., *Antiviral Chem. Chemother.*, **6**, 385-390 (1995).
- 4) Yang H., Zhao G., *Zhongguo Zhongliu Linchang*, **25**, 669-672 (1998).
- 5) Liu X., Wang M., Wu H., Zhao X., Li H., *Tianran Chanwu Yanjiu Yu Kaifa*, **6**, 23-31 (1994).
- 6) Isaev M. I., Gorovits M. B., Abubakirov N. K., *Khim. Prir. Soedin.*, **1998**, 156-175.
- 7) Calis I., Zor M., Saracoglu I., Isimer A., Rüggeger H., *J. Nat. Prod.*, **59**, 1019-1023 (1996).
- 8) Bedir E., Calis I., Zerbe O., Sticher O., *J. Nat. Prod.*, **61**, 503-505 (1998).
- 9) Bedir E., Calis I., Aquino R., Piacente S., Pizza C., *J. Nat. Prod.*, **61**, 1469-1472 (1998).
- 10) Bedir E., Calis I., Aquino R., Piacente S., Pizza C., *J. Nat. Prod.*, **62**, 563-568 (1999).
- 11) Bedir E., Calis I., Aquino R., Piacente S., Pizza C., *Phytochemistry*, **51**, 1017-1020 (1999).
- 12) Calis I., Yusufoglu H., Zerbe O., Sticher O., *Phytochemistry*, **50**, 843-847 (1999).
- 13) Calis I., Yuruker A., Tasdemir D., Wright A. D., Sticher O., Luo Y. D., Pezzuto J. M., *Planta Med.*, **63**, 183-186 (1997).
- 14) Agrawal P. K., Jain D. C., Gupta R. K., Thakur R. S., *Phytochemistry*, **24**, 2479-2496 (1985).
- 15) Fadeev Y. M., Isaev M. I., Akimov Y. A., Kintya P. K., Gorovits M. B., Abubakirov N. K., *Khim. Prir. Soedin.*, **1987**, 817-824.
- 16) Hirotoni M., Zhou Y., Rui H., Furuya T., *Phytochemistry*, **37**, 1403-1407 (1994).
- 17) Isaev M. I., Gorovits M. B., Abdullaev N. D., Abubakirov N. K., *Khim. Prir. Soedin.*, **1982**, 458-464.
- 18) Isaev M. I., Gorovits M. B., Abdullaev N. D., Abubakirov N. K., *Khim. Prir. Soedin.*, **1984**, 732-735.