



Correction to: Proposal for a simple algorithm to differentiate adult-onset Still's disease with other fever of unknown origin causes: a longitudinal prospective study

Emre Bilgin¹ · Mutlu Hayran² · Abdulsamet Erden¹ · Berkan Armağan¹ · Alper Sarı¹ · Levent Kılıç¹ · Ali Akdoğan¹ · Ömer Karadağ¹ · Şule Apraş Bilgen¹ · Sedat Kiraz¹ · İhsan Ertenli¹ · Umut Kalyoncu¹

Published online: 19 February 2019

© International League of Associations for Rheumatology (ILAR) 2019

Correction to: Clinical Rheumatology

<https://doi.org/10.1007/s10067-019-04455-y>

The author regrets that the original version of this article contained error. Figure 1 was shown in the wrong version, thus corrected figure is shown in this article.


The online version of the original article can be found at <https://doi.org/10.1007/s10067-019-04455-y>

✉ Umut Kalyoncu
umut.kalyoncu@yahoo.com

¹ Department of Internal Medicine, Division of Rheumatology, Hacettepe University Medical School, Sıhhiye, 06100 Ankara, Turkey

² Department of Internal Medicine, Department of Preventive Oncology, Hacettepe University Medical School, Ankara, Turkey

Fig. 1 Algorithm for discrimination of adult onset Still's disease and other causes of fever of unknown origin. F.U.O.: Fever of unknown origin, UNL: Upper normal limit (for ferritin: 336 ng/ml), PPV: Positive Predictive Value



Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.