



# *PKNOX2* expression and regulation in the bone marrow mesenchymal stem cells of Fanconi anemia patients and healthy donors

Ilgin Cagnan<sup>1,5</sup> · Erdal Cosgun<sup>2,6</sup> · Ozlen Konu<sup>3</sup> · Duygu Uckan<sup>1,4</sup> · Aysen Gunel-Ozcan<sup>1</sup>

Received: 5 September 2018 / Accepted: 22 November 2018 / Published online: 4 December 2018  
© Springer Nature B.V. 2018

## Abstract

HOX and TALE transcription factors are important regulators of development and homeostasis in determining cellular identity. Deregulation of this process may drive cancer progression. The aim of this study was to investigate the expression of these transcription factors in the bone marrow derived mesenchymal stem cells (BM-MSCs) of Fanconi anemia (FA) patients, which is a cancer-predisposing disease. Expression levels of HOX and TALE genes in BM-MSCs were obtained from FA patients and healthy donors by RT-qPCR and highly conserved expression levels were observed between patient and donor cells, except *PKNOX2*, which is a member of TALE class. *PKNOX2* was significantly downregulated in FA cells compared to donors ( $P < 0.05$ ). *PKNOX2* expression levels did not change with diepoxybutane (DEB), a DNA crosslinking agent, in either donor or FA cells except one patient's with a truncation mutation of *FANCA*. A difference of *PKNOX2* protein level was not obtained between FA patient and donor BM-MSCs by western blot analysis. When human TGF- $\beta$ 1 (rTGF- $\beta$ 1) recombinant protein was provided to the cultures, *PKNOX2* as well as *TGF- $\beta$ 1* expression increased both in FA and donor BM-MSCs in a dose dependent manner. 5 ng/mL rTGF- $\beta$  stimulation had more dominant effect on the gene expression of donor BM-MSCs compared to FA cells. Decreased *PKNOX2* expression in FA BM-MSCs may provide new insights into the molecular pathophysiology of the disease and TGF- $\beta$ 1 levels of the microenvironment may be the cause of *PKNOX2* downregulation.

**Keywords** *PKNOX2* · HOX genes · TALE class · TGF- $\beta$ 1 · Fanconi anemia · Bone marrow mesenchymal stem cells

**Electronic supplementary material** The online version of this article (<https://doi.org/10.1007/s11033-018-4522-z>) contains supplementary material, which is available to authorized users.

✉ Aysen Gunel-Ozcan  
agozcan@hacettepe.edu.tr

Ilgin Cagnan  
icagnan@hotmail.com

Erdal Cosgun  
ercosgun@microsoft.com

Ozlen Konu  
konu@fen.bilkent.edu.tr

Duygu Uckan  
duckan@hacettepe.edu.tr

<sup>1</sup> Department of Stem Cell Sciences, Graduate School of Health Sciences, Center for Stem Cell Research and Development, Hacettepe University, Sıhhiye, 06100 Ankara, Turkey

## Introduction

Cellular identity is established during developmental process when cells progressively gain specific lineage properties through the guidance of transcriptional networks.

<sup>2</sup> Department of Biostatistics, Faculty of Medicine, Hacettepe University, Ankara, Turkey

<sup>3</sup> Department of Molecular Biology and Genetics, Bilkent University, Ankara, Turkey

<sup>4</sup> Department of Pediatrics, Division of Bone Marrow Transplantation Unit, Faculty of Medicine, Hacettepe University, Ankara, Turkey

<sup>5</sup> Present Address: Blood Bank, Burhan Nalbantoglu State Hospital, Nicosia, North Cyprus

<sup>6</sup> Present Address: Microsoft Research, 14820 NE 36th Street, Building 99, Redmond, WA 98052, USA

Organ-specifically expressed *HOX* genes encode ‘master regulatory’ homeodomain transcription factors that function in specifying anterior-posterior patterning and establish regional identity during embryonic development [1–3]. In mammals, *HOX* genes are found in four clusters, designated as A, B, C and D, which are located on different chromosomes. DNA binding specificity of HOX proteins is increased through protein–protein interactions with the members of three-amino-acid loop extension (TALE) class homeodomain proteins, known as Meis (*MEIS1*, *MEIS2*, *MEIS3*), Pknox (*PKNOX1*, *PKNOX2*) and Pbx (*PBX1*, *PBX2*, *PBX3*, *PBX4*) gene families [4, 5]. Besides their role in embryonic development, TALE members act as oncogenes (e.g. *MEIS1*) and tumor suppressors (e.g. *PKNOX1*), as well as function in DNA repair and maintain genomic stability (e.g. *PKNOX1*) [6, 7].

During adult life, tightly regulated *HOX* expression pattern continues to provide a “biological fingerprint” for different cell types [8–10]. Loss of cellular identity through alterations in HOX pathway is one of the driving mechanisms of cancer development such as solid tumors and leukemia [11, 12]. Genomic instability in patients (e.g. Fanconi anemia, Wemer syndrome, Bloom syndrome and Ataxia telangiectasia) with defective DNA damage repair pathway (i.e. direct reversal, homologous recombination, non-homologous end joining, mismatch repair, nucleotide excision repair and base excision repair) is also a contributor of cancer progression [13, 14]. We hypothesize that HOX code may change in the diseases with defective DNA repair pathway and predisposition to cancer. To test this hypothesis, we profiled HOX and TALE gene expression in Fanconi anemia (FA) patients, which is a rare inherited disorder with an estimated incidence of 1 in 160,000–360,000 live births [15, 16]. FA patients are characterized with congenital malformations, predisposition to leukemia and solid organ cancers and bone marrow (BM) failure [15]. Mutations in twenty-two different genes, which encode FA complementation group (FANC) proteins and are involved in DNA repair pathway, are responsible for the disease [15, 17]. Moreover, patient cells display hypersensitivity to DNA interstrand crosslinking agents, such as diepoxybutane (DEB), which lead to DNA damage through high levels of chromosomal breaks [18]. The molecular basis of the FA pathophysiology has not been completely elucidated. A study by Zhang et al. shows hyperactive transforming growth factor-beta (TGF- $\beta$ ) signaling as a cause of BM failure in the patients [19]. Members of TGF- $\beta$  signaling pathway is reported to interact with HOX genes [20–23], thus deregulation of TGF- $\beta$  signaling in FA patients may disturb HOX and TALE gene expression as well. Therefore, we also investigated the degree of association between TGF- $\beta$  and

modulation of PKNOX2, which we found out differentially expressed in FA.

## Materials and methods

### Bone marrow mesenchymal stem cells from FA patients and donors

Bone marrow mesenchymal stem cells (BM-MSCs) obtained from FA patients (HUSCS-FA1 -12; n = 12) and donors (HUSCS-D1-16; n = 16) were used. Cells were maintained in DMF10 medium, which contained 100 U/mL penicillin, 100  $\mu$ g/mL streptomycin, 2 mM L-glutamine (Biochrom AG, Germany) and 10% heat-inactivated fetal bovine serum (GIBCO, UK) in a mixture of 60% Dulbecco’s modified Eagle’s medium–low glucose (GIBCO) and 40% MCDB-201 medium (Sigma-Aldrich, USA). Passage 3 BM-MSCs were used in the following experiments. Characterization of BM-MSCs was published previously [24, 25]. Informed consent was obtained from FA patients and donors enrolled in this study. This study was approved by the Local Ethical Committee (Number 14, 24/08/2009) and Hacettepe University Non-interventional Clinical Research Ethics Board (GO 14/403-12, 23/07/2014).

### HOX and TALE gene expression profiling of BM-MSCs from FA patients and donors

Details of RNA isolation, cDNA synthesis and reverse transcriptase quantitative polymerase chain reaction (RT-qPCR) analysis were outlined previously [24, 26]. cDNAs were synthesized from 260 ng RNA samples per 20  $\mu$ l. Expression of 39 HOX and 8 TALE genes were analyzed using RealTime ready Assay (Roche, USA). Target gene expression was normalized against house keeping gene, *ACTB*. Relative gene expression was determined by  $\Delta$ Ct method, calculated by log transformation of  $2^{-\Delta Ct}$ . To enhance the efficiency of statistical analyses, missing  $\Delta$ Ct values were imputed by Multivariate Imputation by Chained Equations (MICE) in R Project for Statistical Computing [27]. MICE method involved assigning a default value for each missing entry. Each column was then updated by appropriate regression or classification algorithm and *Number of Iterations* parameter showed number of times the updates were repeated [28, 29].

### Western blot analysis

The differentially expressed gene obtained by RT-qPCR profiling was also determined at protein level. Total protein lysates from BM-MSCs were prepared using Pierce® RIPA Buffer (Thermo Scientific, USA) containing 1X protease inhibitor cocktail (Sigma-Aldrich). Protein lysates in

Laemmli Buffer (Bio-Rad, USA) containing 355 mM 2-mercaptoethanol (Bio-Rad) were denatured by boiling for 5 min, and then separated by SDS-PAGE, using 10% TGX Stain-Free FastCast Acrylamide kit (Bio-Rad) following manufacturer's protocol. Proteins were transferred to a PVDF membrane by Trans-Blot® Turbo™ Transfer System (7 min, 2.5 A and  $\leq 25$  V; Bio-Rad). Membranes were blocked in TBS containing 0.1% Tween 20 (TBS-T; Bio-Rad) and 5% dry milk (Bio-Rad) for 1 h at room temperature, followed by incubation with 1:100 diluted mouse-anti-PKNOX2 primary antibody (Santa Cruz Biotechnology, USA, Cat# sc-101857) overnight at 4 °C. Membranes were washed with TBS-T, followed by incubation with HRP-goat-anti-mouse secondary antibody (1:2000 dilution; Abclonal, USA) for 1 h at room temperature. Peroxidase activity was measured using Clarity Western ECL Substrate kit (Bio-Rad), following manufacturer's protocol and images were obtained by Kodak Gel Logic 1500 Imaging System (Thermo Fisher Scientific). Membranes were washed, re-blocked, and re-blotted with 1:2500 diluted rabbit-anti- $\beta$ -ACTIN (Cell Signaling Technology, USA, Cat# 8457). Subsequent steps were same as described above, but HRP-goat-anti-rabbit secondary antibody (Abclonal) was used. PKNOX2 protein levels were compared between samples according to signal intensity of PKNOX2 protein bands normalized to loading control  $\beta$ -ACTIN. Densitometry analyses were performed by evaluating band intensity of mean grey value using ImageJ software [30, 31].

### DEB treatment of BM-MSCs

FA cells show sensitivity to DNA interstrand crosslinking agents, such as DEB. Once treated with DEB, patient cells acquire chromosome breaks and undergo cell cycle arrest as well as genomic instability [18]. BM-MSCs derived from FA patients ( $n=6$ ) and donors ( $n=3$ ) were treated with 0.1  $\mu$ g/mL DEB (Sigma-Aldrich) in DMF10 medium, as outlined previously [25]. Untreated cells cultured in DMF10 medium were used as control. cDNAs synthesized from 260 ng RNA samples were used in RT-qPCR analysis to determine the effect of DEB treatment on *PKNOX2* relative expression. When Ct value was not acquired,  $\Delta$ Ct was accepted as  $-25$ .

### Culture of BM-MSCs with recombinant human TGF- $\beta$ 1 protein

BM-MSCs from FA patients or donor were plated into six-well plates and kept in a 5% CO<sub>2</sub> incubator at 37 °C for 24 h. Cells were then induced with 0.1 or 5 ng/mL of recombinant human TGF- $\beta$ 1 protein (rTGF- $\beta$ 1; BioLegend, USA) containing DMF10 medium for 24 h. Uninduced cells maintained in DMF10 medium were included as controls. The effect of rTGF- $\beta$ 1 treatment on *PKNOX2*, *MEIS1*, *PBX1* and

*TGF- $\beta$ 1* expression in BM-MSCs from FA patients ( $n=5$ ) and donors ( $n=5$ ) was determined. Following induction, BM-MSCs were trypsinized in 0.25% trypsin (Invitrogen, UK) containing 1 mM EDTA (Invitrogen) and washed with PBS (Applichem, Germany), followed by RNA isolation and cDNAs synthesis (i.e. 130 ng RNA was used), according to above protocol. Fold change (FC) in gene expression between induced and control cells were calculated by applying a log transformation to  $2^{-\Delta\Delta C_t}$  [32]. The effect of rTGF- $\beta$ 1 induction on *PKNOX2* protein level of BM-MSCs from FA patients ( $n=3$ ) and donors ( $n=3$ ) was determined using western blot analysis, following the above protocol.


### Statistical analysis

Statistical analyses were performed using IBM SPSS Statistical software, V24 and graphics were constructed using GraphPad Prism 7, unless stated otherwise. To compare two independent groups, Student's *t*-test or Mann Whitney *U* (MWU) test was performed and *P*-value less than 0.05 was considered as statistically significant. To compare three dependent groups, Friedman's 2-way ANOVA by ranks test was used and if asymptotic *P*-value was less than 0.05, pairwise test with Bonferroni correction was applied to test the significance within two groups (i.e. adjusted  $P < 0.05$ ). Heatmap (clustering method: single linkage; distance method: Euclidean), as well as scatter-plot showing differentially expressed genes between groups was constructed using Exiqon GenEx qPCR analysis software. Spearman correlation analysis was performed on GraphPad Prism 7 software and was expressed as correlation coefficient (*r*).

## Results

### HOX and TALE profile of BM-MSCs

HOX and TALE genes had a conserved expression between FA patient and donor BM-MSCs. Cells had no *HOXB1* expression, whereas they had low and inconsistent expression of *HOXB13*, *HOXC12*, *HOXD10*, *HOXD11*, *HOXD12* and *HOXD13* (Online Resource 1), thus these genes were excluded from imputation and further analysis. HOX and TALE gene expression was grouped into six clusters (Fig. 1a). The first cluster included *HOXA13*, *HOXB4*, *HOXB8*, *HOXD3*, *HOXD4*, *HOXD9* and *PBX4* ( $\Delta C_{t_{\min}} = -18.79$ ,  $\Delta C_{t_{\max}} = -11.91$ ). The second cluster was consisted of *HOXA1*, *HOXA2*, *HOXA3*, *HOXA4*, *HOXA5*, *HOXA6*, *HOXA7*, *HOXA9*, *HOXA10*, *HOXA11*, *HOXB2*, *HOXB3*, *HOXB5*, *HOXB6*, *HOXB7*, *HOXC4*, *HOXC5*, *HOXC6*, *HOXC8*, *HOXC9*, *HOXC11*, *HOXD8*, *MEIS1*, *MEIS2*, *PBX1*, *PBX2*, *PBX3*, *PKNOX1* and *PKNOX2* ( $\Delta C_{t_{\min}} = -17.59$ ,  $\Delta C_{t_{\max}} = -5.45$ ).



**Fig. 1** HOX and TALE gene expression was conserved between FA patient and donor BM-MSCs. **a** Heat-map illustration of gene expression of BM-MSCs from FA patients ( $n=12$ ) and donors ( $n=16$ ). Dendrograms showed clustering of genes or FA and donor samples. Red color indicates high expression, whereas green color indicates

low gene expression. **b** HOX and TALE gene expression was highly correlated between FA patient ( $n=12$ ) and donor ( $n=16$ ) BM-MSCs. However, *PKNOX2* and *HOXC13* were differentially expressed between groups. (Color figure online)


*HOXD1* ( $\Delta Ct_{\min} = -18.05$ ,  $\Delta Ct_{\max} = -12.73$ ), *HOXB9* ( $\Delta Ct_{\min} = -18.74$ ,  $\Delta Ct_{\max} = -11.25$ ), *HOXC13* ( $\Delta Ct_{\min} = -18.97$ ,  $\Delta Ct_{\max} = -13.90$ ) or *HOXC10* ( $\Delta Ct_{\min} = -7.10$ ,  $\Delta Ct_{\max} = -4.03$ ) were clustered alone (Fig. 1a). Additionally, *HOXC10* had the highest relative expression level in all BM-MSCs (Fig. 1a). Correlation analysis revealed that gene expression was highly associated ( $r=0.9861$ ,  $P<0.0001$ ) between FA patients and donors, while *PKNOX2* and *HOXC13* were differentially expressed between groups (Fig. 1b).

Statistical analysis revealed that expression of HOX (Fig. 2a–d) genes, as well as *MEIS1*, *MEIS2*, *PBX1*, *PBX2*, *PBX3*, *PBX4* and *PKNOX1* (Fig. 2e) were not significantly ( $P>0.05$ ) different between groups. However, *PKNOX2* expression of FA patients ( $-15.19 \pm 1.49$ ) was significantly lower than ( $P<0.05$ ) the expression of donors ( $-13.24 \pm 1.37$ ; Fig. 2e). DEB treatment had no effect on *PKNOX2* relative

expression levels of FA patients and donors, except one patient's BM-MSCs (HUSCS-FA04) which lost the expression of *PKNOX2* ( $\Delta Ct = -25$ ) by DEB treatment (Fig. 2f). Western blot analysis revealed that BM-MSCs probably had more than one *PKNOX2* isoform (Fig. 3a). When variants corresponding to 70 kDa (i.e. large) and 52 kDa (i.e. small) were quantified, cells had higher levels of large variant compared to small one (Fig. 3b). Additionally, level of large and small *PKNOX2* isoforms did not differ ( $P>0.05$ ) between FA patient ( $1.21 \pm 0.28$  and  $0.30 \pm 0.07$ , respectively) and donor cells ( $1.19 \pm 0.33$  and  $0.24 \pm 0.06$ , respectively; Fig. 3b).

### Effect of rTGF- $\beta$ 1 induction on *PKNOX2* and TGF- $\beta$ 1 levels

For each experimental condition (i.e. control, 0.1 or 5 ng/mL rTGF- $\beta$ 1 protein), fold change in gene expressions of both




**Fig. 2** Relative expression of **a** HOXA, **b** HOXB, **c** HOXC, **d** HOXD, as well as **e** TALE class genes were highly conserved between FA (n=12) and donor (n=16) BM-MSCs. However, *PKNOX2* expression of FA BM-MSCs was significantly lower than donor cells. Data are shown as means  $\pm$  standard deviation (SD). Asterisk (\*) depicted

statistically significant difference ( $P < 0.05$ ). **f** DEB treatment had no effect on *PKNOX2* expression of FA (n=6) and donor (n=3) BM-MSCs. BM-MSCs from a patient (HUSCS-FA04) had no expression of *PKNOX2* ( $\Delta Ct = -25$ ) following DEB treatment

FA and donor BM-MSCs were upregulated as the dose of rTGF- $\beta$ 1 increased (Fig. 4a, b). When compared to their corresponding uninduced controls, increase in *PKNOX2* expression was significant in both FA ( $\text{Log}_2 \text{FC} = 2.37 \pm 0.84$ ) and donor ( $\text{Log}_2 \text{FC} = 3.09 \pm 0.58$ ) BM-MSCs induced with 5 ng/mL rTGF- $\beta$ 1 protein (adjusted  $P < 0.05$ ; Fig. 4a). The same dose also provided a significant increase in *TGF- $\beta$ 1* expression of the donor BM-MSCs ( $\text{Log}_2 \text{FC} = 1.10 \pm 0.16$ ;

adjusted  $P < 0.05$ ; Fig. 4b), but not of the FA BM-MSCs. Fold change differences in either *PKNOX2* (Fig. 4a) or *TGF- $\beta$ 1* (Fig. 4b) were not significant when FA patients compared to donors ( $P > 0.05$ ). rTGF- $\beta$ 1 treatment of BM-MSCs did not alter *MEIS1* (Fig. 4c) or *PBX1* (Fig. 4d) expression levels ( $P > 0.05$ ).

Level of PKNOX2 protein was determined prior to and after induction with 0.1 and 5 ng/mL rTGF- $\beta$ 1 protein



**Fig. 3** PKNOX2 protein level of BM-MSCs derived from FA patients ( $n=7$ ) and donors ( $n=6$ ) was not significantly different ( $P>0.05$ ). BM-MSCs expressed two different PKNOX2 isoforms (70 and 52 kDa). **a** Protein samples (45  $\mu$ g per lane) were run on two different

10% SDS-PAGE gels and the pictures were taken at the same time (exposure time = 5 min). **b** Ratio of PKNOX2 isoforms normalized to  $\beta$ -ACTIN was calculated


(Fig. 5). All samples had higher level of large variant (70 kDa) compared to small isoform (52 kDa; Fig. 5a). PKNOX2 protein level remained unchanged ( $P>0.05$ ) between FA and donor BM-MSCs at any experimental condition (Fig. 5b). Additionally, PKNOX2 protein level within either FA patients or donors did not change significantly ( $P>0.05$ ) upon induction (Fig. 5b).

## Discussion

The molecular signature of HOX expression is organ-specific [8, 9]. Changes in HOX pathway may result in alterations in the cellular identity and trigger cancer progression [11, 12]. One of the aims of this study was to assess, whether HOX and TALE gene expression pattern changed in diseases predisposed to cancer, like FA. At the third passage, FA BM-MSCs had comparable HOX gene expression levels with donor cells, but *HOXC13* expression was relatively lower in patients. In conjunction with other studies, we also found that most HOX genes, except *HOXB1*, *HOXB13*, *HOXC12*, *HOXD10*, *HOXD11*, *HOXD12* and *HOXD13*, were actively expressed by BM-MSCs [33, 34]. Expression of *HOXA9*, *HOXA10*, *HOXC6*, *HOXC8*, *HOXC10* and *HOXD8* in

BM-MSCs is known to be higher than other HOX genes, as also observed in our study [33, 34]. Liedtke and co-authors (2010) have reported that *HOXA3*, *HOXA11*, *HOXA13*, *HOXB2*, *HOXB3*, *HOXB8*, *HOXB9*, *HOXC11*, *HOXC13* and *HOXD1* expression were absent in BM-MSCs [33], but we observed that they were expressed at low to intermediate level.

In our study, BM-MSCs also actively expressed members of TALE gene class. Intriguingly, BM-MSCs derived from FA patients had significantly lower *PKNOX2* expression compared to donors. We analyzed two different GEO datasets (GSE61853 and GSE87806) containing gene expression profile of bone marrow mesenchymal stromal cells from other bone marrow diseases (myelodysplastic syndrome—MDS-, polycythemia vera—PV-, and essential thrombocythemia—ET-, chronic myeloid leukemia—CML-) to test whether *PKNOX2* expression level changed under other disease states. We found that *PKNOX2* expression of mesenchymal stromal cells from MDS, PV, ET or CML patients did not significantly differ from controls (adjusted  $P>0.05$ ; Online Resource 2) [35, 36]. According to the results of these datasets, decrease in *PKNOX2* expression is restricted to FA patients. Additionally, it would be intriguing to compare, whether *PKNOX2* expression changes upon




**Fig. 4** Fold change in **a** *PKNOX2*, **b** *TGF-β1*, **c** *MEIS1* and **d** *PBX1* expression of FA (n=5) and donor (n=5) BM-MSCs induced with 0.1 or 5 ng/mL rTGF-β1 for 24 h was determined. Increase in *PKNOX2* expression was significant in both FA and donor BM-MSCs induced with 5 ng/mL rTGF-β1 protein compared to their corresponding controls (adjusted  $P < 0.05$ ). *TGF-β1* expression of

the donor BM-MSCs induced with 5 ng/mL recombinant protein was also significantly higher than uninduced donor cells (adjusted  $P < 0.05$ ), while the expression level in FA BM-MSCs fluctuated within individuals. Asterisk (\*) depicted statistically significant difference ( $P < 0.05$ )

cell passaging. Due to their low frequency (0.001–0.01% of nucleated cells) in the bone marrow [37], BM-MSCs have to be expanded in-vitro, which prevents the use of fresh explants. However, passaging BM-MSCs for a long-term trigger senescence and affect their proliferative capacity [38, 39]. Also, FA BM-MSCs have defects in their proliferation capacity and undergo senescence in-vitro [40]. Therefore, the passage number is one of the limitations of this study.

Unlike its mRNA level, PKNOX2 protein level did not change between FA patients and healthy donors. Protein and mRNA levels of a gene may not always correlate with each other due to post-transcriptional modifications or half-lives of proteins [41]. Herein, we obtained expression of two different PKNOX2 isoforms in BM-MSCs, corresponding to 70 and 52 kDa. On the SDS-PAGE gel, smaller variant had the expected molecular weight of PKNOX2, while the molecular weight of large isoform corresponded to the in-vitro synthesized protein by Fognani et al. [42]. Similar to our findings, NIH3T3 mouse embryo fibroblast cell line is shown to have variants of PKNOX2 protein [43]. Clear function of PKNOX2 is not well understood apart from its

role in regulation of transcription through sequence-specific DNA binding and actin filament/monomer binding. Furthermore, Pknox2 overexpression in mice limb bud mesenchyme results in hypoplastic radius and ulna, which are common defects observed in FA patients [44]. *PKNOX2* has a high structural similarity to its paralogous gene *PKNOX1*, which is known to function as a tumor suppressor gene with roles in DNA repair and maintenance of genomic stability [6, 7, 42]. This might implicate that *PKNOX2* could also be a potential player in the DNA repair of FA stromal environment. Indeed, a whole genome RNA interference (RNAi) study showed that *PKNOX2* silencing increased cellular sensitivity to ionizing radiation [45]. However, our study showed that DEB treatment of BM-MSCs did not change the expression of *PKNOX2* in either donor or FA patients, except one that possessed a novel deletion of exon 1–2 in *FANCA* gene, reported in our previous study [25]. *PKNOX2* expression is lost by DEB treatment in that patient's BM-MSCs. Truncation mutation of that patient is probable to be more deleterious and can increase cellular sensitivity to cross linking agents such as DEB by *PKNOX2* silencing or



**Fig. 5** rTGF- $\beta$ 1 protein induction had no effect on PKNOX2 protein level of FA ( $n=3$ ) and donor ( $n=3$ ) BM-MSCs. **a** Protein samples (20  $\mu$ g per lane) were run on two different 10% SDS-PAGE gels and the pictures were taken at the same time (exposure time=5 min). **b** Ratio of PKNOX2 protein isoforms normalized to  $\beta$ -ACTIN were

calculated. PKNOX2 protein level did not differ ( $P>0.05$ ) between FA and donor BM-MSCs at any experimental condition. Besides, PKNOX2 protein level within either FA patients or donors groups remained unchanged ( $P>0.05$ ) upon induction

increased cellular sensitivity to DEB may be the cause of *PKNOX2* silencing. It is also possible that *FANCA* exon 1–2 is required for *PKNOX2* expression, which should be tested by further functional assays.

From many ( $n=1639$ ) transcription factors found in humans [46], we focused on HOX and TALE transcription factors that are strictly under epigenetic control during adult life. TGF- $\beta$  signaling interacts with HOX genes [20–23], and we previously showed fluctuation of TGF- $\beta$  secretion from FA BM-MSCs [25]. Deregulated TGF- $\beta$  signaling may disturb *PKNOX2* expression in FA BM-MSCs and trigger disease progression, as seen in FA HSCs [19]. Dose- and time-dependent effects of TGF- $\beta$ 1 on cell cultures are well known [47, 48]. We performed the preliminary experiment by stimulating BM-MSCs from a donor with 0.1 or 5 ng/mL rTGF- $\beta$ 1 protein for 24, 48 or 72 h (Online Resource 3). Following 24 h of incubation, *TGF- $\beta$ 1* expression increased linearly in a dose-dependent manner (Online Resource 3), thus further experiments were only performed on this timepoint. Additionally, 5 ng/mL rTGF- $\beta$ 1 protein was the maximum induction dose used, because higher concentrations

stimulate chondrogenic differentiation [48]. Wu et al. also show that increase in TGF- $\beta$ 1 concentration increases senescence activity of BM-MSCs [47].

*PKNOX2* expression of both FA and donor BM-MSCs were increased by rTGF- $\beta$ 1 in a similar dose-dependent manner, suggesting TGF- $\beta$ 1 signaling may not be perturbed in FA BM-MSCs. Our data confirm the results of an expression microarray study deposited to GEO database (GSE46019) that shows an increase in *PKNOX2* expression of BM-MSCs following TGF- $\beta$ 1 stimulation [49]. Also, Zhou et al. report that overexpressed *Pknox2* decreases p-Smad1/5/8 levels in mice [44]. All in all, our data in conjunction with these studies suggest that *PKNOX2* and TGF- $\beta$  signaling pathway are associated with each other. Secondly, we investigated whether change in rTGF- $\beta$ 1 level altered the expression of any other TALE factors. To answer this question, MEIS1, an oncogenic transcription factor, as well as its cofactor, PBX1, were chosen [7, 50]. Stimulation of BM-MSCs did not affect the mRNA level of these genes, thus the dose-dependent effect of rTGF- $\beta$ 1 treatment is possibly constrained to *PKNOX2* expression. Moreover, this



study displayed that rTGF- $\beta$ 1 treatment up-regulated *TGF- $\beta$ 1* expression of BM-MSCs in a dose-dependent manner, confirming a positive feedback loop shown previously [51]. It was intriguing that the dose dependence of this loop was slightly disrupted in the BM-MSCs of FA patients. Although mouse models of FA do not resemble the complete characteristics of patients [52], it will be important to investigate whether in-vitro effect of rTGF- $\beta$ 1 treatment on the gene expression of BM-MSCs could be correlated with in-vivo studies.

In conclusion, *PKNOX2* expression was downregulated in FA patient BM-MSCs compared to controls. Our results suggest that fluctuation in TGF- $\beta$ 1 levels may change *PKNOX2* expression. Being one of the important members of bone marrow microenvironment, MSCs with deregulated *PKNOX2* expression may impair the function of niche and would contribute to hematopoietic defects seen in FA patients, which needs to be elucidated further with functional analysis.

**Acknowledgements** This study was supported by The Scientific and Technological Research Council of Turkey (TUBITAK; Project No: 110S021 in conjunction with EU COST Action BM0805 designated as ‘HOX and TALE transcription factors in Development and Disease’ and TUBITAK Project No: 214Z033). The data in this study is a part of Ilgin Cagnan’s Ph.D. thesis.

## Compliance with ethical standards

**Conflict of interest** The authors declare that they have no conflict of interest.

## References

- Illig R, Fritsch H, Schwarzer C (2013) Spatio-temporal expression of HOX genes in human hindgut development. *Dev Dyn* 242:53–66. <https://doi.org/10.1002/dvdy.23893>
- McGinnis W, Krumlauf R (1992) Homeobox genes and axial patterning. *Cell* 68:283–302
- Williamson I, Eskeland R, Lettice LA et al (2012) Anterior-posterior differences in HoxD chromatin topology in limb development. *Development* 139:3157–3167. <https://doi.org/10.1242/dev.081174>
- Holland PW, Booth HA, Bruford EA (2007) Classification and nomenclature of all human homeobox genes. *BMC Biol* 5:47
- Merabet S, Mann RS (2016) To Be Specific or Not: The Critical Relationship Between Hox And TALE Proteins. *Trends Genet* 32:334–347. <https://doi.org/10.1016/j.tig.2016.03.004>
- Iotti G, Longobardi E, Masella S et al (2011) Homeodomain transcription factor and tumor suppressor Prep1 is required to maintain genomic stability. *Proc Natl Acad Sci U S A* 108:E314–E322. <https://doi.org/10.1073/pnas.1105216108>
- Thorsteinsdottir U, Kroon E, Jerome L, Blasi F, Sauvageau G (2001) Defining roles for HOX and MEIS1 genes in induction of acute myeloid leukemia. *Mol Cell Biol* 21:224–234
- Ackema KB, Charite J (2008) Mesenchymal stem cells from different organs are characterized by distinct topographic Hox codes. *Stem Cells Dev* 17:979–991. <https://doi.org/10.1089/scd.2007.0220>
- Takahashi Y, Hamada J, Murakawa K et al (2004) Expression profiles of 39 HOX genes in normal human adult organs and anaplastic thyroid cancer cell lines by quantitative real-time RT-PCR system. *Exp Cell Res* 293:144–153
- Yamamoto M, Takai D, Yamamoto F (2003) Comprehensive expression profiling of highly homologous 39 hox genes in 26 different human adult tissues by the modified systematic multiplex RT-pCR method reveals tissue-specific expression pattern that suggests an important role of chromosomal structure in the regulation of Hox gene expression in adult tissues. *Gene Expr* 11:199–210
- Bhatlekar S, Fields JZ, Boman BM (2014) HOX genes and their role in the development of human cancers. *J Mol Med* 92:811–823. <https://doi.org/10.1007/s00109-014-1181-y>
- Platais C, Hakami F, Darda L, Lambert DW, Morgan R, Hunter KD (2016) The role of HOX genes in head and neck squamous cell carcinoma. *J Oral Pathol Med* 45:239–247. <https://doi.org/10.1111/jop.12388>
- Bhattacharjee S, Nandi S (2018) Rare genetic diseases with defects in DNA repair: opportunities and challenges in orphan drug development for targeted cancer therapy. *Cancers* 10(298). <https://doi.org/10.3390/cancers10090298>
- Hakem R (2008) DNA-damage repair; the good, the bad, and the ugly. *EMBO J* 27:589–605. <https://doi.org/10.1038/emboj.2008.15>
- Mamrak NE, Shimamura A, Howlett NG (2016) Recent discoveries in the molecular pathogenesis of the inherited bone marrow failure syndrome Fanconi anemia. *Blood Rev* 31:93–99. <https://doi.org/10.1016/j.blre.2016.10.002>
- Zhan-He W (2013) The concept and practice of Fanconi Anemia: from the clinical bedside to the laboratory bench. *Transl Pediatr* 2:112–119. <https://doi.org/10.3978/j.issn.2224-4336.2013.07.01>
- Knies K, Inano S, Ramirez MJ et al (2017) Biallelic mutations in the ubiquitin ligase RFWF3 cause Fanconi anemia. *J Clin Invest* 127:3013–3027. <https://doi.org/10.1172/JCI92069>
- Auerbach AD (1993) Fanconi anemia diagnosis and the diepoxybutane (DEB) test. *Exp Hematol* 21:731–733
- Zhang H, Kozono DE, O’Connor KW et al (2016) TGF-beta Inhibition rescues hematopoietic stem cell defects and bone marrow failure in Fanconi anemia. *Cell Stem Cell* 18:668–681. <https://doi.org/10.1016/j.stem.2016.03.002>
- Wang N, Kim HG, Cotta CV et al (2006) TGFbeta/BMP inhibits the bone marrow transformation capability of Hoxa9 by repressing its DNA-binding ability. *EMBO J* 25:1469–1480
- Walsh CM, Carroll SB (2007) Collaboration between Smads and a Hox protein in target gene repression. *Development* 134:3585–3592
- Ruiz i Altaba A, Melton DA (1989) Interaction between peptide growth factors and homeobox genes in the establishment of antero-posterior polarity in frog embryos. *Nature* 341:33–38
- Kloen P, Visker MH, Olijve W, van Zoelen EJ, Boersma CJ (1997) Cell-type-specific modulation of Hox gene expression by members of the TGF-beta superfamily: a comparison between human osteosarcoma and neuroblastoma cell lines. *Biochem Biophys Res Commun* 233:365–369
- Cagnan I (2018) HOX and TALE transcription factors in Fanconi anemia bone-marrow mesenchymal stem cells: gene expression and protein interactions. Dissertation, Hacettepe University
- Cagnan I, Gunel-Ozcan A, Aerts-Kaya F et al (2018) Bone marrow mesenchymal stem cells carrying FANCD2 mutation differ from the other Fanconi anemia complementation groups in terms of TGF-beta1 production. *Stem Cell Rev* 14:425–437. <https://doi.org/10.1007/s12015-017-9794-5>

26. Cagnan I, Aerts Kaya F, Cetinkaya D, Gunel Ozcan G (2017) Stably expressed reference genes during differentiation of bone marrow-derived mesenchymal stromal cells. *Turkish J Biol* 41:88–97. <https://doi.org/10.3906/biy-1511-93>
27. Van Buuren S, Groothuis-Oudshoorn K (2011) mice: Multivariate Imputation by Chained Equations in R. *J Statistical Software* 45:1–67. <https://doi.org/10.18637/jss.v045.i03>
28. Azur MJ, Stuart EA, Frangakis C, Leaf PJ (2011) Multiple imputation by chained equations: what is it and how does it work? *Int J Methods Psychiatr Res* 20:40–49. <https://doi.org/10.1002/mpr.329>
29. Waljee AK, Mukherjee A, Singal AG et al (2013) Comparison of imputation methods for missing laboratory data in medicine. *BMJ Open* 3. <https://doi.org/10.1136/bmjopen-2013-002847>
30. Davarinejad H (2017) Quantifications of western blots with Image J. <http://www.yorku.ca/yisheng/Internal/Protocols/ImageJ.pdf>. (Accessed 01 October 2017)
31. Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* 9:671–675
32. Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>(-delta delta C(T))</sup> method. *Methods* 25:402–408
33. Liedtke S, Buchheiser A, Bosch J et al (2010) The HOX Code as a “biological fingerprint” to distinguish functionally distinct stem cell populations derived from cord blood. *Stem Cell Res* 5:40–50. <https://doi.org/10.1016/j.scr.2010.03.004>
34. Picchi J, Trombi L, Spugnese L et al (2013) HOX and TALE signatures specify human stromal stem cell populations from different sources. *J Cell Physiol* 228:879–889. <https://doi.org/10.1002/jcp.24239>
35. Kim M, Hwang S, Park K, Kim SY, Lee YK, Lee DS (2015) Increased expression of interferon signaling genes in the bone marrow microenvironment of myelodysplastic syndromes. *PLoS ONE* 10:e0120602. <https://doi.org/10.1371/journal.pone.0120602>
36. Ramos TL, Sánchez-Abarca LI, Rosón-Burgo B et al (2017) Mesenchymal stromal cells (MSC) from JAK2 + myeloproliferative neoplasms differ from normal MSC and contribute to the maintenance of neoplastic hematopoiesis. *PLoS ONE* 12:e0182470. <https://doi.org/10.1371/journal.pone.0182470>
37. Pittenger MF, Mackay AM, Beck SC et al (1999) Multilineage potential of adult human mesenchymal stem cells. *Science* 284:143–147
38. Baxter M, Wynn RF, Jowitt SN, Wraith JE, Fairbairn LJ, Bellantuono I (2004) Study of telomere length reveal rapid aging of human marrow stromal cells following in vitro expansion. *Stem Cells* 22:675–682
39. Wagner W, Horn P, Castoldi M, Diehlmann A, Bork S, Saffrich R, Benes V, Blake J, Pfiser S, Ecstein V et al (2008) Replicative senescence of mesenchymal stem cells: a continuous and organized process. *PLoS ONE* 3. <https://doi.org/10.1371/journal.pone.0002213>
40. Xu J, Li X, Cole A, Sherman Z, Du W (2018) Reduced cell division control protein 42 activity compromises hematopoiesis-supportive function of Fanconi anemia mesenchymal stromal cells. *Stem Cells* 36:785–795. <https://doi.org/10.1002/stem.2789>
41. Greenbaum D, Colangelo C, Williams K, Gerstein M (2003) Comparing protein abundance and mRNA expression levels on a genomic scale. *Genome Biol* 4:117
42. Fognani C, Kilstrup-Nielsen C, Berthelsen J, Ferretti E, Zappavigna V, Blasi F (2002) Characterization of PREP2, a paralog of PREP1, which defines a novel sub-family of the MEINOX TALE homeodomain transcription factors. *Nucleic Acids Res* 30:2043–2051
43. Haller K, Rambaldi I, Daniels E, Featherstone M (2004) Subcellular localization of multiple PREP2 isoforms is regulated by actin, tubulin, and nuclear export. *J Biol Chem* 279:49384–49394
44. Zhou W, Zhu H, Zhao J et al (2013) Misexpression of Pknox2 in mouse limb bud mesenchyme perturbs zeugopod development and deltoid crest formation. *PloS ONE* 8:e64237. <https://doi.org/10.1371/journal.pone.0064237>
45. Hurov KE, Cotta-Ramusino C, Elledge SJ (2010) A genetic screen identifies the Triple T complex required for DNA damage signaling and ATM and ATR stability. *Genes Dev* 24:1939–1950. <https://doi.org/10.1101/gad.1934210>
46. Lambert SA, Jolma A, Campitelli LF et al (2018) The human transcription factors. *Cell* 172:650–665. <https://doi.org/10.1016/j.cell.2018.01.029>
47. Wu J, Niu J, Li X, Wang X, Guo Z, Zhang F (2014) TGF-β1 induces senescence of bone marrow mesenchymal stem cells via increase of mitochondrial ROS production. *BMC Dev Biol* 14:21. <https://doi.org/10.1186/1471-213X-14-21>
48. Schedel J, Lowin T, Kujat R et al (2010) RAP-PCR fingerprinting reveals time-dependent expression of matrix-related molecules following stem-cell based TGFβ1-induced chondrocyte development. *Int J Mol Med* 27:519–523. <https://doi.org/10.3892/ijmm.2011.608>
49. Walenda G, Abnaof K, Jousen S et al (2013) TGF-beta1 does not induce senescence of multipotent mesenchymal stromal cells and has similar effects in early and late passages. *PLoS ONE* 8:e77656. <https://doi.org/10.1371/journal.pone.0077656>
50. Dardaei L, Longobardi E, Blasi F (2014) Prep1 and Meis1 competition for Pbx1 binding regulates protein stability and tumorigenesis. *Proc Natl Acad Sci U S A* 111:E896–E905. <https://doi.org/10.1073/pnas.1321200111>
51. Li J, Tripathi BJ, Chalam KV, Tripathi RC (1996) Transforming growth factor-beta 1 and -beta 2 positively regulate TGF-beta 1 mRNA expression in trabecular cells. *Invest Ophthalmol Vis Sci* 37:2778–2782
52. Bakker ST, de Winter JP, te Riele H (2013) Learning from a paradox: recent insight into Fanconi anaemia through studying mouse models. *Dis Model Mech* 6:40–47. <https://doi.org/10.1002/stem.2789>