

# catena-Poly[[aquabis(3-chlorobenzoato- $\kappa^2O,O'$ )cadmium]- $\mu$ -*N,N*-diethylnicotinamide- $\kappa^2N^1:O$ ]

Nihat Bozkurt,<sup>a</sup> Tuncay Tunc,<sup>b</sup> Nagihan Çaylak Delibaş,<sup>c</sup> Hacı Necefoğlu<sup>a</sup> and Tuncer Hökelek<sup>d\*</sup>

<sup>a</sup>Department of Chemistry, Kafkas University, 36100 Kars, Turkey, <sup>b</sup>Aksaray University, Science Education Department, 68100, Aksaray, Turkey, <sup>c</sup>Department of Physics, Sakarya University, 54187 Esentepe, Sakarya, Turkey, and <sup>d</sup>Department of Physics, Hacettepe University, 06800 Beytepe, Ankara, Turkey  
Correspondence e-mail: merzifon@hacettepe.edu.tr


Received 11 July 2013; accepted 16 July 2013

Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(C-C) = 0.006$  Å;  $R$  factor = 0.046;  $wR$  factor = 0.093; data-to-parameter ratio = 20.0.

In the crystal of the title  $Cd^{II}$  polymeric complex,  $[Cd(C_7H_4ClO_2)_2(C_{10}H_{14}N_2O)(H_2O)]_n$ , the  $Cd^{II}$  cation is chelated by two chlorobenzoate anions and coordinated by two *N,N*-diethylnicotinamide (DNA) ligands and one water molecule in a distorted  $NO_6$  pentagonal–bipyramidal geometry. The  $Cd^{II}$  cations are bridged by the pyridine N atom and carbonyl O atom of the DNA ligand to form a polymeric chain running along the  $b$  axis. Intermolecular  $O-H \cdots O$  hydrogen bonds between coordinating water molecules and carboxylate groups link adjacent chains into layers parallel to the  $bc$  plane.  $\pi-\pi$  contacts between benzene rings [shortest centroid–centroid distance = 3.912 (2) Å] further stabilizes the crystal structure. In the molecule, weak  $C-H \cdots O$  hydrogen bonds occur between the pyridine ring and carboxylate groups; the dihedral angles between the carboxylate groups and adjacent benzene rings are 4.6 (3) and 12.8 (3)°, while the benzene rings are oriented at a dihedral angle of 1.89 (13)°.

## Related literature

For niacin, see: Krishnamachari (1974). For *N,N*-diethylnicotinamide, see: Bigoli *et al.* (1972). For related structures, see: Çaylak Delibaş *et al.* (2013); Greenaway *et al.* (1984); Hökelek *et al.* (1995); Hökelek & Necefoğlu (1996); Hökelek *et al.* (2009*a,b,c,d,e,f,g*); Hökelek *et al.* (2011); Necefoğlu *et al.* (2010*a,b*); Sertçelik *et al.* (2013).



## Experimental

### Crystal data

$[Cd(C_7H_4ClO_2)_2(C_{10}H_{14}N_2O)(H_2O)]_n$   
 $M_r = 619.76$   
Monoclinic,  $C2/c$   
 $a = 25.1809$  (5) Å  
 $b = 7.0161$  (3) Å  
 $c = 30.6755$  (6) Å

$\beta = 106.203$  (3)°  
 $V = 5204.2$  (3) Å<sup>3</sup>  
 $Z = 8$   
Mo  $K\alpha$  radiation  
 $\mu = 1.09$  mm<sup>-1</sup>  
 $T = 296$  K  
 $0.35 \times 0.15 \times 0.10$  mm

### Data collection

Bruker SMART BREEZE CCD diffractometer  
Absorption correction: multi-scan (SADABS; Bruker, 2012)  
 $T_{min} = 0.823$ ,  $T_{max} = 0.897$

100329 measured reflections  
6531 independent reflections  
6170 reflections with  $I > 2\sigma(I)$   
 $R_{int} = 0.026$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$   
 $wR(F^2) = 0.093$   
 $S = 1.35$   
6531 reflections  
326 parameters  
4 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{max} = 0.77$  e Å<sup>-3</sup>  
 $\Delta\rho_{min} = -1.02$  e Å<sup>-3</sup>

**Table 1**

Selected bond lengths (Å).

Cd1–O1	2.504 (3)	Cd1–O5	2.410 (3)
Cd1–O2	2.323 (3)	Cd1–O6	2.314 (3)
Cd1–O3	2.421 (3)	Cd1–N1	2.305 (3)
Cd1–O4	2.360 (3)		

**Table 2**

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O6–H61 $\cdots$ O2 <sup>i</sup>	0.85 (4)	1.94 (4)	2.753 (5)	160 (4)
O6–H62 $\cdots$ O4 <sup>i</sup>	0.86 (4)	2.11 (4)	2.838 (4)	142 (5)
C15–H15 $\cdots$ O1	0.93	2.52	3.181 (5)	128
C19–H19 $\cdots$ O3	0.93	2.47	3.130 (5)	128

Symmetry code: (i)  $-x + 1, -y + 2, -z$ .

Data collection: APEX2 (Bruker, 2012); cell refinement: SAINT (Bruker, 2012); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics:

ORTEP-3 for Windows (Farrugia, 2012); software used to prepare material for publication: WinGX (Farrugia, 2012) and PLATON (Spek, 2009).

The authors acknowledge the Aksaray University, Science and Technology Application and Research Center, Aksaray, Turkey, for the use of the Bruker SMART BREEZE CCD diffractometer (purchased under grant No. 2010K120480 of the State of Planning Organization).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5721).

## References

- Bigoli, F., Braibanti, A., Pellinghelli, M. A. & Tiripicchio, A. (1972). *Acta Cryst.* **B28**, 962–966.
- Bruker (2012). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Çaylak Delibaş, N., Necefoğlu, H. & Hökelek, T. (2013). *Acta Cryst.* **E69**, m191–m192.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Greenaway, F. T., Pezeshk, A., Cordes, A. W., Noble, M. C. & Sorenson, J. R. J. (1984). *Inorg. Chim. Acta*, **93**, 67–71.
- Hökelek, T., Dal, H., Tercan, B., Aybirdi, Ö. & Necefoğlu, H. (2009c). *Acta Cryst.* **E65**, m1582–m1583.
- Hökelek, T., Dal, H., Tercan, B., Aybirdi, Ö. & Necefoğlu, H. (2009e). *Acta Cryst.* **E65**, m627–m628.
- Hökelek, T., Dal, H., Tercan, B., Aybirdi, Ö. & Necefoğlu, H. (2009f). *Acta Cryst.* **E65**, m1037–m1038.
- Hökelek, T., Dal, H., Tercan, B., Aybirdi, Ö. & Necefoğlu, H. (2009g). *Acta Cryst.* **E65**, m1365–m1366.
- Hökelek, T. & Necefoğlu, H. (1996). *Acta Cryst.* **C52**, 1128–1131.
- Hökelek, T., Necefoğlu, H. & Balcı, M. (1995). *Acta Cryst.* **C51**, 2020–2023.
- Hökelek, T., Sağlam, E. G., Tercan, B., Aybirdi, Ö. & Necefoğlu, H. (2011). *Acta Cryst.* **E67**, m28–m29.
- Hökelek, T., Yılmaz, F., Tercan, B., Aybirdi, Ö. & Necefoğlu, H. (2009a). *Acta Cryst.* **E65**, m955–m956.
- Hökelek, T., Yılmaz, F., Tercan, B., Aybirdi, Ö. & Necefoğlu, H. (2009b). *Acta Cryst.* **E65**, m1328–m1329.
- Hökelek, T., Yılmaz, F., Tercan, B., Gürgen, F. & Necefoğlu, H. (2009d). *Acta Cryst.* **E65**, m1416–m1417.
- Krishnamachari, K. A. V. R. (1974). *Am. J. Clin. Nutr.* **27**, 108–111.
- Necefoğlu, H., Çimen, E., Tercan, B., Dal, H. & Hökelek, T. (2010a). *Acta Cryst.* **E66**, m334–m335.
- Necefoğlu, H., Çimen, E., Tercan, B., Dal, H. & Hökelek, T. (2010b). *Acta Cryst.* **E66**, m485–m486.
- Sertçelik, M., Çaylak Delibaş, N., Necefoğlu, H. & Hökelek, T. (2013). *Acta Cryst.* **E69**, m290–m291.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

## supporting information

*Acta Cryst.* (2013). E69, m466–m467 [doi:10.1107/S160053681301965X]

## **catena-Poly[[aquabis(3-chlorobenzoato- $\kappa^2O,O'$ )cadmium]- $\mu$ -*N,N*-diethyl-nicotinamide- $\kappa^2N^1:O$ ]**

**Nihat Bozkurt, Tuncay Tunç, Nagihan Çaylak Delibaş, Hacali Necefoğlu and Tuncer Hökelek**

### **S1. Comment**

As a part of our ongoing investigation on transition metal complexes of nicotinamide (NA), one form of niacin (Krishnamachari, 1974), and/or the nicotinic acid derivative *N,N*-diethylnicotinamide (DENA), an important respiratory stimulant (Bigoli *et al.*, 1972), the title compound was synthesized and its crystal structure is reported herein.

The structures of some DENA and/or NA complexes of the  $\text{Cu}^{2+}$ ,  $\text{Zn}^{2+}$  and  $\text{Co}^{2+}$  ions,  $[\text{Cu}_2(\text{C}_6\text{H}_5\text{COO})_4(\text{C}_{10}\text{H}_{14}\text{N}_2\text{O})_2]$  (Hökelek *et al.*, 1995);  $[\text{Cu}_2(\text{C}_8\text{H}_7\text{O}_2)_4(\text{C}_6\text{H}_6\text{N}_2\text{O})_2]$  (Necefoğlu *et al.*, 2010a);  $[\text{Zn}_2(\text{C}_{11}\text{H}_{14}\text{NO}_2)_4(\text{C}_{10}\text{H}_{14}\text{N}_2\text{O})_2]$  (Hökelek *et al.*, 2009a);  $[\text{Zn}_2(\text{C}_8\text{H}_8\text{NO}_2)_4(\text{C}_{10}\text{H}_{14}\text{N}_2\text{O})_2] \cdot 2\text{H}_2\text{O}$  (Hökelek *et al.*, 2009b);  $[\text{Zn}_2(\text{C}_9\text{H}_{10}\text{NO}_2)_4(\text{C}_{10}\text{H}_{14}\text{N}_2\text{O})_2]$  (Hökelek *et al.*, 2009c);  $[\text{Zn}_2(\text{C}_8\text{H}_7\text{O}_2)_4(\text{C}_{10}\text{H}_{14}\text{N}_2\text{O})_2]$  (Necefoğlu *et al.*, 2010b) and  $[\text{Co}_2(\text{C}_{11}\text{H}_{14}\text{NO}_2)_4(\text{C}_{10}\text{H}_{14}\text{N}_2\text{O})_2]$  (Hökelek *et al.*, 2011) have also been determined. In these structures, the benzoate ion acts as a bidentate ligand.

The asymmetric unit of the title  $\text{Cd}^{\text{II}}$  complex,  $[\text{Cd}(\text{CB})_2(\text{DENA})(\text{H}_2\text{O})]_n$ , contains two 3-chlorobenzoate (CB), one *N,N*-diethylnicotinamide (DENA) ligands and one coordinated water molecule; the CB ions act as bidentate ligands (Fig. 1). The coordination number of the  $\text{Cd}^{\text{II}}$  ion is six. Intramolecular  $\text{C}—\text{H} \cdots \text{O}$  hydrogen bonds (Fig. 1 and Table 2) link the DENA ligand to the carboxyl groups. The  $\text{O1}—\text{Cd1}—\text{O2}$  and  $\text{O3}—\text{Cd1}—\text{O4}$  angles are  $53.75(10)^\circ$  and  $54.23(9)^\circ$ , respectively. The corresponding  $\text{O}—\text{M}—\text{O}$  (where *M* is a metal) angles are  $53.50(14)^\circ$  in  $[\text{Cu}_2(\text{C}_8\text{H}_5\text{O}_3)_4(\text{C}_6\text{H}_6\text{N}_2\text{O})_4]$  (Sertçelik *et al.*, 2013),  $53.45(4)^\circ$  and  $51.97(4)^\circ$  in  $[\text{Cd}(\text{C}_7\text{H}_5\text{O}_3)_2(\text{C}_6\text{H}_6\text{NO})(\text{H}_2\text{O})_2] \cdot 2(\text{H}_2\text{O})$  (Çaylak Delibaş *et al.*, 2013),  $52.91(4)^\circ$  and  $53.96(4)^\circ$  in  $[\text{Cd}(\text{C}_8\text{H}_5\text{O}_3)_2(\text{C}_6\text{H}_6\text{N}_2\text{O})_2(\text{H}_2\text{O})] \cdot \text{H}_2\text{O}$  (Hökelek *et al.*, 2009d),  $60.70(4)^\circ$  in  $[\text{Co}(\text{C}_9\text{H}_{10}\text{NO}_2)_2(\text{C}_6\text{H}_6\text{N}_2\text{O})(\text{H}_2\text{O})_2]$  (Hökelek *et al.*, 2009e),  $58.45(9)^\circ$  in  $[\text{Mn}(\text{C}_9\text{H}_{10}\text{NO}_2)_2(\text{C}_6\text{H}_6\text{N}_2\text{O})(\text{H}_2\text{O})_2]$  (Hökelek *et al.*, 2009f),  $60.03(6)^\circ$  in  $[\text{Zn}(\text{C}_8\text{H}_8\text{NO}_2)_2(\text{C}_6\text{H}_6\text{N}_2\text{O})_2] \cdot \text{H}_2\text{O}$  (Hökelek *et al.*, 2009g),  $58.3(3)^\circ$  in  $[\text{Zn}_2(\text{C}_{10}\text{H}_{14}\text{N}_2\text{O})_2(\text{C}_7\text{H}_5\text{O}_3)_4] \cdot 2\text{H}_2\text{O}$  (Hökelek & Necefoğlu, 1996) and  $55.2(1)^\circ$  in  $[\text{Cu}(\text{Asp})_2(\text{py})_2]$  (where Asp is acetyl-salicylate and py is pyridine) (Greenaway *et al.*, 1984). The dihedral angles between the planar carboxylate groups  $[(\text{O1}/\text{O2}/\text{C1})$  and  $(\text{O3}/\text{O4}/\text{C8})]$  and the adjacent benzene rings A ( $\text{C2}—\text{C7}$ ) and B ( $\text{C9}—\text{C14}$ ) are  $4.56(28)^\circ$  and  $12.84(26)^\circ$ , respectively, while that between rings A, B and C ( $\text{N1}/\text{C15}—\text{C19}$ ) are  $\text{A}/\text{B} = 1.89(13)$ ,  $\text{A}/\text{C} = 34.83(13)$  and  $\text{B}/\text{C} = 35.13(11)^\circ$ .


In the crystal, the  $\text{Cd}^{\text{II}}$  ions  $[\text{Cd1} \cdots \text{Cd1a} = 7.0161(5) \text{ \AA}$ ; symmetry code: (a)  $x, y - 1, z$ ] are bridged by the N and O atoms of the DENA ligands forming polymeric chains running along the *b*-axis direction, where the coordination number of each  $\text{Cd}^{\text{II}}$  atom is seven within a  $\text{CdO}_6\text{N}$  donor set (Fig. 2). The average  $\text{Cd}—\text{O}$  distance is  $2.389(3) \text{ \AA}$  (Table 1). The Cd atom lies  $0.0972(3) \text{ \AA}$  below and  $0.0127(3) \text{ \AA}$  above of the carboxylate groups  $[(\text{O1}/\text{O2}/\text{C1})$  and  $(\text{O3}/\text{O4}/\text{C8})]$ , respectively. Strong intermolecular  $\text{O}—\text{H} \cdots \text{O}$  hydrogen bonds (Table 2) between water molecules and carboxylate groups link the adjacent chains into layers parallel to the *bc* plane.  $\pi \cdots \pi$  contacts between the benzene rings  $\text{Cg1}—\text{Cg2}^i$ , [symmetry code: (i)  $1 - x, 2 - y, -z$ , where Cg1 and Cg2 are the centroids of the rings A ( $\text{C2}—\text{C7}$ ) and B ( $\text{C9}—\text{C14}$ ), respectively] may further stabilize the structure, with centroid-centroid distance of  $3.912(2) \text{ \AA}$ .

## S2. Experimental

The title compound was prepared by the reaction of  $\text{CdSO}_4 \cdot 8\text{H}_2\text{O}$  (1.283 g, 5 mmol) in  $\text{H}_2\text{O}$  (100 ml) and diethylnicotinamide (1.780 g, 10 mmol) in  $\text{H}_2\text{O}$  (10 ml) with sodium 3-chlorobenzoate (1.790 g, 10 mmol) in  $\text{H}_2\text{O}$  (100 ml). The mixture was filtered and set aside to crystallize at ambient temperature for ten days, giving colorless single crystals.


## S3. Refinement

Atoms H61 and H62 (for  $\text{H}_2\text{O}$ ) were located in a difference Fourier map and were refined freely. The C-bound H-atoms were positioned geometrically with  $\text{C}-\text{H} = 0.93, 0.97$  and  $0.96 \text{ \AA}$ , for aromatic, methylene and methyl H-atoms, respectively, and constrained to ride on their parent atoms, with  $U_{\text{iso}}(\text{H}) = k \times U_{\text{eq}}(\text{C})$ , where  $k = 1.5$  for methyl H-atoms and  $k = 1.2$  for all other H-atoms.



**Figure 1**

The asymmetric unit of the title molecule with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. Hydrogen bonds are shown as dashed lines.


**Figure 2**

Part of the polymeric chain of the title compound [symmetry codes: (a)  $x, y - 1, z$ ; (b)  $x, 1 + y, z$ ]. Hydrogen atoms have been omitted for clarity.

**catena-Poly[[aquabis(3-chlorobenzoato- $\kappa^2O, O'$ )cadmium]- $\mu$ -*N,N*-diethylnicotinamide- $\kappa^2N^1:O$ ]**

*Crystal data*

$[\text{Cd}(\text{C}_7\text{H}_4\text{ClO}_2)_2(\text{C}_{10}\text{H}_{14}\text{N}_2\text{O})(\text{H}_2\text{O})]$

$M_r = 619.76$

Monoclinic,  $C2/c$

Hall symbol:  $-C2yc$

$a = 25.1809 (5) \text{ \AA}$

$b = 7.0161 (3) \text{ \AA}$

$c = 30.6755 (6) \text{ \AA}$

$\beta = 106.203 (3)^\circ$

$V = 5204.2 (3) \text{ \AA}^3$

$Z = 8$

$F(000) = 2496$

$D_x = 1.582 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 9713 reflections

$\theta = 2.5\text{--}28.4^\circ$

$\mu = 1.09 \text{ mm}^{-1}$

$T = 296 \text{ K}$

Rod-shaped, colourless

$0.35 \times 0.15 \times 0.10 \text{ mm}$

*Data collection*

Bruker SMART BREEZE CCD  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2012)

$T_{\min} = 0.823, T_{\max} = 0.897$

100329 measured reflections

6531 independent reflections

6170 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.026$

$\theta_{\max} = 28.4^\circ, \theta_{\min} = 1.4^\circ$

$h = -33 \rightarrow 33$

$k = -9 \rightarrow 9$

$l = -41 \rightarrow 40$

*Refinement*

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.046$

$wR(F^2) = 0.093$

$S = 1.35$

6531 reflections

326 parameters

4 restraints

Primary atom site location: structure-invariant direct methods  
 Secondary atom site location: difference Fourier map  
 Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.0097P)^2 + 22.0143P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 0.77 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -1.02 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional R-factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and R- factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cd1	0.430006 (10)	0.86384 (4)	0.032753 (9)	0.03649 (8)
Cl1	0.67865 (6)	0.6435 (2)	0.24132 (4)	0.0797 (4)
Cl2	0.20678 (5)	1.2006 (2)	-0.17853 (4)	0.0741 (4)
O1	0.50773 (11)	0.7328 (5)	0.09538 (10)	0.0534 (7)
O2	0.50591 (11)	1.0331 (5)	0.07578 (9)	0.0533 (7)
O3	0.35198 (12)	0.9354 (4)	-0.03179 (10)	0.0547 (8)
O4	0.41552 (11)	1.1524 (4)	-0.00840 (9)	0.0500 (7)
O5	0.39003 (12)	0.9782 (4)	0.09065 (10)	0.0498 (7)
O6	0.47410 (12)	0.7107 (5)	-0.01422 (11)	0.0485 (7)
H61	0.472 (2)	0.785 (7)	-0.0367 (12)	0.083 (19)*
H62	0.5081 (12)	0.709 (10)	0.0016 (18)	0.13 (3)*
N1	0.38036 (12)	0.6005 (4)	0.04343 (10)	0.0349 (6)
N2	0.42885 (14)	1.1630 (5)	0.15138 (11)	0.0453 (8)
C1	0.52769 (14)	0.8941 (6)	0.10063 (12)	0.0420 (9)
C2	0.57856 (13)	0.9332 (6)	0.13935 (12)	0.0368 (8)
C3	0.60309 (14)	0.7865 (6)	0.16792 (12)	0.0410 (8)
H3	0.5892	0.6631	0.1627	0.049*
C4	0.64831 (15)	0.8243 (6)	0.20430 (13)	0.0451 (9)
C5	0.66979 (17)	1.0049 (8)	0.21233 (15)	0.0575 (12)
H5	0.7004	1.0287	0.2369	0.069*
C6	0.6455 (2)	1.1492 (8)	0.18368 (16)	0.0648 (13)
H6	0.6597	1.2721	0.1888	0.078*
C7	0.59975 (17)	1.1136 (6)	0.14707 (15)	0.0516 (10)
H7	0.5835	1.2125	0.1278	0.062*
C8	0.37094 (14)	1.0967 (5)	-0.03489 (11)	0.0367 (8)
C9	0.33859 (14)	1.2296 (5)	-0.07074 (11)	0.0328 (7)
C10	0.29377 (14)	1.1625 (5)	-0.10465 (11)	0.0364 (7)
H10	0.2847	1.0337	-0.1060	0.044*
C11	0.26293 (16)	1.2868 (6)	-0.13623 (13)	0.0458 (9)

C12	0.2747 (2)	1.4770 (7)	-0.13478 (15)	0.0583 (12)
H12	0.2530	1.5599	-0.1561	0.070*
C13	0.3196 (2)	1.5445 (6)	-0.10090 (16)	0.0606 (12)
H13	0.3280	1.6737	-0.0994	0.073*
C14	0.35155 (18)	1.4214 (6)	-0.06956 (13)	0.0465 (9)
H14	0.3821	1.4673	-0.0474	0.056*
C15	0.40358 (14)	0.4560 (5)	0.07000 (12)	0.0363 (7)
H15	0.4418	0.4552	0.0822	0.044*
C16	0.37313 (15)	0.3070 (5)	0.08017 (12)	0.0346 (7)
C17	0.31657 (15)	0.3072 (6)	0.06121 (13)	0.0442 (9)
H17	0.2951	0.2065	0.0665	0.053*
C18	0.29234 (16)	0.4580 (7)	0.03445 (15)	0.0523 (10)
H18	0.2542	0.4629	0.0220	0.063*
C19	0.32558 (15)	0.6014 (6)	0.02650 (13)	0.0442 (9)
H19	0.3091	0.7037	0.0085	0.053*
C20	0.39878 (15)	1.1380 (5)	0.10824 (13)	0.0408 (8)
C21	0.43608 (18)	1.3440 (7)	0.17587 (14)	0.0530 (10)
H21A	0.4277	1.3260	0.2046	0.064*
H21B	0.4101	1.4362	0.1583	0.064*
C22	0.4944 (2)	1.4228 (8)	0.18486 (17)	0.0724 (14)
H22A	0.4972	1.5408	0.2012	0.109*
H22B	0.5026	1.4446	0.1565	0.109*
H22C	0.5203	1.3328	0.2026	0.109*
C23	0.4519 (2)	0.9923 (7)	0.17814 (16)	0.0606 (12)
H23A	0.4863	1.0261	0.2004	0.073*
H23B	0.4601	0.8969	0.1581	0.073*
C24	0.4129 (3)	0.9093 (9)	0.2021 (2)	0.097 (2)
H24A	0.4306	0.8052	0.2210	0.146*
H24B	0.3802	0.8641	0.1802	0.146*
H24C	0.4029	1.0056	0.2207	0.146*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cd1	0.03456 (13)	0.03444 (13)	0.03743 (13)	-0.00848 (11)	0.00502 (9)	0.00642 (11)
Cl1	0.0728 (8)	0.0926 (10)	0.0612 (7)	0.0290 (8)	-0.0022 (6)	0.0143 (7)
Cl2	0.0523 (6)	0.1018 (11)	0.0533 (6)	-0.0002 (7)	-0.0100 (5)	0.0020 (7)
O1	0.0415 (15)	0.0632 (19)	0.0483 (16)	-0.0173 (14)	0.0008 (12)	-0.0009 (14)
O2	0.0414 (15)	0.070 (2)	0.0435 (15)	-0.0062 (14)	0.0043 (12)	0.0154 (15)
O3	0.0528 (16)	0.0446 (16)	0.0570 (17)	-0.0091 (13)	-0.0005 (13)	0.0198 (14)
O4	0.0392 (14)	0.0561 (17)	0.0474 (15)	-0.0072 (13)	-0.0002 (11)	0.0094 (14)
O5	0.0609 (17)	0.0322 (14)	0.0653 (18)	-0.0061 (13)	0.0322 (15)	-0.0015 (13)
O6	0.0440 (16)	0.0526 (17)	0.0512 (17)	-0.0086 (13)	0.0173 (13)	0.0004 (14)
N1	0.0369 (14)	0.0292 (15)	0.0384 (15)	-0.0081 (12)	0.0103 (12)	0.0011 (12)
N2	0.0537 (19)	0.0402 (18)	0.0457 (18)	0.0070 (15)	0.0200 (15)	0.0106 (14)
C1	0.0291 (16)	0.063 (3)	0.0349 (18)	-0.0060 (17)	0.0107 (14)	0.0010 (18)
C2	0.0260 (15)	0.049 (2)	0.0356 (17)	-0.0057 (14)	0.0095 (13)	-0.0025 (15)
C3	0.0344 (17)	0.048 (2)	0.0401 (19)	-0.0016 (16)	0.0102 (15)	-0.0038 (17)

C4	0.0351 (18)	0.060 (3)	0.0383 (19)	0.0069 (17)	0.0074 (15)	-0.0016 (18)
C5	0.041 (2)	0.080 (3)	0.045 (2)	-0.013 (2)	0.0015 (18)	-0.014 (2)
C6	0.062 (3)	0.059 (3)	0.068 (3)	-0.023 (2)	0.008 (2)	-0.014 (2)
C7	0.049 (2)	0.046 (2)	0.058 (2)	-0.0066 (19)	0.0110 (19)	0.002 (2)
C8	0.0339 (17)	0.045 (2)	0.0324 (17)	-0.0001 (15)	0.0108 (13)	0.0058 (15)
C9	0.0377 (17)	0.0326 (17)	0.0300 (16)	-0.0011 (14)	0.0128 (13)	0.0032 (13)
C10	0.0373 (17)	0.0345 (18)	0.0376 (18)	-0.0006 (14)	0.0109 (14)	0.0006 (14)
C11	0.041 (2)	0.060 (3)	0.0341 (19)	0.0080 (18)	0.0072 (15)	0.0051 (18)
C12	0.070 (3)	0.053 (3)	0.048 (2)	0.018 (2)	0.011 (2)	0.018 (2)
C13	0.085 (3)	0.033 (2)	0.061 (3)	0.003 (2)	0.016 (2)	0.007 (2)
C14	0.059 (2)	0.037 (2)	0.040 (2)	-0.0086 (18)	0.0097 (17)	-0.0011 (16)
C15	0.0350 (17)	0.0350 (18)	0.0388 (18)	-0.0089 (14)	0.0102 (14)	0.0001 (15)
C16	0.0413 (18)	0.0294 (16)	0.0366 (17)	-0.0043 (14)	0.0168 (14)	-0.0004 (14)
C17	0.0406 (19)	0.041 (2)	0.054 (2)	-0.0147 (16)	0.0187 (17)	0.0038 (17)
C18	0.0332 (18)	0.060 (3)	0.061 (3)	-0.0097 (18)	0.0097 (17)	0.012 (2)
C19	0.0403 (19)	0.039 (2)	0.052 (2)	-0.0046 (16)	0.0100 (16)	0.0093 (17)
C20	0.0464 (19)	0.0345 (18)	0.049 (2)	-0.0065 (16)	0.0265 (17)	0.0047 (16)
C21	0.061 (3)	0.055 (3)	0.043 (2)	0.009 (2)	0.0142 (19)	0.0029 (19)
C22	0.070 (3)	0.079 (4)	0.062 (3)	-0.010 (3)	0.010 (2)	0.000 (3)
C23	0.069 (3)	0.056 (3)	0.065 (3)	0.022 (2)	0.032 (2)	0.024 (2)
C24	0.116 (5)	0.085 (4)	0.117 (5)	0.037 (4)	0.076 (4)	0.059 (4)

*Geometric parameters (Å, °)*

Cd1—O1	2.504 (3)	C9—C10	1.386 (5)
Cd1—O2	2.323 (3)	C9—C14	1.383 (5)
Cd1—O3	2.421 (3)	C10—C11	1.372 (5)
Cd1—O4	2.360 (3)	C10—H10	0.9300
Cd1—O5	2.410 (3)	C11—C12	1.365 (6)
Cd1—O6	2.314 (3)	C12—C13	1.388 (7)
Cd1—N1	2.305 (3)	C12—H12	0.9300
Cd1—C1	2.752 (4)	C13—H13	0.9300
Cd1—C8	2.732 (3)	C14—C13	1.373 (6)
C11—C4	1.732 (4)	C14—H14	0.9300
C12—C11	1.739 (4)	C15—C16	1.383 (5)
O1—C1	1.230 (5)	C15—H15	0.9300
O3—C8	1.242 (4)	C16—C17	1.380 (5)
O4—C8	1.251 (4)	C16—C20 <sup>i</sup>	1.502 (5)
O6—H61	0.856 (17)	C17—C18	1.373 (6)
O6—H62	0.86 (2)	C17—H17	0.9300
O5—C20	1.237 (5)	C18—H18	0.9300
N1—C15	1.330 (4)	C19—C18	1.374 (5)
N1—C19	1.331 (5)	C19—H19	0.9300
N2—C21	1.461 (5)	C20—N2	1.340 (5)
N2—C23	1.476 (5)	C20—C16 <sup>ii</sup>	1.502 (5)
C1—O2	1.265 (5)	C21—C22	1.521 (6)
C2—C1	1.509 (5)	C21—H21A	0.9700
C2—C3	1.381 (5)	C21—H21B	0.9700



C2—C7	1.368 (6)	C22—H22A	0.9600
C3—C4	1.380 (5)	C22—H22B	0.9600
C3—H3	0.9300	C22—H22C	0.9600
C4—C5	1.372 (6)	C23—C24	1.500 (6)
C5—C6	1.368 (7)	C23—H23A	0.9700
C5—H5	0.9300	C23—H23B	0.9700
C6—H6	0.9300	C24—H24A	0.9600
C7—C6	1.389 (6)	C24—H24B	0.9600
C7—H7	0.9300	C24—H24C	0.9600
C9—C8	1.497 (5)		
O1—Cd1—C1	26.54 (11)	C7—C6—H6	119.8
O1—Cd1—C8	161.03 (10)	C2—C7—C6	120.1 (4)
O2—Cd1—O1	53.75 (10)	C2—C7—H7	119.9
O2—Cd1—O3	135.22 (10)	C6—C7—H7	119.9
O2—Cd1—O4	81.10 (10)	O3—C8—Cd1	62.38 (19)
O2—Cd1—O5	81.83 (10)	O3—C8—O4	122.0 (3)
O2—Cd1—C1	27.21 (11)	O3—C8—C9	118.8 (3)
O2—Cd1—C8	108.24 (11)	O4—C8—Cd1	59.57 (19)
O3—Cd1—O1	170.39 (10)	O4—C8—C9	119.2 (3)
O3—Cd1—C1	162.33 (11)	C9—C8—Cd1	178.0 (3)
O3—Cd1—C8	27.04 (10)	C10—C9—C8	120.1 (3)
O4—Cd1—O1	134.15 (9)	C14—C9—C8	120.7 (3)
O4—Cd1—O3	54.23 (9)	C14—C9—C10	119.1 (3)
O4—Cd1—O5	94.28 (10)	C9—C10—H10	120.1
O4—Cd1—C1	108.10 (11)	C11—C10—C9	119.7 (4)
O4—Cd1—C8	27.19 (10)	C11—C10—H10	120.1
O5—Cd1—O1	87.40 (10)	C10—C11—C12	119.2 (3)
O5—Cd1—O3	97.07 (11)	C12—C11—C10	121.6 (4)
O5—Cd1—C1	83.43 (10)	C12—C11—C12	119.2 (3)
O5—Cd1—C8	96.30 (10)	C11—C12—C13	118.8 (4)
O6—Cd1—O1	84.21 (11)	C11—C12—H12	120.6
O6—Cd1—O2	97.43 (11)	C13—C12—H12	120.6
O6—Cd1—O3	90.46 (11)	C12—C13—H13	119.8
O6—Cd1—O4	95.42 (11)	C14—C13—C12	120.4 (4)
O6—Cd1—O5	170.03 (11)	C14—C13—H13	119.8
O6—Cd1—C1	91.39 (11)	C9—C14—H14	119.8
O6—Cd1—C8	93.37 (11)	C13—C14—C9	120.4 (4)
N1—Cd1—O1	86.30 (10)	C13—C14—H14	119.8
N1—Cd1—O2	136.21 (10)	N1—C15—C16	122.6 (3)
N1—Cd1—O3	86.21 (10)	N1—C15—H15	118.7
N1—Cd1—O4	139.00 (10)	C16—C15—H15	118.7
N1—Cd1—O5	78.90 (10)	C15—C16—C20 <sup>i</sup>	123.4 (3)
N1—Cd1—O6	95.16 (10)	C17—C16—C15	118.4 (3)
N1—Cd1—C1	111.11 (11)	C17—C16—C20 <sup>i</sup>	118.1 (3)
N1—Cd1—C8	112.67 (10)	C16—C17—H17	120.4
C8—Cd1—C1	135.29 (12)	C18—C17—C16	119.1 (3)
C1—O1—Cd1	88.1 (2)	C18—C17—H17	120.4

C1—O2—Cd1	95.6 (2)	C17—C18—C19	118.7 (4)
C8—O3—Cd1	90.6 (2)	C17—C18—H18	120.7
C8—O4—Cd1	93.2 (2)	C19—C18—H18	120.7
C20—O5—Cd1	124.3 (2)	N1—C19—C18	122.9 (4)
Cd1—O6—H61	107 (4)	N1—C19—H19	118.5
Cd1—O6—H62	103 (5)	C18—C19—H19	118.5
H61—O6—H62	107 (4)	O5—C20—N2	122.2 (4)
C15—N1—Cd1	122.3 (2)	O5—C20—C16 <sup>ii</sup>	118.0 (3)
C15—N1—C19	118.2 (3)	N2—C20—C16 <sup>ii</sup>	119.8 (3)
C19—N1—Cd1	119.0 (2)	N2—C21—C22	112.6 (4)
C20—N2—C21	125.2 (3)	N2—C21—H21A	109.1
C20—N2—C23	118.0 (4)	N2—C21—H21B	109.1
C21—N2—C23	116.5 (3)	C22—C21—H21A	109.1
O1—C1—Cd1	65.4 (2)	C22—C21—H21B	109.1
O1—C1—O2	122.5 (3)	H21A—C21—H21B	107.8
O1—C1—C2	119.7 (4)	C21—C22—H22A	109.5
O2—C1—Cd1	57.16 (19)	C21—C22—H22B	109.5
O2—C1—C2	117.7 (4)	C21—C22—H22C	109.5
C2—C1—Cd1	173.1 (3)	H22A—C22—H22B	109.5
C3—C2—C1	119.7 (3)	H22A—C22—H22C	109.5
C7—C2—C1	120.6 (4)	H22B—C22—H22C	109.5
C7—C2—C3	119.7 (3)	N2—C23—C24	112.3 (4)
C2—C3—H3	120.3	N2—C23—H23A	109.2
C4—C3—C2	119.5 (4)	N2—C23—H23B	109.2
C4—C3—H3	120.3	C24—C23—H23A	109.2
C3—C4—Cl1	120.2 (3)	C24—C23—H23B	109.2
C5—C4—Cl1	118.6 (3)	H23A—C23—H23B	107.9
C5—C4—C3	121.2 (4)	C23—C24—H24A	109.5
C4—C5—H5	120.5	C23—C24—H24B	109.5
C6—C5—C4	119.0 (4)	C23—C24—H24C	109.5
C6—C5—H5	120.5	H24A—C24—H24B	109.5
C5—C6—C7	120.5 (4)	H24A—C24—H24C	109.5
C5—C6—H6	119.8	H24B—C24—H24C	109.5
O2—Cd1—O1—C1	1.3 (2)	Cd1—O5—C20—N2	-109.6 (3)
O4—Cd1—O1—C1	13.0 (3)	Cd1—O5—C20—C16 <sup>ii</sup>	72.2 (4)
O5—Cd1—O1—C1	-80.5 (2)	C15—N1—Cd1—O1	-9.4 (3)
O6—Cd1—O1—C1	104.9 (2)	C15—N1—Cd1—O2	-31.9 (3)
N1—Cd1—O1—C1	-159.5 (2)	C15—N1—Cd1—O3	164.6 (3)
C8—Cd1—O1—C1	21.4 (5)	C15—N1—Cd1—O4	178.8 (2)
O1—Cd1—O2—C1	-1.3 (2)	C15—N1—Cd1—O5	-97.5 (3)
O3—Cd1—O2—C1	-176.7 (2)	C15—N1—Cd1—O6	74.4 (3)
O4—Cd1—O2—C1	-172.8 (2)	C15—N1—Cd1—C1	-19.0 (3)
O5—Cd1—O2—C1	91.5 (2)	C15—N1—Cd1—C8	170.3 (3)
O6—Cd1—O2—C1	-78.4 (2)	C19—N1—Cd1—O1	163.1 (3)
N1—Cd1—O2—C1	27.0 (3)	C19—N1—Cd1—O2	140.6 (3)
C8—Cd1—O2—C1	-174.5 (2)	C19—N1—Cd1—O3	-22.9 (3)
O2—Cd1—O3—C8	4.5 (3)	C19—N1—Cd1—O4	-8.6 (4)

O4—Cd1—O3—C8	-0.2 (2)	C19—N1—Cd1—O5	75.1 (3)
O5—Cd1—O3—C8	90.0 (2)	C19—N1—Cd1—O6	-113.0 (3)
O6—Cd1—O3—C8	-96.5 (2)	C19—N1—Cd1—C1	153.5 (3)
N1—Cd1—O3—C8	168.3 (2)	C19—N1—Cd1—C8	-17.2 (3)
C1—Cd1—O3—C8	-0.5 (5)	Cd1—N1—C15—C16	173.3 (3)
O1—Cd1—O4—C8	174.0 (2)	C19—N1—C15—C16	0.7 (5)
O2—Cd1—O4—C8	-176.5 (2)	Cd1—N1—C19—C18	-174.4 (3)
O3—Cd1—O4—C8	0.2 (2)	C15—N1—C19—C18	-1.5 (6)
O5—Cd1—O4—C8	-95.5 (2)	C20—N2—C21—C22	-109.8 (5)
O6—Cd1—O4—C8	86.8 (2)	C23—N2—C21—C22	76.9 (5)
N1—Cd1—O4—C8	-17.5 (3)	C20—N2—C23—C24	-89.3 (5)
C1—Cd1—O4—C8	-179.9 (2)	C21—N2—C23—C24	84.5 (6)
O1—Cd1—C1—O2	177.7 (4)	O1—C1—O2—Cd1	2.4 (4)
O2—Cd1—C1—O1	-177.7 (4)	C2—C1—O2—Cd1	-174.6 (3)
O3—Cd1—C1—O1	-170.0 (3)	C3—C2—C1—O1	1.6 (5)
O3—Cd1—C1—O2	7.8 (5)	C3—C2—C1—O2	178.7 (3)
O4—Cd1—C1—O1	-170.3 (2)	C7—C2—C1—O1	-176.8 (4)
O4—Cd1—C1—O2	7.5 (2)	C7—C2—C1—O2	0.4 (5)
O5—Cd1—C1—O1	97.4 (2)	C1—C2—C3—C4	-177.5 (3)
O5—Cd1—C1—O2	-84.9 (2)	C7—C2—C3—C4	0.9 (6)
O6—Cd1—C1—O1	-74.1 (2)	C1—C2—C7—C6	177.9 (4)
O6—Cd1—C1—O2	103.6 (2)	C3—C2—C7—C6	-0.5 (6)
N1—Cd1—C1—O1	22.0 (3)	C2—C3—C4—C11	178.4 (3)
N1—Cd1—C1—O2	-160.3 (2)	C2—C3—C4—C5	-0.8 (6)
C8—Cd1—C1—O1	-170.3 (2)	C11—C4—C5—C6	-178.9 (4)
C8—Cd1—C1—O2	7.4 (3)	C3—C4—C5—C6	0.3 (7)
O1—Cd1—C8—O3	166.4 (3)	C4—C5—C6—C7	0.1 (7)
O1—Cd1—C8—O4	-13.3 (5)	C2—C7—C6—C5	0.0 (7)
O2—Cd1—C8—O3	-176.6 (2)	C10—C9—C8—O3	-11.4 (5)
O2—Cd1—C8—O4	3.7 (2)	C10—C9—C8—O4	170.1 (3)
O3—Cd1—C8—O4	-179.7 (4)	C14—C9—C8—O3	166.1 (4)
O4—Cd1—C8—O3	179.7 (4)	C14—C9—C8—O4	-12.4 (5)
O5—Cd1—C8—O3	-93.2 (2)	C8—C9—C10—C11	177.2 (3)
O5—Cd1—C8—O4	87.1 (2)	C14—C9—C10—C11	-0.3 (5)
O6—Cd1—C8—O3	84.4 (2)	C8—C9—C14—C13	-175.9 (4)
O6—Cd1—C8—O4	-95.3 (2)	C10—C9—C14—C13	1.6 (6)
N1—Cd1—C8—O3	-12.6 (3)	C9—C10—C11—C12	-179.9 (3)
N1—Cd1—C8—O4	167.7 (2)	C9—C10—C11—C12	-1.1 (6)
C1—Cd1—C8—O3	179.8 (2)	C10—C11—C12—C13	1.1 (7)
C1—Cd1—C8—O4	0.1 (3)	C12—C11—C12—C13	-180.0 (4)
Cd1—O1—C1—O2	-2.3 (4)	C11—C12—C13—C14	0.2 (7)
Cd1—O1—C1—C2	174.8 (3)	C9—C14—C13—C12	-1.6 (7)
Cd1—O3—C8—O4	0.3 (4)	N1—C15—C16—C17	1.3 (5)
Cd1—O3—C8—C9	-178.2 (3)	N1—C15—C16—C20 <sup>i</sup>	176.9 (3)
Cd1—O4—C8—O3	-0.3 (4)	C15—C16—C17—C18	-2.4 (6)
Cd1—O4—C8—C9	178.1 (3)	C20 <sup>i</sup> —C16—C17—C18	-178.4 (4)
C20—O5—Cd1—O1	91.2 (3)	C16—C17—C18—C19	1.7 (6)
C20—O5—Cd1—O2	37.5 (3)	N1—C19—C18—C17	0.3 (7)

C20—O5—Cd1—O3	-97.3 (3)	O5—C20—N2—C21	-173.2 (4)
C20—O5—Cd1—O4	-42.9 (3)	O5—C20—N2—C23	0.0 (5)
C20—O5—Cd1—N1	178.0 (3)	C16 <sup>ii</sup> —C20—N2—C21	5.0 (5)
C20—O5—Cd1—C1	64.9 (3)	C16 <sup>ii</sup> —C20—N2—C23	178.2 (3)
C20—O5—Cd1—C8	-70.1 (3)		

Symmetry codes: (i)  $x, y-1, z$ ; (ii)  $x, y+1, z$ .

*Hydrogen-bond geometry (Å, °)*

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
O6—H61...O2 <sup>iii</sup>	0.85 (4)	1.94 (4)	2.753 (5)	160 (4)
O6—H62...O4 <sup>iii</sup>	0.86 (4)	2.11 (4)	2.838 (4)	142 (5)
C15—H15...O1	0.93	2.52	3.181 (5)	128
C19—H19...O3	0.93	2.47	3.130 (5)	128

Symmetry code: (iii)  $-x+1, -y+2, -z$ .