

Crystal structure of 1,4:5,8-dimethano-1,1a,4,4a,5,5a,8,8a-octahydroanthracene-9,10-dione, C₁₆H₁₆O₂

B. Güneş, Soylu

Gazi University, Gazi Education Faculty, Physics Department, 06500 Beşevler Ankara, Turkey


S. Özbey

Hacettepe University, Engineering Faculty, Physics Engineering Department, 06532 Beytepe Ankara, Turkey

and A. Aydın

Gazi University, Gazi Education Faculty, Chemistry Department, 06500 Beşevler Ankara, Turkey

Received March 13, 1998, CSD-No. 409341



Abstract

C₁₆H₁₆O₂, monoclinic, *P*12₁/*n*1 (No. 14), *a* = 12.0005(7) Å, *b* = 6.2388(9) Å, *c* = 16.973(2) Å, β = 110.609(6)°, *V* = 1189.5 Å³, *Z* = 4, *R*_g(*F*) = 0.047, *R*_w(*F*²) = 0.108, *T* = 293 K.

Source of material

The title compound is a Diels-Alder adduct of a [4+2] cycloaddition of 1,3-cyclopentadiene with *p*-benzoquinone. Cyclopentadiene freshly distilled (40.32 g, 0.6 mol) in methanol (20 ml) was added to a solution of *p*-benzoquinone (32.43 g, 0.3 mol) in methanol (100 ml) at 203 K. The resultant colorless solid was recrystallized from ethanol.

Discussion

The *p*-benzoquinone (PBQ) molecule used to obtain the title compound was found to be planar at room temperature [1], then, at 113 K, the significant deviations from planarity were observed, and the PBQ molecule was in the chair conformation [2]. In contrast to these results, it has been found that the 1,4-cyclohexadione, which is a chemically changed form of PBQ molecule, is not planar and has a boat conformation. The near zero torsion angles of this boat are −0.6(3)° and 0.7(3)°. The dihedral angles between 1,4-cyclohexadione ring and the planes formed by norbornenyl atoms of C₂,C₃,C₇,C₁₀ and C₅,C₆,C₁₁,C₁₄ are 70.68(9)° and 39.08(7)°, respectively. In the 1,4-cyclohexadione ring, the bond length of O=C is 1.215(2) Å, the C–C distance in

the range 1.499(3) Å – 1.562(3) Å. O=C–C, C(=O)–C–C and C–C(=O)–C angles in the range 120.0(2)° – 120.9(2)°, 115.8(2)° – 116.6(2)° and 119.1(2)°, respectively. For the norbornenyl atoms, the C–C and C=C distances in the range 1.501(3) Å – 1.579(3) Å and 1.317(3) Å – 1.318(3) Å, respectively. The bond lengths and angles are comparable with corresponding values observed in related molecules [3, 4]. The molecules are linked together by weak C–H⋯O hydrogen bonds. The bond lengths and angles are comparable with corresponding values observed in related molecules [3, 4].

Table 1. Data collection and handling.

Crystal:	colorless, prismatic, size 0.20 x 0.36 x 0.64 mm
Wavelength:	Mo K _α radiation (0.71073 Å)
μ:	0.87 cm ^{−1}
Diffractometer, scan mode:	Enraf-Nonius CAD4, ω/2θ
2θ _{max} :	52.58°
<i>N</i> (<i>hkl</i>) _{measured} , <i>N</i> (<i>hkl</i>) _{unique} :	2424, 2293
Criterion for <i>I</i> _{obs} , <i>N</i> (<i>hkl</i>) _{gt} :	<i>I</i> _{obs} > 2 σ(<i>I</i> _{obs}), 1774
<i>N</i> (<i>param</i>) _{refined} :	228
Programs:	MolEN [5], SHELXS-86 [6], SHELXL-93 [7], ORTEP [8]

Table 2. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso}
H(2)	4e	1.125(1)	0.596(4)	0.448(1)	0.043(6)
H(3)	4e	1.223(1)	0.611(4)	0.362(1)	0.044(6)
H(5)	4e	0.987(1)	0.280(3)	0.203(1)	0.036(5)
H(6)	4e	0.887(1)	0.267(3)	0.291(1)	0.033(5)
H(7)	4e	1.203(2)	0.346(4)	0.559(1)	0.051(6)
H(8)	4e	1.193(2)	−0.014(5)	0.493(2)	0.069(8)
H(9)	4e	1.307(2)	0.001(5)	0.393(2)	0.075(8)
H(10)	4e	1.388(2)	0.377(4)	0.401(1)	0.049(6)
H(11)	4e	0.986(2)	0.614(4)	0.144(1)	0.050(6)
H(12)	4e	1.075(2)	0.868(4)	0.263(1)	0.056(7)
H(13)	4e	0.963(2)	0.860(4)	0.360(1)	0.060(7)
H(14)	4e	0.800(2)	0.590(3)	0.304(1)	0.045(6)
H(15A)	4e	1.411(2)	0.359(4)	0.553(2)	0.066(8)
H(15B)	4e	1.347(2)	0.587(5)	0.516(1)	0.064(8)
H(16A)	4e	0.791(2)	0.728(4)	0.159(1)	0.063(7)
H(16B)	4e	0.793(2)	0.468(4)	0.157(1)	0.061(7)

Table 3. Atomic coordinates and displacement parameters (in Å²)

Atom	Site	x	y	z	U ₁₁	U ₂₂	U ₃₃	U ₁₂	U ₁₃	U ₂₃
O(1)	4e	0.9709(1)	0.2632(3)	0.4431(1)	0.063(1)	0.056(1)	0.0544(9)	-0.0062(9)	0.0353(8)	0.0083(8)
O(2)	4e	1.2023(1)	0.2869(3)	0.24604(9)	0.060(1)	0.063(1)	0.0520(9)	0.0151(9)	0.0329(8)	-0.0043(8)
C(1)	4e	1.0134(2)	0.3560(3)	0.3974(1)	0.046(1)	0.027(1)	0.043(1)	0.0023(9)	0.025(1)	-0.0012(9)
C(2)	4e	1.1358(2)	0.4531(3)	0.4320(1)	0.048(1)	0.031(1)	0.037(1)	-0.0003(9)	0.0203(9)	-0.0042(9)
C(3)	4e	1.2075(2)	0.4633(3)	0.3712(1)	0.039(1)	0.031(1)	0.044(1)	-0.0013(9)	0.0198(9)	0.0004(9)
C(4)	4e	1.1457(2)	0.3720(3)	0.2843(1)	0.048(1)	0.029(1)	0.039(1)	0.0044(9)	0.0245(9)	0.0034(9)
C(5)	4e	1.0130(2)	0.3913(3)	0.2448(1)	0.047(1)	0.032(1)	0.034(1)	0.0002(9)	0.0177(9)	-0.0059(9)
C(6)	4e	0.9418(2)	0.3813(3)	0.3053(1)	0.039(1)	0.031(1)	0.043(1)	-0.0075(9)	0.0181(9)	-0.0046(9)
C(7)	4e	1.2233(2)	0.3314(4)	0.5092(1)	0.060(1)	0.053(1)	0.033(1)	0.002(1)	0.016(1)	-0.002(1)
C(8)	4e	1.2339(2)	0.1066(4)	0.4797(1)	0.063(2)	0.043(1)	0.047(1)	0.005(1)	0.011(1)	0.009(1)
C(9)	4e	1.2946(2)	0.1166(4)	0.4286(2)	0.060(1)	0.047(1)	0.049(1)	0.014(1)	0.013(1)	0.002(1)
C(10)	4e	1.3258(2)	0.3484(4)	0.4222(1)	0.041(1)	0.056(1)	0.052(1)	0.002(1)	0.018(1)	0.000(1)
C(11)	4e	0.9690(2)	0.6087(4)	0.1963(1)	0.052(1)	0.048(1)	0.044(1)	0.009(1)	0.024(1)	0.009(1)
C(12)	4e	1.0114(2)	0.7886(4)	0.2591(2)	0.050(1)	0.030(1)	0.076(2)	0.004(1)	0.021(1)	0.012(1)
C(13)	4e	0.9504(2)	0.7805(4)	0.3102(2)	0.055(1)	0.030(1)	0.060(1)	0.010(1)	0.018(1)	-0.002(1)
C(14)	4e	0.8661(2)	0.5943(4)	0.2832(1)	0.039(1)	0.044(1)	0.049(1)	0.006(1)	0.021(1)	0.002(1)
C(15)	4e	1.3421(2)	0.4256(5)	0.5113(2)	0.053(1)	0.062(2)	0.051(1)	-0.004(1)	0.005(1)	-0.009(1)
C(16)	4e	0.8377(2)	0.5996(5)	0.1879(1)	0.046(1)	0.055(2)	0.046(1)	0.007(1)	0.014(1)	0.004(1)

Acknowledgments. This research was partly supported by Gazi University, Research Fund Department. Research No: GEF 04/97-7.

References

1. Trotter, J.: A three-dimensional analysis of the crystal structure of p-benzoquinone. Acta Crystallogr. **13** (1960) 86-95.
2. Bolhuis, F. van; Kiers, C. Th.: Refinement of the crystal structure of p-Benzoquinone at -160° C. Acta Crystallogr. **B34** (1978) 1015-1016.
3. Garcia, J. G.; Fronczek, F. R.; McLaughlin, M. L.: *endo-endo* Diels-Alder diadduct of 5,5-dimethoxy-1,2,3,4-tetrachloro-1,3-cyclopentadiene with 1,5-cyclooctadiene, (1), and the dechlorinated Diels-Alder diadduct, (2). Acta Crystallogr. **C47** (1991) 206-209.
4. Garcia, J. G.; Morales, G.; Fronczek, F. R.; McLaughlin, M. L.: The structures of *endo*-13-*syn*-methyltricyclo(8.2.1.0^{2,9})trideca-5,11-dien-13-antiol, *endo*-13-*syn*-methyltricyclo(8.2.1.0^{2,9})trideca-5,11-dien-13-antiyl Acetate and its corresponding *endo-endo-anti* Diels-Alder adducts with 1,2,3,4-tetrachloro-5,5-dimethoxy-1,3-cyclopentadiene and hexachloro-rocyclopentadiene. Acta Crystallogr. **C50** (1994) 317-324.
5. Fair, C. K.: MolEN. An Interactive Intelligent System for Crystal Structure Analysis. Enraf-Nonius, Delft, The Netherlands 1990.
6. Sheldrick, G. M.: SHELXS-86. Program for the Solution of Crystal Structures. University of Göttingen, Germany 1986.
7. Sheldrick, G. M.: SHELXL-93. Program for the Refinement of Crystal Structures. University of Göttingen, Germany 1993.
8. Johnson, C. K.: ORTEPII. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA 1976.