

Turkey

Demographic and Health Survey 1998



Hacettepe University
Institute of Population Studies



MEASURE DHS+
Macro International Inc.

World Summit for Children Indicators: Turkey 1998

		Value
BASIC INDICATORS		
Childhood mortality	Infant mortality rate	42.7 per 1,000
	Under-five mortality rate	52.1 per 1,000
Childhood undernutrition	Percent stunted	16.0
	Percent wasted	1.9
	Percent underweight	8.3
Clean water supply	Percent of households within 15 minutes of a safe water supply ¹	74.4
Sanitary excreta disposal	Percent of households with flush toilets or VIP latrines	87.7
Basic education	Percent of women 15-49 with completed primary education	78.3
	Percent of men 15-49 with completed primary education	94.5
	Percent of girls 6-12 attending school	69.7
	Percent of boys 6-12 attending school	78.7
	Percent of women 15-49 who are literate	84.5
Children in especially difficult situations	Percent of children who are orphans (both parents dead)	0.1
	Percent of children who do not live with their natural mother	3.5
	Percent of children who live in single adult households	2.1
SUPPORTING INDICATORS		
Women's Health		
Birth spacing	Percent of births within 24 months of a previous birth	26.2
Safe motherhood	Percent of births with medical antenatal care	67.5
	Percent of births with antenatal care in first trimester	46.4
	Percent of births with medical assistance at delivery	80.6
	Percent of births in a medical facility	72.5
	Percent of births at high risk	40.2
Family planning	Contraceptive prevalence rate (any method, currently married women)	63.9
	Percent of currently married women with an unmet demand for family planning	10.1
	Percent of currently married women with an unmet need for family planning to avoid a high-risk birth	7.7
Nutrition		
Maternal nutrition	Percent of mothers with low BMI	2.5
Low birth weight	Percent of births at low birth weight (of those reporting numeric weight)	14.4
Breastfeeding	Percent of children under 4 months who are exclusively breastfed	9.2
Child Health		
Vaccinations	Percent of children whose mothers received tetanus toxoid vaccination during pregnancy	43.8
	Percent of children 12-23 months with measles vaccination	78.5
	Percent of children 12-23 months fully vaccinated	45.7
Diarrhoea control	Percent of children with diarrhoea in preceding 2 weeks who received oral rehydration therapy (sugar-salt-water solution)	26.9

¹ Piped, well, and bottled water

Turkish Demographic and Health Survey 1998

**Hacettepe University, Institute of Population Studies
Ankara, Turkey**

**Macro International Inc.
Calverton, Maryland, USA**

with the contributions of

**General Directorate of
Mother and Child
Health/Family Planning,
Ministry of Health, Ankara,
Turkey**

**United Nations
Population Fund,
New York,
USA**

**U.S. Agency for
International Development,
Washington DC,
USA**

October 1999

This report summarises the findings of the 1998 Turkish Demographic and Health Survey (TDHS-98) conducted by the Institute of Population Studies, Hacettepe University (HIPS) in collaboration with the General Directorate of Mother and Child Health/Family Planning, Ministry of Health. Technical and financial support for the survey were provided both by the United Nations Population Fund (UNFPA) and Macro International Inc. through its MEASURE/DHS+ project, a project sponsored primarily by the United States Agency for International Development (USAID) to carry out population and health surveys in developing countries.

The TDHS-98 is part of the worldwide Demographic and Health Surveys (MEASURE/DHS+) program, which is designed to provide decision-makers in survey countries with a database and analyses useful for informed policy choices, to expand the international population and health database, to advance survey methodology, and to develop in participating countries the skills and resources necessary to conduct high-quality demographic and health surveys. The TDHS-98 survey is the most recent in a series of demographic surveys carried out in Turkey by HIPS to provide information on fertility and child mortality levels; family planning awareness, approval and use; and basic indicators of maternal and child health.

Additional information on the TDHS-98 can be obtained from Hacettepe University, Institute of Population Studies, 06100 Ankara, Turkey (Telephone: 312-3107906; Fax: 312-3118141; E-mail: hps@hacettepe.edu.tr). Information on the worldwide MEASURE/DHS+ program may be obtained by writing: MEASURE/DHS+, Macro International Inc., 11785 Beltsville Drive, Suite 300, Calverton, MD 20705, USA (Telephone: 301-572-0200; Fax: 301-572-0999).

CONTENTS

	Page
Tables.....	vii
Figures.....	xi
Foreword.....	xiii
Summary of Findings.....	xv
Map of Turkey.....	xviii
CHAPTER 1 INTRODUCTION	
Sunday Üner and Banu Akadlı Ergöçmen.....	1
1.1 Geography.....	1
1.2 History.....	1
1.3 Administrative Divisions and Political Organisation.....	2
1.4 Social and Cultural Features.....	2
1.5 Economy.....	3
1.6 Regional Breakdown.....	4
1.7 Population.....	5
1.8 Population and Family Planning Policies and Programs.....	6
1.9 Health Priorities and Programs.....	6
1.10 Health Care System in Turkey.....	7
1.11 Objectives and Organisation of the Survey.....	7
CHAPTER 2 CHARACTERISTICS OF HOUSEHOLDS AND RESPONDENTS	
İsmet Koç and Attila Hancıoğlu.....	13
2.1 Population by Age and Sex.....	7
2.2 Household Composition.....	16
2.3 Fosterhood and Orphanhood.....	16
2.4 Educational Level of the Household Population.....	17
2.5 School Enrollment.....	19
2.6 Housing Characteristics.....	19
2.7 Household Durable Goods.....	22
2.8 Background Characteristics.....	23
2.9 Respondents' Level of Education by Background Characteristics.....	24
2.10 Reasons for Leaving School.....	25
2.11 Differentials in Characteristics of Couples.....	27
2.12 Exposure to Print Media.....	28
2.13 Employment and Occupation.....	29
2.14 Decision on Use of Earnings.....	31
2.15 Child Care While Working.....	31

CHAPTER 3 FERTILITY		
	Aykut Toros	35
3.1	Current Fertility.....	35
3.2	Children Ever Born and Living.....	40
3.3	Birth Intervals.....	41
3.4	Age at First Birth.....	42
3.5	Teenage Pregnancy and Motherhood	43
CHAPTER 4 FERTILITY REGULATION		
	Turgay Ünalán and İsmet Koç.....	45
4.1	Knowledge of Family Planning Methods.....	45
4.2	Ever Use of Family Planning Methods.....	46
4.3	Current Use of Contraceptive Method.....	47
4.4	Trend in Contraceptive Use.....	51
4.5	Number of Children at First Use of Contraception.....	52
4.6	Knowledge of the Fertile Period.....	53
4.7	Timing of Female Sterilisation	55
4.8	Sources for Family Planning Methods	55
4.9	Discontinuation of Contraceptive Use.....	57
4.10	Future Use of Family Planning.....	59
4.11	Exposure to Family Planning Messages in the Electronic Media.....	61
4.12	Acceptability of Use of Electronic Media to Disseminate Family Planning Messages.....	63
4.13	Exposure to Family Planning Messages in the Print Media	63
4.14	Attitudes of Couples Toward Family Planning	65
CHAPTER 5 ABORTIONS AND STILLBIRTHS		
	Banu Akadhı Ergöçmen and Turgay Ünalán	69
5.1	Life-time Experience of Women	69
5.2	Current Levels and Trends	71
5.3	Contraceptive Use Before and After Induced Abortions.....	72
5.4	Reasons for Induced Abortion.....	73
5.5	Timing of Induced Abortions	74
5.6	Provider	75
CHAPTER 6 OTHER PROXIMATE DETERMINANTS OF FERTILITY		
	Banu Akadhı Ergöçmen and İsmet Koç.....	77
6.1	Current Marital Status	77
6.2	Age at First Marriage	78
6.3	Age At First Sexual Intercourse	81
6.4	Postpartum Amenorrhoea, Postpartum Abstinence, and Insusceptibility	82
6.5	Termination of Exposure to Pregnancy.....	85

	Page
CHAPTER 7 FERTILITY PREFERENCES	
Turgay Ünalın and Elif Kurtuluş	87
7.1 Desire for More Children	87
7.2 Need for Family Planning Services	91
7.3 Ideal and Actual Number of Children	92
7.4 Fertility Planning.....	95
CHAPTER 8 INFANT AND CHILD MORTALITY	
Attila Hancıođlu	97
8.1 Definitions of Infant and Child Mortality.....	97
8.2 Assessment of Data Quality	97
8.3 Levels and Trends in Infant and Child Mortality	98
8.4 Differentials in Infant and Child Mortality.....	100
8.5 High-risk Fertility Behaviour	102
CHAPTER 9 MATERNAL AND CHILD HEALTH	
Banu Akadh Ergöçmen	105
9.1 Antenatal Care and Delivery Assistance	105
9.2 Immunisation of Children	114
9.3 Prevalence and Treatment of Diarrhoea.....	117
CHAPTER 10 INFANT FEEDING, MATERNAL AND CHILDHOOD NUTRITION	
Ergül Tunçbilek, Elif Kurtuluş, and Attila Hancıođlu	123
10.1 Breastfeeding and Supplementation.....	123
10.2 Nutritional Status	128
10.3 Maternal Nutrition.....	132
CHAPTER 11 KNOWLEDGE OF AIDS AND SEXUALLY TRANSMITTED DISEASES	
Attila Hancıođlu and A. Sinan Türkyılmaz.....	135
11.1 Source of Information about AIDS	135
11.2 Knowledge of Ways to Prevent AIDS.....	138
11.3 Perception of Risk of AIDS.....	141
REFERENCES.....	145
APPENDIX A PERSONNEL INVOLVED IN THE TURKISH DEMOGRAPHIC AND HEALTH SURVEY	147

APPENDIX B SURVEY DESIGN	
A. Sinan Türkyılmaz and Alfredo Aliaga	151
B.1 Sample Design and Implementation.....	153
B.2 Sample Frame.....	154
B.3 Stratification.....	154
B.4 Sample Allocation.....	155
B.5 Sample Selection.....	156
B.6 Questionnaire Development and Pre-Test.....	157
B.7 Data Collection Activities	160
B.8 Data Processing and Analysis	161
B.9 Calculation of Sample Weights	161
APPENDIX C ESTIMATES OF SAMPLING ERRORS	
Alfredo Aliaga and A. SinanTurkyılmaz	169
APPENDIX D DATA QUALITY TABLES	183
APPENDIX E QUESTIONNAIRES	191

TABLES

	Page
Table 1.1	Results of the household and individual interviews..... 11
Table 2.1	Household population by age, residence and sex..... 14
Table 2.2	Population by age from selected sources 15
Table 2.3	Household composition 16
Table 2.4	Fosterhood and orphanhood 17
Table 2.5	Educational level of the household population 18
Table 2.6	School enrolment..... 19
Table 2.7	Housing characteristics..... 21
Table 2.8	Household durable goods 22
Table 2.9	Background characteristics of respondents..... 23
Table 2.10	Level of education 25
Table 2.11	Reasons for leaving school 26
Table 2.12	Differential characteristics between spouses..... 27
Table 2.13	Exposure to print media..... 28
Table 2.14	Employment..... 29
Table 2.15	Occupation..... 30
Table 2.16	Decision on use of women's earnings..... 32
Table 2.17	Child care while working..... 33
Table 3.1	Current fertility..... 36
Table 3.2	Fertility by background characteristics 37
Table 3.3	Age-specific fertility rates..... 38
Table 3.4	Fertility by marital duration 39
Table 3.5	Children ever born and living..... 40
Table 3.6	Birth intervals 41
Table 3.7	Age at first birth..... 42
Table 3.8	Median age at first birth by background characteristics..... 43
Table 3.9	Teenage pregnancy and motherhood 44
Table 4.1	Knowledge of contraceptive methods and source for methods..... 45
Table 4.2	Couples' knowledge of contraceptive methods..... 46
Table 4.3	Ever use of contraception 47
Table 4.4	Current use of contraception..... 48
Table 4.5	Current use of contraception by background characteristics..... 50
Table 4.6	Trends in current use of contraception..... 52
Table 4.7	Trends in current use of contraception by residence and region 52
Table 4.8	Number of children at first use of contraception..... 53
Table 4.9	Knowledge of fertile period..... 54
Table 4.10	Timing of sterilisation..... 55
Table 4.11	Source of supply for modern contraceptive methods..... 56
Table 4.12	Source of supply for selected modern methods, 1993 and 1998..... 57
Table 4.13	Contraceptive discontinuation rates..... 58
Table 4.14	Reasons for discontinuation of contraception 58
Table 4.15	Future use of contraception 59
Table 4.16	Reasons for not using contraception..... 60
Table 4.17	Preferred method of contraception for future use 61
Table 4.18	Heard about family planning on radio and television 62
Table 4.19	Acceptability of family planning messages on radio and television..... 63

	Page
Table 4.20	Heard about family planning through print media..... 64
Table 4.21	Wives' perceptions of couple's attitude toward family planning..... 65
Table 4.22	Attitudes of couples toward family planning..... 67
Table 5.1	Number of abortions and stillbirths 70
Table 5.2	Induced abortions by background characteristics 70
Table 5.3	Abortions and stillbirths per 100 pregnancies..... 71
Table 5.4	Trend in induced abortions 71
Table 5.5	Method used before abortion..... 72
Table 5.6	Method used after abortion..... 73
Table 5.7	Reasons for induced abortion 74
Table 5.8	Timing of induced abortions..... 75
Table 5.9	Abortion providers..... 75
Table 6.1	Current marital status..... 77
Table 6.2	Age at first marriage 79
Table 6.3	Median age at first marriage 80
Table 6.4	Age at first sexual intercourse..... 82
Table 6.5	Median age at first intercourse..... 82
Table 6.6	Postpartum amenorrhoea, abstinence and insusceptibility..... 83
Table 6.7	Median duration of postpartum insusceptibility by background characteristics..... 85
Table 6.8	Termination of exposure to risk of pregnancy 85
Table 7.1	Fertility preference by number of living children 88
Table 7.2	Fertility preference by age 89
Table 7.3	Desire to limit (stop) childbearing 90
Table 7.4	Need for family planning services 91
Table 7.5	Ideal and actual number of children..... 93
Table 7.6	Mean ideal number of children by background characteristics..... 94
Table 7.7	Fertility planning status 95
Table 7.8	Wanted fertility rates 96
Table 8.1	Infant and child mortality 99
Table 8.2	Neonatal, post-neonatal, infant, child, and under-five mortality by socioeconomic characteristics..... 101
Table 8.3	Neonatal, post-neonatal, infant, child, and under-five mortality by biodemographic characteristics 102
Table 8.4	High-risk fertility behaviour 103
Table 9.1	Antenatal care..... 106
Table 9.2	Number of antenatal care visits and stage of pregnancy 108
Table 9.3	Tetanus toxoid vaccinations..... 109
Table 9.4	Place of delivery 110
Table 9.5	Assistance during delivery..... 112
Table 9.6	Delivery characteristics: caesarean section, birth weight and size 113
Table 9.7	Delivery complications 114
Table 9.8	Vaccinations by source of information 115
Table 9.9	Vaccinations by background characteristics 116
Table 9.10	Vaccinations in first year of life by current age 117
Table 9.11	Prevalence of diarrhoea 118
Table 9.12	Treatment of diarrhoea 120
Table 9.13	Feeding practices during diarrhoea..... 121

Table 10.1	Initial breastfeeding.....	124
Table 10.2	Breastfeeding status.....	125
Table 10.3	Median duration and frequency of breastfeeding.....	126
Table 10.4	Types of food received by children in the preceding 24 hours	127
Table 10.5	Nutritional status of children by background characteristics	129
Table 10.6	Anthropometric indicators of maternal nutritional status.....	130
Table 10.7	Nutritional status of mothers by background characteristics.....	134
Table 11.1.1	Knowledge of AIDS and sources of AIDS information: women.....	136
Table 11.1.2	Knowledge of AIDS and sources of AIDS information: husbands.....	137
Table 11.2	Knowledge of AIDS and sexually transmitted diseases (STDs).....	138
Table 11.3.1	Knowledge of ways to avoid HIV/AIDS: women.....	139
Table 11.3.2	Knowledge of ways to avoid HIV/AIDS: husbands	140
Table 11.4.1	Knowledge of AIDS-related issues: women	142
Table 11.4.2	Knowledge of AIDS-related issues: husbands.....	143
Table B.1	Allocation of sample households.....	155
Table B.2	Distribution of sample clusters	156
Table B.3.1	Design weights and nonresponse factors: women sample	163
Table B.3.2	Design weights and nonresponse factors: husbands sample	164
Table B.4	Final sample weights	165
Table B.5	Sample implementation: women.....	167
Table B.6	Sample implementation: husbands	168
Table C.1	List of selected variables for sampling errors	173
Table C.2	Sampling errors - National sample.....	174
Table C.3	Sampling errors – Urban areas.....	175
Table C.4	Sampling errors - Rural areas	176
Table C.5	Sampling errors – West region	177
Table C.6	Sampling errors – South region	178
Table C.7	Sampling errors – Central region.....	179
Table C.8	Sampling errors – North region	180
Table C.9	Sampling errors – East region.....	181
Table D.1	Household age distribution	185
Table D.2	Age distribution of eligible and interviewed women	186
Table D.3	Completeness of reporting.....	186
Table D.4	Births by calendar years.....	187
Table D.5	Reporting of age at death in days.....	188
Table C.6	Reporting of age at death in months	189

FIGURES

		Page
Figure 2.1	Population Pyramid	14
Figure 2.2	Number of Persons Reported at Each Age by Sex.....	15
Figure 2.3	School Enrolment by Age and Place of Residence.....	20
Figure 3.1	Age-Specific Fertility Rates by Urban-Rural Residence.....	36
Figure 3.2	Age-Specific Fertility Rates during the Last 20 Years.....	39
Figure 4.1	Current Use of Family Planning Methods, Turkey 1993 and 1998	49
Figure 4.2	Current Use of Family Planning by Region and Method.....	51
Figure 4.3	Knowledge of Fertile Period among All Women.....	54
Figure 4.4	Source of Supply of Modern Contraceptive Methods.....	56
Figure 4.5	Ever Use of Contraception among Non-Users.....	60
Figure 4.6	Attitudes toward Family Planning as Reported by Women	66
Figure 6.1	Marital Status of Women Age 15-49.....	78
Figure 6.2	Median Age at First Marriage among Women 25-49 and Men 25-64	81
Figure 6.3	Percentage of Births Whose Mothers are Amenorrhoeic, Abstaining, or Insusceptible.....	84
Figure 7.1	Fertility Preferences of Currently Married Women, Age 15-49.....	87
Figure 7.2	Under-Five Mortality by Selected Demographic Characteristics.....	85
Figure 8.1	Trends in Infant and Child Mortality	100
Figure 9.1	Source of Antenatal Care by Maternal Age and Birth Order.....	106
Figure 9.2	Antenatal Care by Region and Residence.....	107
Figure 9.3	Place of Delivery by Maternal Age and Birth Order.....	111
Figure 9.4	Percentage of Children Under Five Years with Diarrhoea, by Age, Sex, Birth Order and Residence	119
Figure 10.1	Growth of Children Under Five Years, Mean Z-scores by Age in Months	131

FOREWORD

It is a great pleasure for me to introduce the main findings of the 1998 Turkish Demographic and Health Survey. The findings presented in this volume highlight the major changes that are taking place in the Turkish demographic and health situation. The population of Turkey is going through a structural transition. Mortality rates are going down. Advancements in infant mortality are impressive. Fertility is becoming a matter of choice rather than fate. Replacement levels of fertility are within sight. Significant changes are taking place in age distribution. The proportion of the population living in urban areas is approaching the level observed in industrial societies. These factors, which are both a cause as well as consequence of each other, illustrate the shifting state of Turkish demographic forces.

In view of the dynamic nature of the demographic situation, there is a continuing need for good demographic data in Turkey. "Valid, reliable, timely, culturally relevant and internationally comparable data form the basis for policy and program development, implementation monitoring and evaluation" writes the ICPD Programme of Action adopted at the International Conference on Population and Development in 1994. Good governance is built upon good data. Within this context, data collected through the World Fertility Survey (WFS) and Demographic and Health Survey (DHS) programs have proven their merit globally.

Hacettepe Institute of Population Studies (HIPS) is one of the first academic institutions to collect survey data in the form defined by the WFS-DHS heritage. In fact, HIPS has been contributing to this heritage since its 1968 survey, and preparatory international pilot studies for WFS in the mid-1970s. In the 30 years since 1968, HIPS has successfully carried out six nationwide surveys at regular five-year intervals. The 1998 Turkey DHS is the seventh in its series. I believe that there are not many examples of a similar survey series in the world.

The 1998 Turkish Demographic and Health Survey (TDHS-98) was conducted through an agreement with Macro International Inc. under the auspices of a the MEASURE DHS+ project supported by the United States Agency for International Development. The TDHS-98 is the second demographic survey carried out in collaboration with Macro International Inc. Contributions of the United Nations Population Fund also were critical in realisation of the survey in its present scope. Without the UNFPA contribution, the scope of the survey would have been much more limited.

The TDHS-98 findings are crucial in monitoring trends in demographic and health variables and in understanding the factors that contribute to differentials in fertility and contraceptive use among various population subgroups. The data also are important for understanding the factors that influence the health and survival of infants and young children. In addition, the TDHS-98 provides a wealth of information for husbands and never-married women. Beyond providing insights into population and health issues in Turkey, it is hoped that the TDHS-98 will also lead to an improved global understanding of population and health problems, as one of more than 80 surveys implemented and sponsored by the Demographic and Health Surveys Program.

This TDHS-98 could have been realised only with the industrious contributions of many individuals. I would like to express my appreciation to the HIPS staff for their devotion and sincere efforts in accomplishing the planned activities on time and to Dr. Banu Ergöçmen and Dr. Edilberto Loaiza for preparing this presentation of the survey findings. I would like to thank Dr. Ann Way who had a large part at the inception of the project as well as at the finalization of the main report. Dr. Pinar Senlet, USAID mission in Turkey, tactfully paved the way through difficulties. I also would like to acknowledge the contributions of Dr. Mona Kaidbey and other UNFPA staff through their representative, Dr. Alain P. Mouchiroud. The Population Sector of the State Planning Organisation through her able director Dr Samira Yener was critically instrumental in getting the survey started. I thank them all. My

thanks are also due to the Steering Committee Members for their valuable contributions and to the State Institute of Statistics for their assistance in sample selection. I would like to express my gratitude to Prof. Dr. Süleyman Sağlam, the Rector of Hacettepe University, for his continuous encouragement and support during all phases of the survey.

Finally, my special thanks go to Dr. Rifat Köse and to his dedicated staff at the General Directorate of Mother and Child Health/Family Planning in the Ministry of Health. Their support was indeed crucial and is a good example of a cooperation between a governmental body and an academic institution.

Altogether it was a tough task to accomplish, although it was a pleasant enterprise for those who did it.

Prof. Dr. Aykut Toros
Director
Institute of Population Studies
Hacettepe University

SUMMARY OF FINDINGS

The 1998 Turkish Demographic and Health Survey (TDHS-98) is a nationally representative sample survey designed to provide information on fertility levels and trends, infant and child mortality, family planning, and maternal and child health. Survey results are presented at the national level, by urban and rural residence and for each of the five regions in the country.

The survey was fielded between August and November 1998. Hacettepe University Institute of Population Studies (HIPS) carried out the TDHS-98 in collaboration with the General Directorate of Mother and Child Health and Family Planning, Ministry of Health. Funding for the TDHS-98 was provided both by the U.S. Agency for International Development through the MEASURE/DHS+ program and United Nations Population Fund.

Interviews were carried out in 8,059 households, with 8,576 women, and with 1,971 husbands. All women at ages 15-49 who were present in the household on the night before the interview or who generally live in that household were eligible for the survey. In half of the selected households for women interview, husbands (of currently married eligible women), who were present in the household on the night before the interview or who generally live in that particular household were eligible husbands for the survey.

Survey results indicate that if Turkish women maintain current fertility rates during their reproductive years, they can expect to have an average of 2.6 children by the end of their reproductive years. The highest fertility rate is observed for the age group 20-24. There are marked regional differences in fertility rates, ranging from 4.2 children per woman in the East to 2.0 children per woman in the West. Fertility also varies widely by urban-rural residence and by education level. A woman living in rural areas will have almost one child more than a woman living in an urban area. However, when compared with evidence from previous surveys it is worth noting that the urban/rural gap appears to be closing. Women who have no education have almost one child more than women who have a primary-level education and almost 2 children more than women with secondary-level education.

The first requirement of success in family planning is the knowledge of family planning methods. Knowledge of modern method is almost universal among Turkish women. The IUD and pill are the most widely known family planning methods among women and husbands followed by the condom and female sterilisation. Eighty-four percent of currently married women have used a method sometime in their life. Thirty-nine percent of currently married women report ever using the IUD. Overall, 64 percent of currently married women are currently using a method. The majority of these women are modern method users (38 percent), but more than one fourth use traditional methods (26 percent). The IUD is the most commonly used modern method (20 percent), followed by the condom (8 percent) and the pill (4 percent). Regional differences are substantial. The level of current use is 42 percent in the East, 71 percent in the West and more than 60 percent in the other three regions.

A basic knowledge of reproductive physiology is necessary, especially in the use of coitus-related methods. However, only 18 percent of all women know the correct time of ovulation. Information on the sources of methods is important for planning the services. The majority of modern method users (56 percent) obtain the methods from public sector. Health centres are the major public sector suppliers (23 percent) and pharmacies are the major private sector suppliers (25 percent).

The discontinuation rate of the IUD is the lowest among all methods. Information on the intentions of current non-users was also collected for the estimation of future demand. Of this group, 44 percent do not intend to use any method in the future whereas half of the currently married women who are not using any

contraception have the intention to use in future. Of the latter women, more than one third report that their method of choice will be the IUD.

During the five year period before the survey, almost one in four pregnancies terminated in other than a live birth. Only about 2 out of every 100 pregnancies ended in a stillbirth and of the 23 abortions per 100 pregnancies 15 were induced. More than one fourth of ever-married women reported ever having had an induced abortion during their reproductive lives. However, among the women who had had an induced abortion, around half had had only one induced abortion. Women living in the East region and in the rural settlements are the least likely to have ever had an induced abortion. There is little variation by education in the proportion of ever-married women who have had an induced abortion.

There is a very important opportunity for family planning counselling after an abortion. However, the results show that this opportunity is not utilised well. In the month after an induced abortion, 32 percent of women did not use any method and 27 percent used withdrawal. The main reason for obtaining an abortion was the desire not to have any more children (62 percent). Overall, 68 percent of abortions took place in the first month of pregnancy, 23 percent in the second month, and 9 percent in the third or later months of pregnancy. Some 74 percent of abortions took place at a private sector; there is variation between regions in terms of the place where induced abortions are performed.

The age at first marriage is one of the important determinants of fertility. TDHS-98 results suggest that there is a steady increase in the median age at first marriage, with respect to cohorts. The median age at first marriage has increased from 18.4 years for the 45-49 age group to 20 years for the 25-29 age group. There are differences in the age at marriage across places of residence and regions. Even more pronounced differences are observed by educational level of women. Among women age 25-49, there is a difference of almost 6 years in the timing of entry into marriage between those with little or no education and those who completed at least the secondary level. The survey findings indicate that husbands enter into first marriage at much later age than women. The median age at first marriage is 23.5 for the husbands in age group 25-29, and it is 24 for the 30-34 and 35-39 age groups.

Two-thirds of currently married women in Turkey say that they do not want any more children or are already sterilised for contraceptive purposes. An additional 14 percent want to wait at least two years before having another child. Sixty percent of husbands at ages 30-34 want to have no more children. A strong desire for a two-child family is evident. Results from the survey suggest that if all unwanted births were eliminated, the total fertility rate at the national level would be 1.9 children per woman, almost one child lower than the actual level of 2.6. Nineteen percent of the births in the five years preceding the survey were unwanted births and 11 percent of them were mistimed. The unmet need for family planning in Turkey indicates that there is potential for further increases in contraceptive use. Ten percent of currently married women are considered to be in need of a family planning method either for spacing or for limiting.

For the five years preceding the TDHS-98, the infant mortality rate is estimated at 43 per thousand, the child mortality rate at 10 per thousand, and the under-five mortality rate at 52 per thousand. For the same period, results show that in Turkey, the neonatal mortality rate is higher than the postneonatal mortality rate, and that all the indicators of infant and child mortality have declined rapidly in recent years.

The TDHS-98 findings point to significant differences in infant and child mortality between regions and urban and rural areas, and show that the educational level of the mother and the presence of medical maternity care are important correlates of infant and child mortality. In addition to the differentials observed between socioeconomic groups, infant and child mortality rates also appear to correlate strongly with demographic variables. Age of mother at birth and order of birth show the expected U-shaped relationship with infant and child mortality. High birth order and short birth intervals are the major factors contributing to elevated risks of mortality.

Among the maternal health indicators, antenatal care was received from trained health personnel by 68 percent of pregnant women. It is preferred that antenatal care is sought early in pregnancy and is continued throughout a pregnancy. In Turkey, in 58 percent of the births, antenatal care was sought before the sixth month of pregnancy and the median number of antenatal care visits is 4.2. Tetanus toxoid coverage for women is low, with 15 percent having one dose and 29 percent having two doses or more. The TDHS-98 shows that 73 percent of all deliveries took place at a health facility.


One of the major child health indicators is immunisation coverage. Among children age 12-23 months, the coverage rates for BCG and the first dose of DPT are 87 and 86 percent respectively, with most of the children receiving those vaccines before age one. The results indicate that only 40 percent of the children had received all vaccinations at some time before the survey. On a regional basis, coverage is significantly lower in the East region, followed by the West and Central regions. Diarrhoea is a prevalent disease of children under age five in Turkey. In the two weeks preceding the survey, the prevalence of diarrhoea was 30 percent for children under age five. Among children with diarrhoea 70 percent were given more fluids than usual.

Breastfeeding in Turkey is widespread. Almost all Turkish children (95 percent) are breastfed for some period of time. The median duration of breastfeeding is 14 months, but supplementary foods and liquids are introduced at an early age. Almost half of the children are being given supplementary food as early as one month of age.

By age five, almost one-quarter of children are stunted (short for their age), compared to an international reference population. Stunting is more prevalent in rural areas, in the East, among children of mothers with little or no education, among children who are of higher birth order, and among those born less than 24 months after a prior birth. Overall, wasting is not a problem. Two percent of children are wasted (thin for their height), and 8 percent of children under five are underweight for their age. The survey results show that obesity is a problem among mothers. According to Body Mass Index (BMI) calculations, 52 percent of mothers are overweight, of which 19 percent are obese.

AIDS is widely known in Turkey. Overall 84 percent of women and 93 percent of husbands have heard about AIDS. However, although women and husbands generally know AIDS, knowledge of ways to avoid it appears to be poor among a substantial minority of both groups. Additionally, the general perception of the disease in Turkey is that it is almost always fatal; 65 percent of women and 74 percent of husbands stated AIDS as an almost always a fatal disease. Sexually transmitted diseases (STD) other than AIDS are not known as widely as AIDS is known. Thirty-five percent of currently married women and 27 percent of never-married women reported having heard of STDs. STDs are more widely known by husbands; 54 percent of the husbands mentioned knowing STDs.

TURKEY



IIIX



REGIONS AND PROVINCES

WEST

09 Aydın
10 Balıkesir
16 Bursa
17 Çanakkale
20 Denizli
22 Edirne
34 İstanbul
35 İzmir
39 Kırklareli
41 Kocaeli
45 Manisa
54 Sakarya
59 Tekirdağ
77 Yalova

SOUTH

01 Adana
07 Antalya
15 Burdur
27 Gaziantep
31 Hatay
32 Isparta
33 İçel
48 Muğla
79 Kilis
80 Osmaniye

CENTRAL

03 Afyon
05 Amasya
06 Ankara
11 Bilecik
14 Bolu
18 Çankırı
19 Çorum
26 Eskişehir
38 Kayseri
40 Kırşehir
42 Konya
43 Kütahya
50 Nevşehir

NORTH

08 Artvin
28 Giresun
37 Kastamonu
52 Ordu
53 Rize
55 Samsun
61 Trabzon
67 Zonguldak
74 Bartın
78 Karabük

EAST

02 Adıyaman
04 Ağrı
12 Bingöl
13 Bitlis
21 Diyarbakır
23 Elazığ
24 Erzincan
25 Erzurum
29 Gümüşhane
30 Hakkari
36 Kars
44 Malatya
46 K. Maraş
47 Mardin
49 Muş
56 Sırt
58 Sivas
62 Tunceli
63 Ş. Urfa
65 Van
69 Bayburt
72 Batman
73 Şırnak
75 Ardahan
76 Iğdır

CHAPTER 1

INTRODUCTION

Sunday Üner and Banu Akadlı Ergöçmen

1.1 Geography

Turkey occupies a surface area of 774,815 square kilometres. About three percent of the total area lies in Southeastern Europe (Thrace) and the remainder in Southwestern Asia (Anatolia or Asia Minor). Turkey has borders with Greece, Bulgaria, Syria, Iraq, Iran, Georgia, Armenia, and Nahcivan (Azerbaijan). The shape of the country resembles a rectangle, stretching in the east-west direction for approximately 1,565 kilometres and in the north-south direction for 650 kilometres. The three sides of Turkey are surrounded by seas: in the north, the Black Sea; in the northwest, the Sea of Marmara; in the west, the Aegean Sea; and in the south, the Mediterranean Sea. The total coastline of Turkey is approximately 8,333 kilometres.

Anatolia lies on a semi-arid central plateau that is surrounded by mountains. It is for the most part an elevated steppe-like plateau enclosed by mountains on all sides but the west. The Taurus Mountains in the south and the Northern Anatolia Mountains in the north stretch parallel to the coastline, these two mountains meet in the eastern part of the country. The average altitude of the country is approximately 1,130 metres above sea level. However, there are vast differences among regions, ranging from an average of 500 metres in the west to 2,000 metres in the east.

The climate is characterized by variations of temperature and rainfall, depending on topography of the country. The average rainfall is 500 millimetres; however, it ranges from 2,000 millimetres in Rize, a province on the Black Sea coast, to less than 300 millimetres in parts of Central Anatolia. The typical climatic conditions of Turkey include dry, hot summers and cold, rainy winters. In summer, temperatures do not vary greatly across the country, whereas in winter, the temperature ranges from an average of -10°C in the east to $+10^{\circ}\text{C}$ in the south.

1.2 History

Before becoming the core of the Ottoman Empire, Anatolia was dominated by the Seljuqs for almost two centuries (1055-1243) and later became the core of the Ottoman Empire. The Ottoman Empire was one of the most powerful forces in the Middle East and Europe for nearly 600 years. Following the War of Independence led by Mustafa Kemal Atatürk, the Republic of Turkey was founded in 1923. Subsequently, the country's borders were shaped by a number of agreements, with its present borders finally established following the annexing in 1939 of Hatay, a province on the southern border.

The founding of the Republic not only marked the end of the Ottoman era and the establishment of the present borders of modern Turkey, but also signified a revolutionary change in the social and economic structure of the country. A modern constitution was introduced, the Sultanate and Caliphate were abolished, as were the veil and the traditional headgear. The Latin alphabet was adopted instead of Arabic script. The schools were taken out of the hands of the religious authorities, and a program of free, compulsory education was set up. Islamic Law was abandoned and replaced with modified versions of the European civil, penal and commercial codes, and religious courts were closed. In short, the direction of change, led by Atatürk, was one from a religious, oriental Empire to a modern, westernised secular Republic.

From the foundation of the Turkish Republic to 1950, the country was governed by one party. This changed in 1950 as a result of the wind of democracy that caught up the country after the Second World War. In the first multiparty election held in 1950, the Democrat Party won, putting the Republican People's Party into the opposition. With the introduction of multi-party period, Turkey achieved a more liberal and democratic environment. Although Turkish political history included three military interventions (1960, 1971, and 1980), Turkey has succeeded in preserving a parliamentary, multi-party system until today.

With the foundation of the republic, Turkey turned her face to the 'Western world', establishing close relations with European countries and especially with the United States of America. Turkey is a member of the United Nations, the Council of Europe and the North Atlantic Treaty Organization (NATO) and an associate member of the European Community. Turkey also maintains close relations with the countries of the Middle East, stemming from deep-rooted cultural and historical links.

1.3 Administrative Divisions and Political Organisation

Since the founding of the Republic, three Constitutions (1924, 1961, and 1982) have shaped the Turkish administrative structure. These three constitutions proclaimed Turkey to be a Republic with a parliamentary system and specified that the will of the people is vested in the Turkish Grand National Assembly (TGNA). All three constitutions adopted basic individual, social and political rights, and accepted the principle of separation of powers.

The legislative body of the Republic is the TGNA. The TGNA is composed of 550 deputies, who are elected for five-year terms. The President of the Republic is elected by the TGNA for a seven-year term. The Prime Minister and other Cabinet Ministers compose the Council of Ministers, the executive branch of the Republic. The judiciary consists of the Court of Appeals, the Court of Jurisdictional Disputes, the Military Court of Appeals, the Constitutional Court, and the civil and military Courts.

Turkey is administratively divided into 80 provinces. These are further subdivided into districts (ilçe), subdivisions (bucak), and villages. The head of the province is the governor, who is appointed by and responsible to the central government. The governor, as the chief administrative officer in the province, carries out the policies of the central government, supervises the overall administration of the province, coordinates the work of the various ministry representatives appointed by the central authority in the capital Ankara, and maintains law and order within his/her jurisdiction.

A mayor and a municipal council administer local government at the municipality level. The municipal electoral body elects them for a term of four years. Every locality with a population of more than 2,000 is entitled to form a municipal administration. Municipalities are expected to provide basic services such as electricity, water, gas, the building and maintenance of roads, and sewage and garbage disposal facilities. Educational and health services are mainly provided by the central government, but municipalities also provide some health services.

1.4 Social and Cultural Features

Turkey has a highly heterogeneous social and cultural structure. There are sharp contrasts between population groups. The "modern" and "traditional" exist simultaneously within the society. Attitudes to life are reminiscent of those in the Western world, especially for the inhabitants of metropolitan areas. On the other hand, people living in rural areas are more conservative and religious. Family ties are strong and influence the formation of values, attitudes, aspirations, and goals. Although laws are considered to be quite liberal on gender equality, patriarchal ideology still may characterize the social life.

Citizens of Turkey are predominantly Muslim. About 98 percent of the population belong to the Sunni and Alawi sects of the Muslim religion, with the Sunnis forming the overwhelming majority. Turks predominate ethnically. Kurdish, Arabic, Greek, Circassian, Georgian, Armenian, and Jewish communities of varying sizes complete the ethnic mosaic of the rich and complex culture of the Turkish society.

One of the most striking achievements since the founding of the Republic has been the increase in both literacy and education. In 1935, only 10 percent of females and 29 percent of males were literate in Turkey. According to the 1990 census figures, the female and male literacy rates for the population age 6 and over were 72 and 89 percent, respectively. Educational attainment has also increased dramatically. The rate for primary school attendance today is around 90 percent. Moderate advances have also been made in increasing the proportions of males and females with higher than primary-level education. In 1998, an eight-year education became compulsory in Turkey, with primary school encompassing the first 5 years and junior high school, 3 years. Despite these achievements, considerable regional and urban-rural differences in literacy and educational attainment continue to exist in the country in addition to the gender differences.

1.5 Economy

The Turkish Republic inherited from the Ottoman Empire not only a bankrupt country, but also centuries-old traditions of instability and insolvency. After the foundation of the Turkish Republic, various economic development strategies were tried. In the early years of the Republic, the Turkish economy was very weak. The economy was almost exclusively based on the agriculture, and it was totally undeveloped and poor. The creation and development of industry was clearly the first step that had to be taken to achieve a healthy and balanced economy. Throughout the 1920s, the government promoted the development of industry through private enterprise, encouraged and assisted by favourable legislation and the introduction of credit facilities. These liberal policies continued until 1929, and moderate improvements were realised in the mechanisation of agriculture. In the following decade, the state, under the so-called *étatiste* system, assumed the role of entrepreneur, owning and developing large sectors of agriculture, industry, mining, commerce and public works. The origins of modern industrialisation in Turkey can be traced to the era of the 1930s. Although the beginnings of the industrialisation drive were evident in the immediate aftermath of the formation of the republic in 1923, the real breakthrough occurred in the context of the 1930s.

Although Turkey did not actually participate in the Second World War (Turkey initially remained neutral during the Second World War but eventually sided with Allies), the country was faced with heavy restraints on the economy, which slowed down the industrialization process. A "mixed economy" regime followed the war, with the transition to democracy in 1950 signifying a shift towards a more liberal economic order; private enterprise gained recognition side by side with the state economic enterprises. Also, more emphasis was placed on trade liberalization, agricultural and infrastructural development, and the encouragement of foreign capital.

A series of Five-Year Development Plans were prepared after the military intervention in 1960. The first of these plans became operative in 1963. Imports were limited, and protectionism was established as the dominant economic strategy during the 1960s and the 1970s. A basic objective was to replace the era of unplanned and uncontrolled expansion during the 1950s.

In the 1980s, governments followed a strategy of renewing economic growth based on an export-oriented strategy. Following the stagnation of the late 1970s, growth recovered in response to a combination of an increased flow of exports and inputs of foreign capital. The liberal economic strategy followed in the 1980s was not unique to that period. The differences between the liberal and *étatiste* phases are not only the nature of the trade regime and the attitude toward foreign direct investment (FDI),

but also the mode of state intervention in the economy. Respectable rates of economic growth were achieved during the 1980s; however, in recent years, macro instability has manifested itself once again.

Industrialization during the 1990s has been shaped by three dynamics. First, the state's direct influence on the distribution of the resources was lessened. Second, competition gained importance, with increased emphasis on industrial performance and reconstruction of the industry. Third, general globalisation and integration into the European Union gained speed. During the 1990s, privatisation also gained importance as a solution to economic problems. A committee was founded in order to regulate privatisation. Some of the state enterprises were privatised during this program.

Turkey is a self-sufficient country in terms of its agricultural production. Wheat, barley, sugar beets, potatoes, and rice are grown in the interior, and cotton, tobacco and citrus are grown for export around the coastal areas. Turkey is not rich in mineral resources. One of the country's main problems is the inadequacy of primary energy resources. Copper, chromium, borax, coal, and bauxite are among the mineral resources in the country. The main industries are steel, cement, textiles, and fertilizers. Machinery, chemicals and metals are imported mainly from the OECD countries. In recent years, there has been a significant increase in the amount of industrial goods exported to Europe and Arab countries.

Turkey can be classified as a middle-income country in the late 1990s. The rate of economic growth has been comparatively high in recent years, and the economy has undergone a radical transformation, from an agricultural base to an industrial one, particularly within the last 20 years.

1.6 Regional Breakdown

The diverse geographical, climatic, cultural, social, and economic characteristics of different parts of the country is the basis for the conventional regional breakdown within Turkey. Five regions (West, South, Central, North, and East) are distinguished, reflecting, to some extent, differences in socioeconomic development levels and demographic conditions among sections of the country. This regional breakdown is frequently used for sampling and analysis purposes in social surveys.

The West region is the most densely settled, the most industrialized, and the most socioeconomically advanced region of the country. The region includes both Istanbul, (previously the capital of the Ottoman Empire), which is Turkey's largest city, and the country's manufacturing and commercial centre, and Izmir, the country's third largest city. Coastal provinces form a relatively urbanized, fast-growing area. The Aegean coast is also a major agricultural area, where cotton is grown in the river valleys, and fruit is cultivated on the hillsides. With dry summers and mild, rainy winters, agricultural yields from the fertile soils are good. Most of the industrial establishments are situated in the West. The region contributes most of the gross domestic product of the country.

The South includes highly fertile plains and some rapidly growing industrial centres. Adana, one of the new metropolises of Turkey, is located in this region. Steep mountains cut off the semitropical coastal plains from the Anatolian highlands to the north. Hot, dry summers and mild, wet winters describe the climatic conditions of the region. Cultivation of cotton and citrus provide high incomes and export earnings; recent decades have witnessed an industrial boom and an inflow of migrants, especially from the East region.

The Central region is an arid grazing area and includes Ankara, the capital and second largest city. Industrial production in the region is low, except for some minor industries located around Ankara. The region specializes in the production of cereals. Given the dry, temperate climate, fruit tree cultivation and sheep and cattle raising are also common.

The North region has a fertile coastal strip, but in most places it is only a few kilometres wide; the region is relatively isolated from the rest of the country by mountainous terrain. The region specialises

in small-scale, labour-intensive crops like hazelnuts and tea. The region receives large quantities of rainfall. Zonguldak, a western province, has extensive coal reserves and is a centre for mining and the steel industry.

The East region is considered as the least developed part of the country. Rugged mountainous terrain, short summers, and the severe climate are suited to animal husbandry rather than settled farming. However, with the "Southeast Anatolia Project", the economy in the East has improved in the recent years. Huge irrigation channels were constructed and water was provided to arid and semi-arid lands, leading to agricultural development in the Southeast Anatolia. In addition to economic benefits, the project is also expected to reverse the migration flow from the region to the rest of the country. Although the capacity of agriculture has increased, the region is still poor in terms of industrial production.

1.7 Population

In 1927, Turkey's population was 13.6 million according to the census, which was conducted four years after the establishment of the Republic. Beginning with the 1935 census, subsequent population censuses were undertaken at 5-year intervals, with the last complete census occurring in 1990. In order to be able to prepare the electoral rolls, a General Population Register was carried out in 1997; the results of the Register put the population of Turkey at 62.8 million. Turkey is among the 20 most populous countries of the world and is the most populous country of the Middle East (State Institute of Statistics, 1999; Population Reference Bureau, 1999).

Intercensal estimates of population growth have been around 20-25 per thousand since the 1970s. The latest estimate of the population growth rate was 15.1 per thousand for the 1990-1997 period. According to the projections, the population of Turkey is expected to reach 76 million in the year 2010 and 88 million in 2025 (Population Reference Bureau, 1999).

Turkey has a young population structure as a result of the high fertility and growth rates of the recent past. Recent decades have witnessed dramatic declines especially in fertility rates. In the early 1970s, the total fertility rate was around 5 children per woman, whereas the estimates in the early 1990s put the total fertility rate at less than 3 children. The crude birth rate is estimated at 22 per thousand for the mid-1990s (Population Reference Bureau, 1999).

There is a shortage of information on mortality in Turkey, particularly adult mortality. However, due to the estimation of the indicator through fertility surveys, infant mortality rates can be traced back for a relatively long period of time. The infant mortality rate in the late 1950s was around 200 per thousand. It declined to about 130 per thousand during the mid-1970s and to an estimated 53 per thousand during the early 1990s. Crude death rates have also declined from around 30 per thousand in the 1940s to 7 per thousand in the mid-1990s. The latest estimates put life expectancy in Turkey at 66 years for males and 71 for females (Population Reference Bureau, 1999).

Marriage, predominantly civil, is widely practised in Turkey. Religious marriages also account for a significant proportion of the marriages; however, the main custom is to undergo a civil as well as a religious ceremony. The universality of marriage in Turkey is observed in the proportions never married. According to the 1990 Population Census, in the age group 45-49 which marks the end of the reproductive ages, only two percent of females had never married, whereas the corresponding figure for males in the same age group was three percent. Marriages in Turkey are also known to be very stable; divorce rates are very low (Hancıoğlu and Akadlı Ergöçmen, 1992).

The population of Turkey has undergone an intensive process of urbanization, especially from the 1950s onwards. According to the 1997 Population Count, 65 percent of the population in Turkey are living in urban settlements (province and district centers). The rate of urbanisation has been approximately 50 per thousand during the 1970-1990 period. The process of urbanisation has inevitably

caused problems in the provision of services and the emergence of large areas of squatter housing in unplanned cities.

Since the early 1960s, Turkey has had a long history of external migration to Western European countries, principally Germany. Migration to Western Europe continued throughout the 1960s and 1970s; at that point, migration increasingly was directed towards oil-producing countries of the Middle East. During the past two decades, however, the political turmoil in region and changes in policies and practices governing the labour force in the European Union have led to a reversal in migration trends. There has been an increase in the number of expatriate workers returning from Europe, either because of loss of opportunity or as a result of having achieved their savings targets. Meanwhile, political conditions in neighbouring countries have led to waves of immigration into Turkey, as well (UNFPA, 1995).

1.8 Population and Family Planning Policies and Programs

Policies related to population have been formulated since the establishment of the Turkish Republic in 1923. The government of the Turkish Republic implemented a somewhat pronatalist population policy aimed at increasing the population size, until the mid-1960s, after which an antinatalist policy was adopted. The shift in policy is manifested in the Population Planning Law of 1965.

During the early years of the Republic there was an apparent need to increase fertility, since the country was suffering from the heavy human losses during the First World War and the War of Independence. The defence needs of the country and the shortage of manpower, as well as high infant and child mortality rates, led Turkey to follow a pronatalist population policy until the late 1950s. A number of laws having direct or indirect implications for encouraging population growth were passed. These laws included monetary awards to women with more than 5 children, prohibitions on the import and sale of contraceptives, and prohibitions on abortions on social grounds.

The high population growth rates prevailing in the 1950s produced medical problems, particularly the high maternal mortality caused by illegal abortions, which brought the population debate into the political agenda. High urban population growth and employment problems were also factors contributing to the new antinatalist environment in government circles. The State Planning Organisation and the Ministry of Health pioneered the policy change, allowing limited importation of contraceptives. As mentioned earlier, the Population Planning Law was enacted in 1965. The law mandated the Ministry of Health with the responsibility for implementing the new family planning policy. In addition, the State Planning Organisation incorporated the notion of population planning in the First Five-Year Development Plan.

In 1983, the Population Planning Law was revised and a more liberal and comprehensive law was passed. The new law legalized abortions (up to the tenth week of pregnancy) and voluntary surgical contraception. It also permitted the training of auxiliary health personnel in inserting IUDs and included other measures to improve family planning services and mother and child health.

1.9 Health Priorities and Programs

Mother and child health and family planning services have been given a priority status in the policies of the government in recent decades. These services gained importance due to the large proportion of women of reproductive ages and children in the Turkish population, the high infant, child and maternal mortality rates, the high demand for family planning services, and the limited prenatal and postnatal care. A number of programs to improve services are being implemented, with special emphasis on provinces which have been designated as priority development areas as well as on squatter housing districts in metropolitan cities, rural areas, and special risk groups. The initiatives include programs in immunisation, childhood diarrhoeal diseases, acute respiratory infections, promotion of breastfeeding and

growth monitoring, nutrition, reproductive health, and antenatal and delivery care, safe motherhood. Information, Education, and Communication programs to promote the mother and child health and family planning activities also are being widely implemented.

1.10 Health Care System in Turkey

The Ministry of Health is officially responsible for designing and implementing nationwide health policies and delivering health-care services. Besides the Ministry of Health, other public sector institutions and non-governmental organisations contribute to providing health services.

At the central level, the Ministry of Health is responsible for the implementation of curative and preventive health-care services throughout the country, within the principles of primary health care. The responsibility for delivering the services and implementing specific Primary Health Care programs is shared by various General Directorates (Primary Health Care, Mother and Child Health and Family Planning, Health Training) and by various Departments (Departments of Tuberculosis Control, Malaria Control, Cancer Control).

At the provincial level, the health care system is the responsibility of Health Directorates, under the supervision of the Governor. The provincial Health Director is responsible for delivering all primary health-care services as well as curative services. The present network of Health Centres and Health Houses was formed on the basis of "Legislation for the Socialization of Health Services" so that services and facilities are extended down to the village level. A substantial proportion of villages has health centres or health houses. These are located so as to provide easy access to other villages.

The most basic element of the health service is the Health House, which serves a population of 2500-3000 and is staffed by a midwife. The Health Centre serves a population of 5,000-10,000 and is staffed by a team consisting of a physician, a nurse, a health officer, midwives, an environmental health technician, and a driver. Health Centres mainly offer integrated, polyvalent primary health-care services. Mother and Child Health and Family Planning Centres and Tuberculosis Dispensaries also offer preventive health services.

This network of health facilities are responsible for delivering primary health services, maternal and child health, family planning, and public education services. These health facilities are also the main sources of the health information system.

1.11 Objectives and Organisation of the Survey

Objectives

The 1998 Turkish Demographic and Health Survey (TDHS-98) is the latest in a series of national-level population and health surveys that have been conducted during the last thirty years in Turkey. The primary objective of the TDHS-98 is to provide data on fertility and mortality, family planning, maternal and child health, and reproductive health. The survey obtained detailed information on these issues from a sample of women in the reproductive ages (15-49) and from the husbands of currently married eligible women.

More specifically, the objectives of the TDHS were to:

Collect data at the national level that allow the calculation of demographic rates, particularly fertility and childhood mortality rates;

- Obtain information on direct and indirect factors that determine levels and trends in fertility and childhood mortality;
- Measure the level of contraceptive knowledge and practice by method, region, and urban-rural residence;
- Collect data on mother and child health, including immunisations, prevalence and treatment of diarrhoea among children under five, antenatal care, assistance at delivery, and breastfeeding;
- Measure the nutritional status of children under five and of their mothers using anthropometric measurements.

The TDHS-98 information is intended to contribute data to assist policy makers and administrators to evaluate existing programs and to design new strategies for improving demographic, social and health policies in Turkey.

Organisation

The TDHS-98 was implemented by the Institute of Population studies, Hacettepe University, in collaboration with the General Directorate of Mother and Child Health/ Family Planning, Ministry of Health. Technical and financial support for the survey were provided both by United Nations Population Fund (UNFPA) and Macro International Inc. Macro's assistance was provided through the MEASURE/DHS+ project, a project sponsored by the United States Agency for International Development (USAID) to carry out population and health surveys in developing countries.

A steering committee consisting representatives of the Institute of Population Studies, Hacettepe University, the General Directorate of Mother and Child Health/Family Planning, Ministry of Health, the State Planning Organization, the State Institute of Statistics, the United Nations Population Fund (UNFPA) and Macro International Inc. participated in all phases of the project.

Questionnaires

Four main types of questionnaires were used in the TDHS-98: the Household Questionnaire and three Individual Questionnaires, one for ever-married women of reproductive ages, one for never-married women, and one for husbands. The contents of the questionnaires were based on the DHS Model 'A' Questionnaire, which was designed for the DHS program for use in countries with high contraceptive prevalence. Additions, deletions and modifications were made to the model questionnaire in order to collect information particularly relevant to Turkey. In developing the questionnaire, close attention was paid to obtaining the data needed for program planning in Turkey as specified during consultations with population and health agencies. Ensuring the comparability of the TDHS-98 findings with previous demographic surveys carried out by the Hacettepe Institute of Population Studies also was a goal during questionnaire development. The questionnaires were developed in English and translated into Turkish.

The Household Questionnaire was used to enumerate all usual members of and visitors to the selected households and to collect information relating to the socioeconomic situation of the households. In the first part of the household questionnaire, basic information was collected on the age, sex, educational attainment, marital status and relationship to the head of household of each person listed as a household member or visitor. The objective of the first part of the Household Questionnaire was to obtain the information needed to identify women and husbands who were eligible for the individual interview as well as to provide basic demographic data for Turkish households. The second part of the Household Questionnaire included questions on the welfare of the aged people. In the third part of the Household

Questionnaire, questions were included on the dwelling unit, such as the number of rooms, the flooring material, the source of water, and the type of toilet facilities, and on the household's ownership of a variety of consumer goods.

The Individual Questionnaire for women¹ covered the following major topics:

- Background characteristics
- Reproduction
- Marriage
- Knowledge and use of family planning
- Maternal care and breastfeeding
- Other issues relating to contraception
- Immunisation and health
- Fertility preferences
- Husband's background
- Women's work and status
- Sexually transmitted diseases and AIDS
- Maternal and child anthropometry.

The ever-married women's questionnaire included a monthly calendar, which was used to record fertility, contraception, marriage and migration histories for an approximate period of six years, beginning in January 1993 through the month of interview. The monthly calendar in the never-married women's questionnaire included only the migration history. In addition, the fieldwork teams measured the heights and weights of children under age five and of all women ages 15-49.

The Individual Questionnaire for husbands covered the following major topics:

- Background characteristics
- Reproduction
- Knowledge and use of family planning
- Marriage
- Fertility preferences
- Sexually transmitted diseases and AIDS
- Attitudes

Sample

The sample for the TDHS-98 was designed to provide estimates of population and health indicators including fertility and mortality rates for the nation as a whole, for urban and rural areas, and for the five major regions of the country (West, South, Central, North and East). A weighted, multi-stage, stratified cluster sampling approach was used in the selection of the TDHS-98 sample.

The optimal distribution with a target sample size of 10,000 selected households was based on the provisional results of the 1997 General Population Count. Selection of the TDHS-98 sample was undertaken in three stages. The sampling units at the first stage were the settlements stratified by population size. The frame for the selection of the primary sampling units (PSU) was prepared using the provisional results of the 1997 Population Count. The frame was divided into two groups, one including those settlements with populations of more than 10,000 and the other including settlements with populations less than 10,000. The selection of the settlement in each group was carried out with probability proportional to size (1997 population).

¹ For the never-married women some of the topics were not covered in the questionnaire.

The second stage of selection required the selection of the assigned number of clusters in each selected settlement. For the majority of the settlements (340 clusters), the selection of clusters was based on the household lists that were available from the 1995 Structure Schedules. The State Institute of Statistics (SIS) selected the clusters and provided to Hacettepe Institute of Population Studies a description of each selected cluster. Each cluster included approximately 100 households. For those settlements where SIS was not able to provide information (140 clusters), the lists of households were prepared in the field.

Following the selection of the secondary sampling units (SSUs), a household listing was prepared or updated for each SSU by the TDHS-98 listing teams. Using the household lists, a systematic random sample of fixed number of households (25 in clusters located in settlements over 10,000 and 15 in those less than 10,000) was chosen within each cluster for the TDHS-98. All women at ages 15-49 who were present in the household on the night before the interview were eligible for the survey. In half of the selected households for women interview, husbands of currently married eligible women, who were present in the household on the night before the interview or who usually lived in the household were eligible for the husband survey.

A more technical and detailed description of the TDHS sample design, selection and implementation is presented in Appendix B.

Fieldwork and Data Processing

The TDHS-98 data collection was carried out by 16 teams²; each team consisted of four to five female interviewers, one male interviewer, a field editor, and a team supervisor. The male interviewer and the field editor worked as measurers as well. There were four regional coordinators who visited the teams on a continuous basis. The field staff was trained during a three-week period in July 1998. The main fieldwork for the TDHS-98, including initial interviews, call-backs and reinterviews, began in the first week of August 1998 and was completed in mid-November.

Questionnaires were returned to the Institute of Population Studies in Ankara for data processing. The office editing staff checked the questionnaires returned from the field. Those questions which had not been pre-coded (e.g., occupation) were coded by the office team. The data were then entered and edited using microcomputers and the ISSA (Integrated System for Survey Analysis) package. The office editing and data processing activities were started in the third week of August, two weeks after the beginning of fieldwork, and were completed by the end of November 1998. During the data entry process, almost hundred percent verification was done by double entry. The relatively few questionnaires that were not verified largely consisted of call-back questionnaires.

The results of the household and individual questionnaires are summarized in Table 1.1. Information is provided on the overall coverage of the sample, including household and individual response rates. In all, 9,970 households were selected for the TDHS-98. At the time of the survey, 8,596 households were considered occupied and, thus, available for interview. The main reasons the field teams were unable to interview some households was because some dwelling units that had been listed were

² Fieldwork for the TDHS-98 began in August with 12 teams. In October, four new teams were formed to complete the fieldwork in the remaining provinces. The teams were composed of staff from the original teams who had completed their assignments.

Table 1.1 Results of the household and individual interviews

Number of households, number of interviews, and response rates.
Turkey 1998

Result	Residence		Total
	Urban	Rural	
WOMEN			
Household interviews			
Dwellings sampled	6,989	2,981	9,970
Households found	5,938	2,658	8,596
Households interviewed	5,480	2,579	8,059
Household response rate	92.3	97.0	93.8
Individual interviews: women			
Eligible women	6,322	3,146	9,468
Eligible women interviewed	5,702	2,874	8,576
Eligible woman response rate	90.2	91.4	90.6
HUSBANDS			
Household interviews			
Dwellings sampled	3,496	1,487	4,983
Households found	2,988	1,333	4,321
Households interviewed	2,765	1,298	4,063
Household response rate	92.5	97.4	94.0
Individual interviews: husbands			
Eligible husbands	2,087	956	3,043
Eligible husbands interviewed	1,312	659	1,971
Eligible husband response rate	62.9	8.9	64.8

found to be vacant at the time of the interview or the household was away for the extended period. Of the 8,596 occupied households 94 percent (8,059 households) were successfully interviewed.

In the interviewed households, 9,468 women were identified as eligible for the individual interview, i.e. they were women at reproductive ages (15-49) who were present in the household on the night before the interview. Interviews were successfully completed with 8,576 of these women (91 percent). Among the eligible women not interviewed in the survey, the principal reason for non-response was the failure to find the women at home after repeated visits to the household.

In half of the selected households, husbands of currently married eligible women who were present in the household on the night before the interview or who usually lived in that particular household were eligible for the husband survey. A total of 4,983 households were selected for the husband interview. In the households interviewed (4,321 households), 3,043 husbands were identified as eligible for the individual interview. These husbands were present in the household on the night before the interview and they were currently married to women at reproductive ages. Of the 3,043 husbands designated as eligible for individual interview, 65 percent (1,971) were successfully interviewed.

A more complete description of the fieldwork, coverage of the sample, and data processing is presented in Appendix B.

CHAPTER 2

CHARACTERISTICS OF HOUSEHOLDS AND SURVEY RESPONDENTS

İsmet Koç and Attila Hancıoğlu

The main objective of this chapter is to describe the general characteristics of the sample population, including age, sex, household composition, and education. This information is not only useful in itself but it can also be used to evaluate the quality of the TDHS-98 data and to investigate changes in social and economic conditions over time. The other objective of this chapter is to describe the environment in which respondents and their children live. Household-level indicators relating to housing facilities and the ownership of durable goods are presented. These data are useful in assessing the household's living standard, which is an important influence on nuptiality, fertility and contraceptive behaviour, as well as maternal care and child mortality.

The TDHS household questionnaire included two questions distinguishing between the de jure population (persons who are usual residents in the selected households) and the de facto population (persons who spent the night before the interview in the selected households). The differences between these two populations are small and since past surveys and censuses were based on de facto populations, all tabulations in this report are based on the de facto population in the selected households, unless otherwise stated. A household was defined as a person or a group of persons living together and sharing a common source of food.

2.1 Population by Age and Sex

Table 2.1 and Figure 2.1 present the age distribution of the de facto population by five-year age groups according to urban-rural residence and sex. In addition to providing a background against which various demographic processes are occurring, the age structure of the population incorporates the past history of the population. As was observed in the 1993 Turkish Demographic and Health Survey (TDHS-93), the age distribution conforms to the pattern typical of countries that experienced relatively high fertility in the recent past. Evidence of recent declines in fertility is reflected in the fact that there is a smaller proportion of children under age 5 than age 5 to 9.

As Table 2.1 shows, the proportion under 15 is greater in the rural population than the urban population. The differences in the age distributions are evidence of lower recent fertility in urban areas compared with rural areas. Overall, the number of women significantly exceeds the number of men. This pattern is especially pronounced at ages 15-29, which may partly be explained by migration of young men for work and for military service.


Figure 2.2 presents the distribution of the male and female household population by single year of age. The data shows the evidence of a preference for reporting ages that end in zero or five that is common in countries where ages are not well known. Digit preference is considerably more pronounced for men than for women. This is probably due to the fact that many of the household interviews were completed with women who were able to provide more accurate information on their own ages than on men's ages.

Table 2.1. Household population by age, residence and sex


Percent distribution of the de facto household population by five-year age groups, according to urban-rural residence and sex, Turkey 1998

Age group	Urban			Rural			Total		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
0-4	10.6	9.6	10.1	11.4	9.5	10.4	10.9	9.5	10.2
5-9	10.5	10.1	10.3	11.7	10.2	10.9	10.9	10.1	10.5
10-14	10.4	9.7	10.0	12.1	12.0	12.0	11.1	10.5	10.8
15-19	11.0	10.6	10.8	10.2	11.6	10.9	10.7	11.0	10.8
20-24	8.4	10.6	9.5	7.5	8.5	8.0	8.1	9.8	9.0
25-29	9.0	9.5	9.3	6.9	6.9	6.9	8.2	8.6	8.4
30-34	7.9	8.0	8.0	5.8	5.9	5.9	7.1	7.2	7.2
35-39	7.2	7.1	7.2	5.1	5.9	5.5	6.5	6.7	6.6
40-44	6.4	6.0	6.2	5.1	5.0	5.1	5.9	5.6	5.8
45-49	5.2	4.7	4.9	4.3	4.2	4.3	4.9	4.5	4.7
50-54	3.7	3.6	3.6	4.0	4.6	4.3	3.8	3.9	3.9
55-59	2.8	3.2	3.0	4.1	4.3	4.2	3.3	3.6	3.4
60-64	2.5	2.3	2.4	4.0	3.5	3.7	3.1	2.7	2.9
65-69	2.0	2.2	2.1	3.6	3.4	3.5	2.5	2.6	2.6
70-74	1.3	1.4	1.3	2.3	2.3	2.3	1.7	1.7	1.7
75-79	0.7	0.7	0.7	1.2	1.0	1.1	0.9	0.8	0.8
80+	0.4	0.8	0.6	0.7	1.2	1.0	0.5	1.0	0.8
Don't know/Missing	0.1	0.0	0.1	0.1	0.1	0.1	0.1	0.0	0.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Number	10,533	10,897	21,430	5,930	6,443	12,374	16,464	17,340	33,804

**Figure 2.1
Population Pyramid, Turkey, 1998**



TDHS 1998



The distribution of the TDHS-93 sample population by broad age groups is presented in Table 2.2, along with comparable data from 1990 Census and TDHS-93. There has been a decline in the proportion of population less than 15 years of age and an increase in the proportion of ages 15-64 and 65 and over. Again this pattern is typical of populations that are experiencing a fertility decline.

Table 2.2 Population by age from selected sources

Percent distribution of the population by age group, selected sources, Turkey 1990-1998

Age group	CP 1990	TDHS 1993	TDHS 1998
Less than 15	35.0	33.0	31.5
15-64	60.7	61.4	62.6
65+	4.3	5.5	5.9
Total	100.0	100.0	100.0
Median age	22.2	23.1	24.3
Dependency ratio	64.7	62.7	59.7

Sources: 1990 Census of Population, SIS, 1993; 1993 TDHS, HIPS, 1994

The dependency ratio, defined as the ratio of the non-reproductive population (persons under age 15 and age 65 and over) to reproductive population (persons age 15-64), is calculated based on broad age distribution of the survey population. The dependency ratio decreased from 65 in 1990 to 60 in 1998, indicating a gradual lessening in the burden placed on persons in the productive ages to support older and younger household members. Table 2.2 also indicates that the median age of household population is 24.3

years, two years higher than the median age in 1990. Both changes in dependency ratio and in the median age of population are consistent with the gradual aging of the population that occurs as fertility declines.

2.2 Household Composition

Table 2.3 presents the distribution of households by the sex of the head of the household, size of the household and presence of foster children. The size and composition of the household may affect the allocation of financial resources among household members, which in turn influences the overall well being of the members. Household size may be associated with crowding in the dwelling, which can lead to unfavourable health conditions. In cases where women are heads of household, financial resources are often limited.

The TDHS-98 shows that 10 percent of households are headed by women. The proportion is slightly higher in urban areas than in rural areas (11 percent and 9 percent, respectively). Seven in ten households have between two and five members. The average household size is 4.3 persons, 4.0 persons in urban areas, and 4.9 persons in rural areas. Only one percent of households include one or more children under age 15 who are living with neither their natural father nor their natural mother.

Characteristic	Residence		Total
	Urban	Rural	
Household headship			
Male	88.9	90.9	89.6
Female	11.1	9.1	10.4
Number of usual members			
0	1.1	2.1	1.4
1	5.3	5.0	5.2
2	16.1	16.1	16.1
3	19.3	12.7	17.2
4	25.2	14.7	21.9
5	15.0	14.4	14.8
6	8.4	11.6	9.4
7	4.0	7.3	5.1
8	2.1	5.3	3.1
9+	3.6	10.8	5.9
Total	100.0	100.0	100.0
Mean size	4.0	4.9	4.3
Percent with foster children	1.2	1.6	1.3
Number of households	5,497	2,562	8,059

2.3 Fosterhood and Orphanhood

Information on the proportion of households including fostered or orphaned children was presented in Table 2.3. Table 2.4 considers information regarding fosterhood and orphanhood for the population of children under 15 years of age. Ninety-four percent of children under 15 years of age are living with both parents, four percent are living with their mothers (but not with their fathers), one percent are living with their fathers (but not with their mothers), and one percent are living with neither their

Table 2.4 Fosterhood and orphanhood

Percent distribution of de jure children under age 15 by survival of parents and child's living arrangements, according to child's age, sex, residence, and region, Turkey 1998

Background characteristic	Living with both parents	Living with mother but not father		Living with father but not mother		Not living with either parent			Missing information on father/mother	Total	Number of children	
		Father alive	Father dead	Mother alive	Mother dead	Both alive	Father only alive	Mother only alive				Both dead
Age												
0-2	97.0	2.0	0.3	0.1	0.1	0.4	0.0	0.0	0.0	0.1	100.0	2,045
3-5	96.1	1.8	0.8	0.3	0.3	0.5	0.0	0.1	0.0	0.1	100.0	1,976
6-9	94.0	2.0	2.0	0.6	0.5	0.6	0.3	0.1	0.0	0.0	100.0	2,871
10-14	90.0	2.2	3.7	0.7	1.5	1.2	0.2	0.4	0.1	0.0	100.0	3,693
Sex												
Male	93.7	2.1	1.9	0.4	0.7	0.7	0.1	0.2	0.1	0.0	100.0	5,386
Female	93.4	1.9	2.1	0.5	0.8	0.8	0.2	0.2	0.0	0.0	100.0	5,198
Residence												
Urban	94.1	2.1	1.6	0.5	0.7	0.7	0.1	0.1	0.1	0.1	100.0	6,608
Rural	92.7	1.9	2.8	0.5	0.8	0.8	0.2	0.3	0.0	0.0	100.0	3,977
Region												
West	94.5	2.1	0.7	0.7	1.0	0.6	0.1	0.2	0.1	0.1	100.0	3,202
South	93.1	1.8	2.5	0.5	0.4	1.4	0.1	0.2	0.1	0.0	100.0	1,574
Central	93.0	3.0	1.6	0.6	0.3	0.8	0.3	0.2	0.1	0.1	100.0	2,228
North	93.2	2.4	2.5	0.1	0.3	1.2	0.0	0.2	0.0	0.0	100.0	825
East	93.2	1.1	3.5	0.2	1.1	0.4	0.2	0.1	0.0	0.1	100.0	2,756
Total	93.6	2.0	2.0	0.5	0.7	0.8	0.2	0.2	0.1	0.0	100.0	10,585

Note: By convention, *foster children* are those who are not living with either biological parent. This includes *orphans*, i.e., children with both parents dead.

natural father nor natural mother. Of children under 15 years of age, two percent have lost their fathers and one percent have lost their mothers. Less than one percent of children have lost both their natural parents.

2.4 Educational Level of the Household Population

The education level of household members is perhaps their most important characteristic. Many phenomena, such as reproductive behaviour, use of contraception, health of children, and proper hygienic habits, are affected by the education of household members. In the TDHS-98, information on educational attainment was collected for every member of the household. Table 2.5 shows the distribution of the de facto female and male household populations age six and over by the highest level of education attended and the median number of years of schooling completed, according to selected background characteristics.

Primary education is compulsory in Turkey; it starts at age 6 or 7 and in accordance with the recent curriculum reform lasts eight years. Seventy-five percent of men and 60 percent of women have completed at least primary school, and 19 percent of men and 12 percent of women have completed secondary school or higher. Table 2.5 also shows the median number of years of schooling attained by males and females. Overall, males have a median duration of schooling of 4.7 years, 0.4 years longer than females. The gap in the median number of years of schooling between males and females is more than 1 year for the population above age 15, but is negligible among those age 10-14 years.

Table 2.5 also presents the educational level of household members by urban-rural residence and region. The proportion of persons with no education is much higher in rural areas than in urban areas, and this difference is observed for both males and females. More than three-fourths of males and nearly two-

Table 2.5 Educational level of the household population

Percent distribution of the de facto household population age six and over by highest level of education attended, according to selected background characteristics, Turkey 1998

Background characteristic	Level of education						Total	Number	Median number of years
	No education	Primary incomplete	Primary complete	Secondary incomplete	Secondary complete+	Don't know/ Missing			
MALE POPULATION									
Age									
6-9	27.1	70.5	0.7	0.2	0.0	1.5	100.0	1,466	0.6
10-14	3.0	30.1	25.1	41.3	0.4	0.2	100.0	1,820	4.6
15-19	1.2	2.8	26.0	45.5	24.3	0.1	100.0	1,757	7.8
20-24	1.5	1.2	32.9	21.8	42.4	0.1	100.0	1,327	7.7
25-29	2.1	1.4	41.6	22.4	32.5	0.1	100.0	1,355	5.8
30-34	3.5	1.8	44.7	21.2	28.6	0.3	100.0	1,175	5.0
35-39	3.9	2.0	48.9	16.3	28.8	0.2	100.0	1,064	4.9
40	4.8	2.0	48.8	17.3	27.0	0.0	100.0	978	4.9
45-49	9.4	4.0	50.2	13.8	22.6	0.0	100.0	801	4.7
50-54	20.3	7.0	42.8	14.0	15.1	0.7	100.0	620	4.5
55-59	26.3	8.9	45.5	6.0	12.0	1.4	100.0	536	4.3
60-64	30.7	10.2	45.8	5.9	5.9	1.5	100.0	504	4.2
65+	41.9	10.3	33.1	6.0	6.8	1.9	100.0	920	2.4
Residence									
Urban	7.5	13.1	30.1	23.8	24.9	0.5	100.0	9,214	4.9
Rural	16.4	15.3	41.5	16.8	9.3	0.6	100.0	5,124	4.4
Region									
West	6.7	11.2	35.2	23.9	22.4	0.5	100.0	5,337	4.9
South	9.1	15.4	39.1	20.2	15.7	0.5	100.0	2,103	4.6
Central	8.0	13.3	34.2	22.2	21.7	0.6	100.0	3,250	4.8
North	12.1	15.0	33.6	21.5	17.2	0.5	100.0	1,130	4.6
East	23.4	18.8	28.1	15.4	13.9	0.4	100.0	2,517	4.3
Total	10.7	13.9	34.2	21.3	19.4	0.5	100.0	14,338	4.7
FEMALE POPULATION									
Age									
6-9	29.8	67.9	1.1	0.1	0.0	1.0	100.0	1,441	0.6
10-14	7.3	24.4	36.1	31.7	0.3	0.2	100.0	1,823	4.5
15-19	6.3	4.3	43.4	27.7	18.2	0.1	100.0	1,907	4.9
20-24	10.2	2.6	50.1	9.2	27.7	0.2	100.0	1,701	4.7
25-29	13.6	2.7	51.3	10.7	21.6	0.1	100.0	1,483	4.7
30-34	15.9	3.6	54.2	8.3	17.9	0.0	100.0	1,256	4.6
35-39	22.1	6.3	49.5	6.3	15.6	0.2	100.0	1,154	4.4
40-44	30.3	9.4	39.8	7.1	13.4	0.0	100.0	975	4.3
45-49	39.5	8.6	36.5	5.4	9.6	0.5	100.0	787	4.0
50-54	50.0	10.0	31.5	2.6	4.5	1.4	100.0	684	0.0
55-59	56.6	12.1	25.8	1.7	3.7	0.0	100.0	619	0.0
60-64	61.3	12.4	20.1	2.1	3.7	0.4	100.0	471	0.0
65+	75.0	8.3	11.6	2.7	1.5	0.9	100.0	1,061	0.0
Don't know/Missing	100.0	0.0	0.0	0.0	0.0	0.0	100.0	8	0.0
Residence									
Urban	20.9	13.2	34.2	14.4	17.0	0.4	100.0	9,662	4.4
Rural	33.1	15.4	40.8	6.8	3.5	0.3	100.0	5,707	4.0
Region									
West	17.0	13.1	39.8	13.7	16.2	0.4	100.0	5,548	4.5
South	25.2	15.3	38.1	11.4	9.8	0.2	100.0	2,186	4.2
Central	20.8	13.5	39.8	12.3	13.3	0.4	100.0	3,524	4.4
North	28.0	13.7	36.7	11.4	9.8	0.5	100.0	1,241	4.2
East	46.4	15.8	25.7	6.8	4.9	0.3	100.0	2,870	0.6
Total	25.4	14.0	36.7	11.6	12.0	0.3	100.0	15,369	4.3

thirds of females in the urban areas are graduates of at least primary school. In rural areas, the proportions completing at least the primary level are lower for both males (68 percent) and females (51 percent). The proportion of secondary school graduates also differs between urban and rural areas for males and, in an even more pronounced way, for females. The proportion of secondary school graduates is around five times higher for females in urban areas than in rural areas.

Regional differences in education are considerable. The highest proportions of women (46 percent) and men (23 percent) with no education are found in the East region. The West region has the lowest proportions of male and female respondents with no education (7 and 17 percent, respectively).

2.5 School Enrolment

Table 2.6 presents information on school enrollment by age, sex, and residence. These rates are simple ratios of the number of enrolled persons in a specific age group to the total number in that age group. According to the TDHS, 72 percent of children age 6-10 were enrolled in school at the survey date. The proportion enrolled drops to 64 percent in the age group 11-15 years. Enrolment after age 15 drops significantly; whereas two in three children age 6-15 are in school, among the population age 16-20, the ratio drops to only one in four, and by age 21-24, only around one in ten are attending school. At all ages, the rural and/or female population has consistently lower school enrolment than the urban and/or male population (Figure 2.3). Moreover, as age increases, the gap between males and females widens. These results show that both gender and residential differences persist in the proportion of the population currently attending school in Turkey.


Table 2.6 School enrolment

Percentage of the de facto household population age 6-24 years enrolled in school, by age group, sex, and urban-rural residence, Turkey 1998

Age group	Male			Female			Total		
	Urban	Rural	Total	Urban	Rural	Total	Urban	Rural	Total
6-10	78.2	68.3	74.4	73.7	65.0	70.4	76.0	66.7	72.4
11-15	79.4	64.5	73.6	67.1	38.3	55.1	73.3	50.6	64.2
6-15	78.8	66.5	74.0	70.4	50.8	62.6	74.7	58.5	68.3
16-20	36.7	22.1	31.6	25.5	10.3	19.6	30.9	15.4	25.1
21-24	18.1	7.5	14.7	11.5	3.0	8.9	14.5	5.1	11.5

2.6 Housing Characteristics

To assess the economic and environmental conditions in which household members live, the household questionnaire included questions about sources of drinking water, time to the water source, sanitation facilities, flooring material, and the number of rooms used for sleeping. Information on these characteristics is useful from a public health point of view, as well as indirectly reflecting the household socioeconomic status.



This information on household characteristics is given in Table 2.7. Overall, more than 5 in 10 households get their drinking water from pipes, mainly within their dwelling. The source for drinking water differs considerably by area of residence. Among urban households, 67 percent get drinking water from pipes in their residence, and only one percent obtains water from a public tap. The second most common source of drinking water in urban areas is bottled water (15 percent), and nine percent get drinking water from a water station (commercial outlets selling spring water). Around two-thirds of rural households report having piped water; however, for nearly half of these households, the source for the piped water is a river, stream or other surface water. Nearly one-fifth of rural households obtain water from a spring. Households in rural Turkey are somewhat more likely to be relying on well water than households in urban Turkey (9 percent and 2 percent respectively).

For most households, the source for drinking water is within their dwelling or not far from the residence. Overall, 91 percent of households obtain drinking water in their dwelling or from within 15 minutes of the residence. Urban households tend to be somewhat closer to the source of water than rural areas; 93 percent of households in urban areas report that the source for drinking water is in the house or within 15 minutes of dwelling compared with 88 percent of rural households.

Two-thirds of households have modern sanitation facilities in Turkey. Another 29 percent use an open pit or latrine, and only two percent report having no toilet facilities. There are substantial differences in the type of toilet facilities available to households by urban-rural residence. Households in urban areas are much more likely to have modern sanitation facilities than households in rural areas (89 percent and 25 percent, respectively). Only ten percent of urban households use pit toilets or report that they have no toilet facilities. In comparison, 70 percent of rural households use a pit or latrine, and five percent report that they have no toilet facilities.

Table 2.7. Housing characteristics			
Percent distribution of households by housing characteristics, according to urban-rural residence, Turkey 1998			
Housing characteristic	Residence		Total
	Urban	Rural	
Source of drinking water			
Piped into residence/garden	65.9	32.7	55.3
Public tap	0.8	1.3	1.0
Well in residence/garden	1.6	5.6	2.8
Public well	0.3	3.0	1.2
Piped surface water			
in house/garden	1.4	34.7	12.0
Spring/public fountain	4.7	19.0	9.2
River/stream	0.2	1.0	0.5
Rainwater	0.0	0.1	0.0
Tanker truck	0.6	0.8	0.7
Bottled water/Demi John	14.9	1.0	10.5
Water station	9.1	0.2	6.3
Other	0.4	0.5	0.5
Total	100.0	100.0	100.0
Time to water source			
Water within 15 min./on premises	92.6	87.6	91.0
Type of toilet facility			
Own flush toilet	87.6	24.0	67.4
Shared flush toilet	1.6	0.7	1.3
Open pit toilet	2.4	25.9	9.9
Closed pit latrine	7.5	43.6	19.0
No facility/bush	0.3	5.0	1.8
Other	0.3	0.7	0.5
Missing	0.3	0.1	0.2
Total	100.0	100.0	100.0
Main floor material			
Earth	2.4	16.7	6.9
Wood planks	12.9	24.4	16.5
Parquet/polished wood	14.5	2.0	10.5
Karo	11.6	3.7	9.1
Cement	23.3	41.8	29.2
Carpet	10.1	6.3	8.9
Marley	19.3	3.5	14.2
Mosaic	4.9	1.0	3.7
Other	0.8	0.6	0.7
Missing	0.3	0.2	0.3
Total	100.0	100.0	100.0
Persons per sleeping room			
1-2	78.6	64.9	74.3
3-4	18.2	27.4	21.1
5-6	2.4	5.1	3.3
7+	0.7	2.4	1.3
Don't know/Missing	0.1	0.1	0.1
Total	100.0	100.0	100.0
Mean persons per room	2.2	2.6	2.3
Number of households	5,497	2,562	8,059

With regard to flooring, 29 percent of the households live in dwellings with cement floors, an additional 27 percent have a wooden floor, and 14 percent have marley floors. There are significant differences in the flooring materials in urban and rural dwellings. Cement is the most common flooring material in both rural areas (42 percent) and urban areas (23 percent). Wood is also a common flooring material; about one-fourth of urban and rural households live in dwellings with wood floors. Seventeen percent of households in rural areas have earth floors, compared to only two percent of households in urban areas.

Information on the number of rooms that a household uses for sleeping was collected to determine the extent of crowding. Table 2.7 shows that 74 percent of households have one or two persons per sleeping room, and 21 percent have three to four persons per sleeping room. The overall mean is 2.3 persons per sleeping room. Rural households are more crowded than urban households. The mean number of persons per sleeping room is 2.2 persons in urban areas compared with 2.6 persons in rural areas.

2.7 Household Durable Goods

The availability of durable consumer goods is a good indicator of household socioeconomic level. Moreover, particular goods have specific benefits. For example, having access to a television exposes household members to innovative ideas, and a refrigerator prolongs the wholesomeness of foods. Table 2.8 presents the availability of selected consumer goods by residence. Most of the population in Turkey enjoys the convenience of electrical appliances. More than nine in ten households own a television set or a refrigerator, while almost eight in ten households have a telephone, and more than six in ten households own an oven, a vacuum cleaner or a washing machine. Ownership of various durable goods varies by place of residence, with higher proportions of ownership for all items reported among households in urban areas than in rural areas.

Durable goods	Residence		Total
	Urban	Rural	
Refrigerator	96.2	86.7	93.2
Gas or electric oven	77.0	47.5	67.6
Dishwasher	22.4	3.3	16.4
Washing machine	76.1	36.9	63.6
Vacuum cleaner	77.7	38.0	65.1
Television	94.8	84.1	91.4
Video recorder	14.3	3.7	10.9
Camera	40.7	18.9	33.8
CD player	19.8	5.0	15.1
Telephone	80.4	69.7	77.0
Mobile telephone	17.9	3.5	13.3
Computer	6.0	0.4	4.2
None of the above	1.5	4.4	2.4
Number of households	5,497	2,562	8,059

2.8 Background Characteristics

A description of the basic characteristics of the women and husbands interviewed in the TDHS is essential as background for interpreting findings presented later in the report. Table 2.9 provides the percent distribution of women and husbands by age, marital status, level of education, urban-rural residence, and region.

Background characteristic	Weighted percent	Number of women		Weighted percent	Number of husbands	
		Weighted	Un-weighted		Weighted	Un-weighted
Age						
15-19	20.1	1,720	1,763	*	*	3
20-24	18.2	1,558	1,539	5.5	109	105
25-29	16.3	1,397	1,373	17.4	342	314
30-34	14.0	1,202	1,195	18.5	364	362
35-39	12.6	1,081	1,104	17.9	352	366
40-44	10.3	885	890	17.0	335	344
45-49	8.5	733	712	12.2	240	245
50+	NA	NA	NA	11.4	226	232
Current marital status						
Never married	27.7	2,380	2,424	NA	NA	NA
Married	69.0	5,921	5,893	100.0	1,971	1,971
Widowed	1.7	147	153	NA	NA	NA
Divorced	1.1	96	77	NA	NA	NA
Separated	0.4	34	29	NA	NA	NA
Residence						
Urban	66.5	5,704	5,702	68.3	1,347	1,312
Rural	33.5	2,872	2,874	31.7	624	659
Region						
West	37.4	3,204	2,170	38.9	767	546
South	14.7	1,258	1,678	14.4	285	400
Central	23.1	1,985	1,706	24.4	481	413
North	8.1	692	1,258	7.6	150	299
East	16.8	1,437	1,764	14.6	287	313
Education						
No education	16.7	1,435	1,590	6.6	131	135
Primary incomplete	5.0	426	442	3.7	73	76
Primary complete	48.0	4,117	4,013	42.8	844	849
Secondary incomplete	12.1	1,041	1,029	20.8	409	401
Secondary complete+	18.1	1,556	1,502	26.0	513	510
Total	100.0	8,576	8,576	100.0	1,971	1,971

NA = Not applicable
 Note: An asterisk indicates that an item is based on fewer than 25 persons and has been suppressed.

Women and husbands were asked two questions in the individual interview to assess their age: "In what month and year were you born?" and "How old are you?" Interviewers were trained to probe in situations in which respondents knew neither their age nor date of birth; as a last resort, interviewers were instructed to record their best estimate of the respondent's age. The age data indicate that 38 percent of women and six percent of husbands are under age 25, 30 percent of women and 36 percent of husbands

are in the age group 25-34 and slightly more than 30 percent of women and 47 percent of husbands are in the age group 35-49. Around 10 percent of husbands are age 50 or older.

Data on the women's marital status at the time of survey indicate that 69 percent were currently married, while 28 percent had never married, and the rest were widowed, divorced, or separated. The latter proportion indicates the rarity of marital dissolution in Turkey. As described earlier, the men's sample included husbands of eligible currently married women living in a subsample of the households covered in the TDHS. Thus, the men's sample does not include never-married men or men who were widowed, divorced or separated.

About two-thirds of women and men live in urban areas, and the rest live in rural areas. According to the data, more than one-third of the respondents live in the West region, one-fourth live in the Central region, around one-sixth live in the East and in the South, and less than 10 percent live in the North region.

The proportion of women who have never attended school is more than twice that of husbands (17 percent and 7 percent, respectively). About 48 percent of women and 43 percent of husbands have a primary education only, while 30 percent of women and 47 percent of husbands have gone beyond primary school education. Fewer than one in five women have completed secondary school or higher.

2.9 Respondents' Level of Education by Background Characteristics

Table 2.10 shows the distributions of women and husbands by the highest level of education attended, according to selected characteristics. The table is shown as a first effort to clarify the relationship between education and other explanatory or background variables used in later tabulations. Differences in the educational composition of respondents from different age groups, regions, and urban-rural backgrounds are highlighted.

As mentioned before, men are generally better educated than women. For both groups, education is inversely related to age; older women and men are generally less educated than younger women and men. The percentage of women with no education rises rapidly with age, from six percent in the 15-19 age group to 40 percent in the age group 45-49. This suggests that younger women have had better educational opportunities than older women. This is also reflected in the higher percentage of women in the age group 20-24 who completed secondary education (27 percent), compared to women age 45-49 (9 percent).

Urban women and men in Turkey are much more likely to have higher education than their rural counterparts. Twenty-two percent of rural women have no education, compared to only 14 percent of urban women. Conversely, while 39 percent of urban women have gone to school beyond the primary level, while only 14 percent of rural women have been educated beyond primary school. Similar urban-rural differentials in education are observed for husbands. A comparison of the results for regions shows that East region has the highest proportions of women and husbands with no education (40 percent and 21 percent respectively), while the West region has the lowest proportion of women and husbands with no education (10 percent and 4 percent, respectively). The highest proportions of women and husbands with secondary or more education is also observed in the West region.

Table 2.10 Level of education							
Percent distribution of women and husbands by the highest level of education attended, according to selected background characteristics, Turkey 1998							
Background characteristic	Level of education					Total	Number of women/husbands
	No education	Primary incomplete	Primary complete	Secondary incomplete	Secondary complete+		
WOMEN							
Age							
15-19	5.8	4.8	44.4	27.7	17.3	100.0	1,720
20-24	9.4	2.4	51.3	9.9	26.9	100.0	1,558
25-29	13.3	2.5	54.1	10.4	19.7	100.0	1,397
30-34	15.5	3.8	54.1	8.2	18.4	100.0	1,202
35-39	22.9	6.6	49.2	6.5	14.9	100.0	1,081
40-44	31.6	9.4	39.7	6.4	12.9	100.0	885
45-49	39.5	9.5	36.2	5.6	9.3	100.0	733
Residence							
Urban	14.1	3.9	43.3	14.5	24.2	100.0	5,704
Rural	21.9	7.1	57.3	7.5	6.2	100.0	2,872
Region							
West	9.9	3.2	49.8	14.3	22.8	100.0	3,204
South	18.2	5.9	49.8	11.4	14.6	100.0	1,258
Central	9.8	5.4	52.4	12.1	20.3	100.0	1,985
North	18.3	4.8	49.1	11.6	16.2	100.0	692
East	39.5	7.6	35.9	8.2	8.8	100.0	1,437
Total	16.7	5.0	48.0	12.1	18.1	100.0	8,576
HUSBANDS							
Age							
15-24	0.0	3.0	42.9	27.7	26.4	100.0	113
25-29	1.8	2.8	38.3	25.3	31.8	100.0	342
30-34	3.9	3.4	38.8	24.2	29.8	100.0	364
35-39	3.9	2.7	45.9	17.9	29.5	100.0	352
40-44	5.9	2.2	47.9	17.7	26.2	100.0	335
45-49	8.5	4.0	47.7	19.3	20.5	100.0	240
50+	25.1	9.6	38.4	15.4	11.5	100.0	226
Residence							
Urban	4.7	2.6	36.4	22.5	33.7	100.0	1,347
Rural	10.8	6.2	56.6	17.0	9.5	100.0	624
Region							
West	3.6	3.2	41.0	22.0	30.3	100.0	767
South	6.4	6.1	49.1	20.7	17.7	100.0	285
Central	4.0	2.2	42.6	23.7	27.5	100.0	481
North	5.1	3.5	50.6	16.0	24.7	100.0	150
East	20.5	5.3	37.8	15.2	21.2	100.0	287
Total	6.6	3.7	42.8	20.8	26.0	100.0	1,971

2.10 Reasons for Leaving School

Information about the reasons that lead women to drop out of school can provide guidance to programs seeking to improve educational opportunities for women. To obtain some insight into this issue, women under age 25 years who were not currently attending school were asked during the TDHS interview about the main reason for leaving school. Table 2.11 shows the distribution of women 15-24 years according to whether they are currently attending school and, if not, their reasons for leaving school, according to the highest level of education attended.

Table 2.11 Reasons for leaving school

Percent distribution of women age 15-24 who had ever attended school but were not currently attending by reason for leaving school, according to highest level of education attended, Turkey 1998

Reason stopped attending school	Highest level of education					Total
	Primary incomplete	Primary complete	Secondary incomplete	Secondary complete	Higher	
URBAN						
Currently attending	3.6	0.6	52.2	16.6	67.5	23.7
Got pregnant	0.0	0.0	0.0	0.1	0.0	0.0
Got married	0.0	2.3	4.1	9.8	3.9	4.3
Take care of children	0.0	0.1	0.9	0.0	0.0	0.3
Family needed help	3.2	1.2	0.6	0.3	0.0	0.8
Could not pay school fees	4.3	7.4	1.4	1.8	0.0	3.9
Needed to earn money	1.9	2.1	0.4	2.4	1.0	1.6
Graduated/had enough school	0.0	0.6	0.1	6.5	21.3	3.9
Did not pass exams	2.4	0.0	0.3	35.8	0.0	7.2
Did not like school	16.3	23.9	17.6	5.4	0.3	16.0
School not accessible	1.3	3.7	0.4	0.2	0.0	1.7
Parents did not send to school	30.5	36.2	9.1	4.9	1.4	19.5
Other	36.5	19.1	11.8	15.4	3.7	15.4
Don't know/Missing	0.0	2.8	1.0	0.7	1.0	1.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Number	55	829	482	386	210	1,963
RURAL						
Currently attending	1.7	0.5	45.6	16.7	(57.1)	9.7
Got pregnant	0.0	0.0	0.0	0.0	0.0	0.0
Got married	1.6	2.0	4.7	11.7	(3.0)	3.2
Take care of children	1.5	0.1	0.0	0.0	0.0	0.2
Family needed help	7.3	2.0	0.0	1.5	0.0	2.0
Could not pay school fees	1.5	3.6	1.6	0.0	0.0	2.8
Needed to earn money	1.0	0.6	0.4	2.7	0.0	0.8
Graduated/had enough school	0.0	0.7	0.0	3.3	(31.0)	1.6
Did not pass exams	0.0	0.2	0.0	36.1	0.0	3.2
Did not like school	6.2	14.4	13.4	7.3	0.0	12.8
School not accessible	4.3	7.5	6.4	0.0	0.0	6.3
Parents did not send to school	40.4	44.6	16.5	3.5	0.0	35.8
Other	31.1	22.0	8.8	17.1	(8.9)	19.9
Don't know/Missing	3.4	1.7	2.6	0.0	0.0	1.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Number	66	734	148	91	29	1,068
TOTAL						
Currently attending	2.6	0.6	50.6	16.7	66.2	18.8
Got pregnant	0.0	0.0	0.0	0.1	0.0	0.0
Got married	0.9	2.1	4.3	10.1	3.8	3.9
Take care of children	0.8	0.1	0.7	0.0	0.0	0.2
Family needed help	5.4	1.6	0.5	0.6	0.0	1.2
Could not pay school fees	2.7	5.6	1.4	1.4	0.0	3.5
Needed to earn money	1.4	1.4	0.4	2.4	0.9	1.3
Graduated/had enough school	0.0	0.7	0.1	5.9	22.5	3.1
Did not pass exams	1.1	0.1	0.2	35.9	0.0	5.8
Did not like school	10.8	19.4	16.6	5.8	0.3	14.8
School not accessible	3.0	5.5	1.8	0.1	0.0	3.4
Parents did not send to school	35.9	40.1	10.8	4.7	1.2	25.2
Other	33.5	20.4	11.1	15.7	4.3	17.0
Don't know/Missing	1.9	2.3	1.4	0.6	0.9	1.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Number	121	1,563	630	478	239	3,031

Note: Parentheses indicate that a figure is based on 25-49 respondents.

Only 19 percent of TDHS respondents 15-24 were attending school at the time of interview. The reasons for leaving school vary with the level of school women had attained at the time they left school. Women who had not completed primary school are most likely to say that they had left because parents did not send them (36 percent) or because they had not liked school (11 percent). Failure to pass examinations (36 percent) is the principal reason for dropping out of school among women who left after completing the secondary level, while eight percent report that they stopped going to school because they married.

2.11 Differentials in Characteristics of Couples

Because the married men interviewed in TDHS were selected from a subgroup of households it is possible to match male respondents with their wives to obtain a couple dataset. Table 2.12 shows husband-wife differentials in age and education for the 1,896 couples interviewed in the TDHS. For most of the couples, the husband is older than the wife; in the case of the majority of couples, the age difference is between 0 and 9 years. Among 10 percent of couples, the wife is older than her husband. On average, men are five years older than their wives.

Regarding educational differences, in about five percent of couples, neither the husband nor the wife has been to school, while among eight in ten couples, both the husband and wife are educated. For the remaining couples, the proportion in which the husband has some education and the wife has none is much higher than that of the wife is educated and the husband is not (14 percent and 2 percent, respectively).

<u>Table 2.12 Differential characteristics between spouses</u>	
Percent distribution of couples by differences in age and level of education between the husband and wife, Turkey 1998	
Characteristic	Percent/ years
Age difference	
Husband younger than wife	9.8
Husband older by:	
0-4 years	44.7
5-9 years	32.9
10-14 years	9.6
15 years+	2.9
Mean age difference	5.0
Education	
Both husband and wife not educated	4.9
Wife educated, husband not	1.7
Husband educated, wife not	13.8
Both husband and wife educated	79.6
Total	100.0
Number of couples	1,896

2.12 Exposure to Print Media

Women and husbands were asked how often they read newspapers or magazines. This information is important to program planners seeking to reach people with family planning and health messages through the media. Table 2.13 presents the percentage of women and husbands exposed to printed media by age, residence, region and level of education. Results show that 32 percent of women and 61 percent of husbands read newspapers or magazines weekly. Access to media is somewhat higher among younger respondents and among those living in urban as opposed to rural areas. Print media access is more widespread among women and husbands living in the West compared with among those living in the other regions. As expected, educated persons are more likely to read newspaper or magazines than less educated persons.

Table 2.13 Exposure to print media				
Percentage of women/husbands who usually read a newspaper at least once a week, by selected background characteristics, Turkey 1998				
Background characteristic	Women		Husbands	
	Read newspaper weekly	Number of women	Read newspaper weekly	Number of husbands
Age				
15-19	36.6	1,720	*	4
20-24	39.1	1,558	56.1	109
25-29	33.1	1,397	66.7	342
30-34	29.8	1,202	66.8	364
35-39	27.8	1,081	64.5	352
40-44	26.2	885	59.5	335
45-49	25.4	733	56.9	240
50-54	NA	NA	43.6	226
Residence				
Urban	40.9	5,704	68.2	1,347
Rural	15.4	2,872	44.1	624
Region				
West	45.1	3,204	74.3	767
South	23.9	1,258	48.0	285
Central	30.3	1,985	57.7	481
North	28.5	692	64.5	150
East	16.3	1,437	39.2	287
Education				
No education	1.9	1,435	12.7	131
Primary incomplete	9.2	426	33.2	73
Primary complete	24.2	4,117	52.2	844
Secondary incomplete	53.2	1,041	67.4	409
Secondary complete+	74.6	1,556	85.1	513
Total	32.4	8,576	60.6	1,971
NA = Not applicable				
Note: An asterisk indicates that an item is based on fewer than 25 persons and has been suppressed.				

2.13 Employment and Occupation

In the TDHS-98, information was collected about current employment. Although data were collected from both women and husbands, the analysis in this section concentrates on women. Table 2.14 indicates that 39 percent of women report being employed during the 12-month period before the TDHS interview with the majority (35 percent) working at the time of the survey. Nearly half of the women who were currently employed at the time of the survey worked all year, 43 percent worked in seasonal jobs, and only 11 percent worked occasionally. The proportion not currently employed is higher among older women, rural women, and women with no education than other women. Women in the West and Central regions are more likely to be employed than women in other regions. The North region has the highest proportion of women reporting seasonal employment. Rural women are much more likely to be employed seasonally than urban women, presumably as agricultural laborers.

Table 2.14 Employment
Percent distribution of women by employment status and continuity of employment, according to selected background characteristics, Turkey 1998

Background characteristic	Not currently employed		Currently employed					Total	Number of women
	Did not work in last 12 months	Worked in last 12 months	All year		Seasonally	Occasionally	Missing		
			5+ days per week	<5 days per week					
Age									
15-19	60.3	6.1	12.2	1.0	17.5	2.5	0.4	100.0	1,720
20-24	59.4	5.4	17.4	0.6	13.1	3.8	0.3	100.0	1,558
25-29	62.5	4.0	16.5	0.9	11.8	4.0	0.3	100.0	1,397
30-34	60.1	3.4	15.1	1.4	14.9	5.1	0.0	100.0	1,202
35-39	57.2	2.3	18.5	2.1	15.0	4.9	0.0	100.0	1,081
40-44	60.6	2.2	14.5	2.0	16.5	3.9	0.2	100.0	885
45-49	66.1	2.0	9.6	0.9	17.6	3.7	0.0	100.0	733
Residence									
Urban	68.3	4.4	17.7	1.2	4.8	3.3	0.3	100.0	5,704
Rural	45.4	3.2	9.8	1.2	35.2	5.1	0.0	100.0	2,872
Region									
West	57.7	5.5	22.6	1.4	8.7	3.5	0.4	100.0	3,204
South	65.1	3.2	10.7	1.2	16.2	3.6	0.1	100.0	1,258
Central	57.5	3.5	12.1	1.2	21.0	4.5	0.1	100.0	1,985
North	44.1	3.6	14.6	1.8	32.0	3.9	0.1	100.0	692
East	75.4	2.2	6.3	0.6	11.5	4.0	0.0	100.0	1,437
Education									
No education	68.7	1.9	6.7	0.9	18.0	3.9	0.0	100.0	1,435
Primary incomplete	63.2	1.9	11.2	1.0	19.4	3.3	0.0	100.0	426
Primary complete	59.1	3.8	11.1	1.0	19.8	4.9	0.2	100.0	4,117
Secondary incomplete	70.2	5.7	10.7	1.0	7.7	4.1	0.6	100.0	1,041
Secondary complete+	50.0	5.9	37.2	2.3	3.3	1.3	0.1	100.0	1,556
Total	60.6	4.0	15.1	1.2	15.0	3.9	0.2	100.0	8,576

Table 2.15 presents the distribution of women who were employed at the time of the survey by occupation, according to selected background characteristics. Forty-nine percent of employed women worked in agriculture, among whom more than six in ten worked on their family land. The table also shows that women who worked outside the agricultural sector were most often employed in professional and technical occupations, followed by skilled manual laborers. Most of the remaining women worked in household and domestic occupations and sales and services.

Table 2.15 Occupation

Percent distribution of currently employed women by occupation and type of agricultural land worked or type of non-agricultural employment, according to selected background characteristics, Turkey 1998

Background characteristic	Agricultural				Nonagricultural						Total	Number of women
	Own land	Family land	Rented land	Other's land	Prof./tech./manag.	Sales/services	Skilled manual	Unskilled manual	Household and domestic	Missing		
Age												
15-19	2.2	38.2	3.8	13.7	10.7	4.5	18.9	4.5	3.1	0.5	100.0	573
20-24	1.3	28.0	3.3	8.5	25.1	6.6	23.2	1.7	2.2	0.2	100.0	545
25-29	2.2	28.4	3.4	8.0	29.1	2.7	21.1	2.3	2.3	0.5	100.0	464
30-34	3.1	31.7	0.8	11.3	23.2	4.0	19.1	1.7	4.1	0.8	100.0	439
35-39	4.9	28.5	3.0	10.7	21.5	3.0	17.5	2.0	8.7	0.3	100.0	438
40-44	6.2	30.0	3.1	15.6	19.5	4.5	10.7	3.1	7.0	0.2	100.0	327
45-49	11.8	37.1	1.5	14.6	8.5	3.8	13.9	1.1	7.4	0.3	100.0	233
Residence												
Urban	1.3	7.2	1.1	7.4	36.2	7.4	26.6	4.2	8.2	0.6	100.0	1,544
Rural	6.2	57.0	4.7	15.6	3.6	1.0	10.2	0.7	0.7	0.3	100.0	1,475
Region												
West	2.4	18.0	2.2	5.2	26.7	6.4	26.9	4.1	7.7	0.4	100.0	1,164
South	3.6	25.0	1.0	28.8	18.0	5.7	10.1	1.4	4.9	1.5	100.0	399
Central	3.4	40.5	3.2	11.1	19.4	2.0	16.3	2.2	1.7	0.2	100.0	773
North	6.4	62.6	1.3	4.0	9.9	2.7	8.5	1.2	3.3	0.0	100.0	362
East	6.4	31.9	8.2	21.1	13.7	2.0	16.0	0.0	0.7	0.2	100.0	322
Education												
No education	9.1	39.5	4.2	24.0	1.6	1.6	10.5	1.8	7.6	0.1	100.0	422
Primary incomplete	7.0	38.3	3.2	17.7	7.1	3.2	12.9	1.9	8.1	0.4	100.0	149
Primary complete	3.8	43.3	3.9	12.6	2.4	3.8	23.3	2.0	4.5	0.3	100.0	1,516
Secondary incomplete	1.5	21.1	1.1	6.5	23.1	5.6	34.8	2.5	2.4	1.4	100.0	246
Secondary complete+	0.2	2.9	0.1	1.3	73.2	6.5	8.5	4.0	2.7	0.5	100.0	686
Total	3.7	31.5	2.8	11.4	20.3	4.2	18.6	2.5	4.5	0.4	100.0	3,019

Note: *Professional/technical/managerial* includes professional, technical, clerical and managerial occupations.

Women's occupations vary by age. The proportion of women who worked in agriculture is highest for those in the youngest and oldest age groups. The proportion declines from 58 percent of working women age 15-19 to 47 percent of women age 30-34, and it increases to 65 percent for women age 45-49. On the other hand, the proportion of women employed as professional and technical workers, has an opposite pattern: low at younger ages, peaking at age 25-29, and then declining for older women. Manual occupations attract younger women.

Table 2.15 also shows that women's occupations vary significantly by urban-rural residence and region. While 84 percent of working women in rural areas were engaged in agricultural sector, the corresponding proportion in urban areas was only 17 percent. In contrast, urban women were more likely to be employed in professional, technical and managerial occupations (36 percent), as skilled manual labors (27 percent), in household and domestic jobs (8 percent), or in sales and services (7 percent). An examination of regional differences shows that non-agricultural employment is more common in the West than in other regions; 72 percent of the employed women in the West work in the non-agricultural sector, with the vast majority working in professional, technical and managerial occupations (27 percent) and unskilled manual jobs (27 percent). As expected, the majority of employed women in the other regions

work in the agricultural sector; almost six in ten women in the South and Central regions and around seven in ten women in the North and East regions are engaged in agricultural jobs.

Women's education is inversely related to their propensity to work in agriculture; employed women with no education were much more likely to have been working in agriculture than better-educated women. For example, 77 percent of working women with no education worked in the agricultural sector, compared with only five percent of women who have completed secondary school. Moreover, women who had completed the secondary level were more likely to be employed in professional and technical occupations or in skilled manual occupations.

2.14 Decision on Use of Earnings

When assessing the status of women, one valuable indicator is their independence in making decisions on the use of their earnings. Table 2.16 shows that around half (49 percent) of employed women make their own decisions about the use of their earnings, while 29 percent decide jointly with their husband, and 15 percent are not involved in making the decisions. Independent decision-making with regard to the use of earnings tend to be higher among younger women, women in urban areas and among single and formerly married women.

There also is variation across regions in the percentages of women who indicate that they alone make decisions about the money they earn. The percentage of women who make their own decisions ranges from 43 percent in the South to 51 percent in the West. The percentage of women who report making the decision together with their husbands varies from a low of 18 percent (East) to a high of 33 percent (North). Twelve percent in the East report that husbands alone decide on how to spend their earnings; in the remaining regions, the percentage citing the husband as the prime decision-maker fall below 10 percent.

2.15 Child Care While Working

The welfare of children under six years of age whose mothers are employed is the focus of Table 2.17. Overall, one in four women who worked in the 12 months prior to the survey have one or more children under age six. This proportion varies by residence, education, region, and occupation. Less educated women, women working in the agricultural sector, women working occasionally, and women living in the rural areas are more likely to have children under six. Reflecting regional fertility differentials, women living in the East are more likely to have children under age six than women living in other regions.

Among working women with children under age six, 34 percent take care of their children while they work. Relatives and older female siblings are the other most common caretakers for children of working women (36 and 11 percent, respectively). The role of female siblings in child care is significant in families where the mother has limited education, works in agriculture or as seasonal worker, lives in the East, or lives in the rural areas. Children whose mothers have attended secondary school, live in urban areas or the West region, are employed all year, or work in non-agricultural occupations are more likely to be cared for by servants or hired help. Across all sub-groups, husbands and male siblings have a very limited role in child minding while the mother is at work.

Table 2.16 Decision on use of women's earnings

Percent distribution of women receiving cash earnings by person who decides how earnings are used, according to selected background characteristics, Turkey 1998

Background characteristic	Person who decides how earnings are used						Total	Number of women
	Self only	Husband/partner	Jointly with husband/partner	Someone else	Jointly with someone else	Missing		
Age								
15-19	47.8	1.9	1.3	32.2	15.7	1.0	100.0	306
20-24	60.4	2.0	16.8	11.4	9.4	0.0	100.0	347
25-29	51.1	4.8	37.0	2.6	4.3	0.3	100.0	291
30-34	44.8	9.3	38.6	3.7	3.6	0.0	100.0	259
35-39	38.7	10.6	49.6	0.0	1.0	0.0	100.0	271
40-44	51.5	12.5	33.9	0.0	1.7	0.4	100.0	180
45-49	44.6	13.2	40.1	0.0	2.1	0.0	100.0	109
Residence								
Urban	54.5	3.4	29.1	6.4	6.3	0.2	100.0	1,316
Rural	33.3	16.1	28.2	15.8	6.1	0.4	100.0	447
Region								
West	50.8	6.0	30.9	5.4	6.7	0.2	100.0	877
South	43.3	5.6	28.8	13.4	8.5	0.3	100.0	242
Central	50.2	7.0	27.5	10.0	5.0	0.3	100.0	379
North	49.1	5.3	33.1	7.8	4.6	0.0	100.0	114
East	46.9	11.7	17.7	18.8	4.5	0.5	100.0	151
Education								
No education	41.9	18.2	25.3	11.0	3.2	0.4	100.0	185
Primary incomplete	46.3	12.4	19.6	16.7	2.1	2.9	100.0	71
Primary complete	42.4	8.7	27.9	13.5	7.3	0.1	100.0	703
Secondary incomplete	62.1	0.9	15.9	8.1	12.5	0.6	100.0	166
Secondary complete+	55.6	1.8	35.4	2.3	4.8	0.0	100.0	639
Marital status								
Not married	67.9	0.0	0.0	18.2	13.5	0.4	100.0	778
Currently married	34.3	11.9	51.7	1.4	0.5	0.1	100.0	985
Total	49.2	6.6	28.9	8.8	6.3	0.3	100.0	1,763

Table 2.17 Child care while working

Percent distribution of currently employed women by whether they have a child under six years of age at home, and the percent distribution of employed mothers who have a child under six by person who cares for child while mother is at work, according to selected background characteristics, Turkey 1998

Background characteristic	No child under six at home	One or more children under six at home	Child's caretaker while mother is at work									Not worked since birth ¹	Other	Missing	Total	Number of employed women
			Re-spondent	Husband/partner	Other relative	Neigh-bor/Friend	Hired help	Child is in school	Other female child	Other male child						
Residence																
Urban	78.1	21.9	36.5	1.9	26.6	0.8	9.3	13.6	6.6	0.4	1.2	1.4	1.8	100.0	1,544	
Rural	70.7	29.3	32.0	0.1	44.1	0.5	0.5	0.4	13.7	1.6	1.1	3.0	3.0	100.0	1,475	
Education																
No education	72.9	27.1	36.5	1.2	19.3	0.0	0.0	0.0	28.2	1.6	1.8	7.2	4.3	100.0	422	
Primary incomplete	80.0	20.0	27.7	0.0	40.5	2.2	3.5	0.0	18.1	3.3	0.0	4.7	0.0	100.0	149	
Primary complete	71.9	28.1	39.3	0.7	42.9	0.5	0.5	0.3	9.0	1.1	0.7	1.9	3.1	100.0	1,516	
Secondary incomplete	78.7	21.3	45.5	0.0	39.7	0.0	0.0	6.7	6.1	0.0	1.3	0.0	0.7	100.0	246	
Secondary complete+	78.3	21.7	13.8	1.8	29.1	1.3	20.7	28.8	1.7	0.4	2.1	0.0	0.5	100.0	686	
Region																
West	79.9	20.1	31.5	0.7	34.6	0.6	6.4	9.8	9.1	0.1	1.2	2.2	3.7	100.0	1,164	
South	80.0	20.0	32.7	1.8	37.1	3.5	5.2	2.6	8.5	2.6	3.4	0.8	1.7	100.0	399	
Central	70.5	29.5	38.2	1.1	37.4	0.0	2.8	7.8	7.1	0.4	1.0	2.1	2.0	100.0	773	
North	67.9	32.1	27.1	0.4	44.9	0.4	2.2	1.8	13.9	3.6	0.7	1.8	3.2	100.0	362	
East	64.8	35.2	38.4	0.6	28.9	0.0	5.1	2.7	18.7	0.5	0.0	4.5	0.6	100.0	322	
Occupation																
Agricultural	70.3	29.7	28.6	0.2	46.1	0.6	0.2	0.0	16.1	1.7	0.4	2.4	3.6	100.0	1,494	
Non-agricultural	78.6	21.4	40.9	1.8	23.1	0.6	10.2	14.8	3.2	0.2	2.1	2.2	1.0	100.0	1,513	
Employment status																
All year, full week	77.8	22.2	31.1	0.8	29.4	0.7	10.4	15.3	6.1	0.2	1.1	2.2	2.9	100.0	1,292	
All year, part week	82.3	17.7	31.1	5.5	28.6	0.0	9.1	21.5	4.1	0.0	0.0	0.0	0.0	100.0	105	
Seasonal	72.2	27.8	27.7	0.8	47.2	0.6	0.4	0.0	15.8	2.0	0.4	2.3	2.8	100.0	1,286	
Occasional	67.8	32.2	62.8	0.7	20.6	0.6	1.0	0.0	6.6	0.3	3.8	3.0	0.7	100.0	334	
Total	74.5	25.5	34.0	0.9	36.4	0.6	4.4	6.2	10.6	1.0	1.1	2.3	2.5	100.0	3,019	

Note: Total includes 3 women for whom information on employment status was not available and 13 women for whom information on occupation is not available.

¹ Respondent is currently employed but has not worked since last birth.

CHAPTER 3

FERTILITY

Aykut Toros

The fertility measures presented in this chapter are based on the retrospective reproductive histories of ever-married women age 15-49 interviewed in the TDHS. Each ever-married woman was asked the number of sons and daughters living with her, the number living elsewhere, and the number who had died. She was then asked for a history of all her births, including the month and year, and the name and sex of each birth; if deceased, the age at death was also asked. If alive, the current age and whether the child was living with the mother were also asked. Based on this information, measures of completed fertility (number of children ever born) and current fertility (age-specific rates) were calculated. These measures are analyzed in connection with various background characteristics.

Cumulative fertility and children ever born are also looked at in this chapter. The tables display the data on children ever born by the woman's current age and by her age at marriage. The chapter concludes with an analysis of information on the age of the woman at the time of her first birth. The data are important because they indicate the beginning of the woman's reproductive life.

3.1 Current Fertility

The current level of fertility is the most important topic in this chapter because of its direct relevance to population policies and programs. Age-specific fertility rates (ASFR) for the three-year period before the survey are presented in Table 3.1 and Figure 3.1 for the country as a whole and for urban and rural areas. In addition, the total fertility rate (TFR) for women 15-44 years is shown in the table.

Numerators for the age-specific fertility rates in Table 3.1 are calculated by isolating live births that occurred during the three years preceding the survey (determined from the date of birth of the child), and classifying them by age of the mother (in five-year age groups) at the time of birth (determined from the date of birth of the mother). The denominators of the rates are the number of woman-years lived in each of the specified five-year age groups during the three years preceding the survey.

The crude birth rate (also shown in Table 3.1) is calculated by summing the product of the age-specific rates multiplied by the proportion of women in the specific age group out of the total *de facto* population, male and female. The general fertility rate is calculated as the number of births per thousand women in the reproductive age groups.

Age-specific fertility rates are estimated for the three years preceding the survey. As is typical, the distribution is skewed towards the younger ages. The highest fertility rate is observed for the age group 20-24. After age 24, the rates decline steadily, implying modern levels of fertility control in upper ages.


Table 3.1 Current fertility

Age-specific and cumulative fertility rates and the crude birth rate for the three years preceding the survey, by urban-rural residence, Turkey 1998

Age group	Residence		Total
	Urban	Rural	
15-19	55	68	60
20-24	141	211	163
25-29	139	178	150
30-34	97	85	93
35-39	32	60	42
40-44	14	12	13
45-49	0	2	1
TFR 15-49	2.39	3.08	2.61
TFR 15-44	2.39	3.08	2.61
GFR	87	107	94
CBR	22.8	24.7	23.4

Note: Rates are for the period 1-36 months preceding the survey. Rates for age group 45-49 may be slightly biased due to truncation.
 TFR: Total fertility rate expressed per woman
 GFR: General fertility rate (births divided by number of women 15-44), expressed per 1,000 women

Figure 3.1
Age-Specific Fertility Rates by Urban-Rural Residence



TDHS 1998

The total fertility rate (number of children a woman would bear if she lived through these rates throughout her reproductive life span) is slightly over three children (3.1) for women living in rural areas, and decreases to around two children (2.4) in urban areas. The national average is 2.6 children per woman. When compared with evidence from previous surveys (see HIPS, 1980, 1987, and 1989) the urban/rural gap appears to be closing.

The crude birth rate has fallen to the lower twenties. As expected, birth rates are higher in rural areas (24.7 per thousand) than in urban areas (22.8 per thousand). The national average (23.4 per thousand) implies a rather low population growth rate even if the crude death rate is very low.

The total fertility rates for the three-year period prior to the TDHS-98 are summarized in Table 3.2 for major groups in the population. The table also provides a basis for inferring trends in fertility by comparing current synthetic measures with the average number of children ever born to women currently 40-49 years of age. Although comparison of completed fertility among women age 40 or more with the total fertility rate can provide an indication of fertility change, such an approach is vulnerable to an understatement of parity for older women. The findings on contraceptive use (Chapter 4) and nuptiality (Chapter 6) are also of crucial importance in reaching a balanced judgment about fertility trends.

<u>Table 3.2 Fertility by background characteristics</u>			
Total fertility rate for the three years preceding the survey and mean number of children ever born to women age 40-49, by selected background characteristics, Turkey 1998			
Background characteristic	Total fertility rate	Percentage currently pregnant	Mean number of children ever born to women age 40-49
Residence			
Urban	2.39	4.8	3.82
Rural	3.08	5.4	5.02
Region			
West	2.03	3.5	3.43
South	2.55	5.3	4.46
Central	2.56	5.5	3.84
North	2.68	4.4	4.36
East	4.19	8.0	7.00
Education			
No educ./Pri. incomp.	3.89	6.0	5.63
Pri. comp./Sec. incomp.	2.55	5.4	3.40
Sec. comp./+	1.61	2.6	1.84
Total	2.61	5.0	4.22
† Rate for women age 15-49 years			

There are clear variations in fertility levels by region and education. These variations are evident for past fertility experience (mean number of children for women age 40-49) as well as current fertility levels (total fertility rates). Regional variations of fertility involve three regional groupings. The Eastern region is notable as a high fertility region, with a total fertility rate exceeding four children (4.2). The North, Central and South regions constitute another group, with rates well under three children (2.7, 2.6, and 2.6, respectively). The lowest rate (2.0) is found in the West region and is comparable to that of many Western European countries.

The regional groupings based on current levels of fertility are also cogent for examining differences in the past fertility experience. Although the mean number of children born to women age 40-49 is much higher (between 50 to 75 percent) than the corresponding TFRs in each of the regions, notable variations are observed as with current fertility. The table suggests an overall decline in fertility, keeping regional differences almost the same, during the last three decades.

Past experience as well as current levels of fertility show strong variations by levels of education. Both the total fertility rate and the number of children ever born are more than fifty percent lower among women with at least a secondary level of education compared to women with no education.

Fertility trends can be analyzed in two ways. One is to compare TDHS data with the results of previous surveys. Fertility trends can also be examined based on TDHS data alone. Having the complete birth history provides more direct evidence on trends, thereby permitting more accurate conclusions. However, use of birth histories for analysis of trends places a great burden on the quality of data, which should always be interpreted with caution. Table 3.3 shows the age-specific fertility rates for five-year periods preceding the survey. The age-specific schedule of rates in Table 3.3 is progressively truncated as time before the survey increases. The bottom diagonal of estimates (enclosed in brackets) is also truncated. Total fertility rates can be calculated from the age-specific rates in Table 3.3, but only by summing across ages unaffected by truncation.

Table 3.3 Age-specific fertility rates				
Age-specific fertility rates for five-year periods preceding the survey, by mother's age, Turkey 1998				
Mother's age	Number of years preceding the survey			
	0-4	5-9	10-14	15-19
15-19	62	79	90	122
20-24	169	190	221	259
25-29	148	166	189	221
30-34	91	103	119	[176]
35-39	40	52	[70]	-
40-44	14	[20]	-	-
45-49	[1]	-	-	-


Note: Age-specific fertility rates are per 1,000 women. Estimates enclosed in brackets are truncated.

The decline of fertility over time, which is implied by the earlier tables, is seen much more clearly in Figure 3.2. Cumulation of the ASFRs up to age 40 shows a recent decline in fertility of fourteen percent, from 3 births per woman in the period 5 to 9 years before the survey to 2.6 births per woman in the five-year period immediately preceding the survey.

Table 3.4 presents fertility rates for ever-married women by duration since first marriage for five-year periods preceding the survey. These rates are similar to those presented in Table 3.3 and the same admonitions apply in their interpretation. Fertility early in marriage often remains resistant to change, even when fertility is declining, because fertility decline usually begins at the older ages (when women start to limit the number of births) and not among young couples postponing births. Therefore, a complete examination of duration-specific trends requires interpretation in the light of other evidence.

Figure 3.2

Age-Specific Fertility Rates during the Last 20 Years



Note: Dates are approximate and refer to September 1998, the mid-point of fieldwork.

TDHS 1998

Fertility rates are declining in general, but as shown earlier, the decline is greater among women who are in their later years of childbearing. Table 3.4 indicates that a decline of fertility by around one-fifth, from 358 to 293, among women married 0-4 years and, thus, in the early years of childbearing; in fact, this decline is slightly more than the decline observed in TDHS-93. An even more substantial decline (by almost 40 percent from 264 to 164) is observed for women married 5-9 years and very dramatic changes (around than sixty percent) occurred among women who had been married for longer durations. Although this pattern is quite common among populations with increasing fertility control, the speed of change is noteworthy.

The table also indicates that the decline in fertility was more rapid during the early 1990s than during the late 1980s.

Table 3.4 Fertility by marital duration

Fertility rates for ever-married women by duration since first marriage in years, for five-year periods preceding the survey. Turkey 1998

Marriage duration at birth	Number of years preceding the survey			
	0-4	5-9	10-14	15-19
0-4	293	309	335	358
5-9	164	182	214	264
10-14	85	107	147	210
15-19	49	75	104	[161]
20-24	23	50	[106]	-
25-29	12	[21]	-	-

Note: Fertility rates are per 1,000 women. Estimates enclosed in brackets are truncated.

3.2 Children Ever Born and Living

The distribution of women by number of children ever born is presented in Table 3.5 for all women and for currently married women. In the TDHS-98 questionnaire, the total number of children ever born was ascertained by a sequence of questions designed to maximize recall. Life-time fertility reflects the accumulation of births over the past 30 years and, therefore, its relevance to the current situation is limited.

Table 3.5 Children ever born and living

Percent distribution of all women and of currently married women by number of children ever born (CEB) and mean number ever born and living, according to five-year age groups, Turkey 1998

Age group	Number of children ever born (CEB)											Total	Number of women	Mean no. of CEB	Mean no of living children	
	0	1	2	3	4	5	6	7	8	9	10+					
ALL WOMEN																
15-19	92.1	6.7	0.9	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	100.0	1,720	0.09	0.09	
20-24	53.9	24.1	15.8	4.3	1.3	0.4	0.1	0.0	0.0	0.0	0.0	100.0	1,558	0.76	0.72	
25-29	22.8	22.5	32.0	13.8	4.5	2.6	1.0	0.4	0.2	0.1	0.0	100.0	1,397	1.71	1.60	
30-34	10.2	12.7	33.2	20.5	12.0	5.1	2.5	2.1	1.0	0.4	0.4	100.0	1,202	2.59	2.39	
35-39	6.0	7.6	29.3	21.2	15.5	7.5	4.6	2.7	2.4	1.6	1.6	100.0	1,081	3.27	2.95	
40-44	3.8	5.6	22.4	20.7	14.6	10.9	7.1	3.9	4.6	3.5	2.9	100.0	885	3.96	3.51	
45-49	3.7	4.5	15.7	17.8	16.0	11.7	10.1	5.8	5.5	3.0	6.1	100.0	733	4.54	3.80	
Total	34.9	13.1	20.3	12.3	7.5	4.3	2.7	1.6	1.4	0.9	1.1	100.0	8,576	2.01	1.80	
CURRENTLY MARRIED WOMEN																
15-19	49.6	42.9	6.2	1.1	0.2	0.0	0.0	0.0	0.0	0.0	0.0	100.0	262	0.59	0.56	
20-24	23.8	39.7	26.3	7.3	2.2	0.6	0.1	0.0	0.0	0.0	0.0	100.0	924	1.27	1.20	
25-29	11.1	25.5	37.1	16.0	5.2	3.1	1.1	0.5	0.2	0.1	0.0	100.0	1,196	1.97	1.85	
30-34	3.7	13.1	35.6	22.2	13.3	5.5	2.7	2.2	1.1	0.4	0.4	100.0	1,090	2.79	2.57	
35-39	3.6	7.0	30.0	22.4	16.3	7.5	4.7	2.7	2.4	1.7	1.7	100.0	1,014	3.37	3.04	
40-44	1.7	5.5	22.3	20.3	15.4	11.7	7.9	4.1	5.0	3.4	2.8	100.0	789	4.08	3.62	
45-49	2.2	3.6	15.7	18.5	16.7	12.4	10.3	6.0	4.9	2.9	6.6	100.0	645	4.64	3.89	
Total	9.9	18.0	28.2	17.1	10.5	5.9	3.7	2.2	1.9	1.2	1.5	100.0	5,921	2.76	2.49	

The results in Table 3.5 for younger women who are currently married differ from those for the sample as a whole because of the large number of unmarried women with minimal fertility. Differences at older ages, though minimal, generally reflect the impact of marital dissolution. The parity distribution for older currently married women provides a measure of primary infertility.

A comparison of the mean number of children ever born with the mean number of children surviving offers a quick evaluation of the survival status of the children. Around one in six children born by women age 45-49 had not survived at the time of the survey (4.5 versus 3.8). The proportion of children surviving among younger women is much higher. This may not only reflect the shorter exposure to risk by the children of the younger cohorts, but also is evidence of improved mortality conditions in general. Of all children born (mean of 2.0), 90 percent (mean of 1.8) were still alive at the time of the survey.

As marriage is universal in Turkey (see Chapter 6), the proportion of women remaining childless is very low. The proportion of women with no children declines in tandem with the proportion remaining single, and almost all women who are married by the age of 45-49 have children. Just over 2 percent of the currently married women who are about to complete their reproductive period remain childless, probably largely due to sterility rather than preference.

3.3 Birth Intervals

There has been a fair amount of research to indicate that short birth intervals are deleterious to the health of babies. This is particularly true for babies born at intervals of less than 24 months. Table 3.6 shows the percent distribution of non-first births in the five years preceding the survey by the number of months since the previous birth.

Table 3.6 Birth intervals

Percent distribution of births in the five years preceding the survey by number of months since previous birth, according to demographic and socioeconomic characteristics, Turkey 1998

Characteristic	Number of months since previous birth					Total	Number of births	Median number of month since previous birth
	7-17	18-23	24-35	36-47	48+			
Age of mother								
15-19	(29.1)	(27.0)	(30.8)	(13.1)	(0.0)	100.0	23	23.4
20-29	18.5	16.4	24.2	17.9	23.0	100.0	1,187	30.7
30-39	8.1	7.8	19.9	15.7	48.6	100.0	919	46.6
40+	3.4	7.9	15.1	9.0	64.6	100.0	125	-
Birth order								
2-3	13.6	13.3	20.2	17.2	35.6	100.0	1,482	37.6
4-6	14.0	9.3	23.6	15.2	37.8	100.0	544	37.5
7+	11.8	14.8	29.7	14.6	29.1	100.0	229	33.6
Sex of prior birth								
Male	13.3	11.3	21.7	16.9	36.8	100.0	1,141	38.0
Female	13.7	13.8	22.4	16.0	34.2	100.0	1,113	36.1
Survival of prior birth								
Living	11.6	12.0	22.0	17.0	37.4	100.0	2,096	38.4
Dead	39.5	20.0	22.0	8.9	9.7	100.0	158	21.6
Residence								
Urban	11.1	10.9	20.0	17.7	40.4	100.0	1,359	40.6
Rural	17.2	15.1	25.1	14.5	28.1	100.0	895	31.6
Region								
West	10.0	7.8	17.6	19.3	45.2	100.0	594	44.3
South	10.0	11.8	20.3	18.1	39.8	100.0	311	39.4
Central	11.2	13.8	20.8	14.6	39.6	100.0	496	38.5
North	15.0	10.9	24.4	13.3	36.3	100.0	182	35.7
East	19.5	16.5	26.9	15.4	21.6	100.0	672	28.7
Education								
No educ./Pri. incomp.	14.4	14.0	29.7	15.0	26.9	100.0	790	32.1
Pri. comp./Sec. incomp.	13.5	12.4	18.5	17.4	38.2	100.0	1,279	39.1
Sec. comp./+	9.9	7.1	13.2	16.1	53.6	100.0	185	49.9
Total	13.5	12.5	22.0	16.5	35.5	100.0	2,254	37.0

Note: First-order births are excluded. The interval for multiple births is the number of months since the preceding pregnancy that ended in a live birth.

() Figures in parentheses are based on 25-49 unweighted births.

The median birth interval is slightly over three years (37 months). This is more than a year longer than the minimum considered safe. One-fourth of non-first births were born with intervals of less than 24 months. This percentage shows striking variations by background variables. Among births to women with at least a secondary education, the percentage born within 24 months of a prior birth is about 40 percent lower than the percentage of short interval births among women with no education (17 percent and 28 percent, respectively). The smallest proportion of short birth intervals is observed in the West region and the highest proportion in the East region (18 percent and 36 percent, respectively). Short intervals following a female birth are more frequent than for male births (28 percent and 25 percent, respectively). Among all the factors presented in the table, the survival status of the preceding child appears to be strongly related to the proportion of short birth intervals (24 percent for surviving children and 60 percent for deceased children).

3.4 Age at First Birth

The age at which childbearing begins has important demographic consequences as well as important consequences for the mother and child. In many countries, postponement of first births, reflecting an increase in the age at marriage, has contributed greatly to overall fertility decline. The proportion of women who become mothers before the age of 20 is also a measure of the magnitude of adolescent fertility, which is a major health and social concern in many countries. Table 3.7 presents the distribution of Turkish women by age at first birth, according to their current age.

Table 3.7 Age at first birth
Percent distribution of women 15-49 by age at first birth, according to current age, Turkey 1998

Current age	Women with no births	Age at first birth						Total	Number of women	Median age at first birth
		<15	15-17	18-19	20-21	22-24	25+			
15-19	92.1	0.3	5.2	2.4	NA	NA	NA	100.0	1,720	a
20-24	53.9	0.9	10.0	15.3	13.5	6.4	NA	100.0	1,558	a
25-29	22.8	1.7	12.7	15.6	18.1	20.2	8.8	100.0	1,397	22.2
30-34	10.2	1.8	13.0	17.2	18.9	20.9	18.0	100.0	1,202	21.9
35-39	6.0	3.0	17.4	21.2	16.2	17.8	18.3	100.0	1,081	20.9
40-44	3.8	1.6	18.4	24.2	20.7	16.7	14.6	100.0	885	20.5
45-49	3.7	3.7	20.0	20.2	19.3	20.0	13.3	100.0	733	20.6

NA = Not applicable
^a Less than 50 percent of the women in group x to $x + 4$ have had a birth by age x .

Age of childbearing is increasing gradually. The median has risen from 20.6 years among women age 45-49 years to 22.2 years among women age 25-29 years. The table indicates dramatic changes in adolescent fertility. Some 26 percent of women age 20-24 during the survey had become mothers before age 20; this percentage is substantially lower than the percentage for women age 30-34 (32 percent). Among women aged 35-44 at the time of the survey, more than 40 percent or more had become mothers in their teens.

The median age at first birth for different cohorts is summarised in Table 3.8 and the entry age into motherhood for different subgroups of the population can be compared (the medians for cohorts 15-19 and 20-24 could not be determined because half the women had not yet had a birth).

Table 3.8 Median age at first birth by background characteristics

Median age at first birth among women 25-49, by current age and selected background characteristics, Turkey 1998

Background characteristic	Current age					Women age 25-49
	25-29	30-34	35-39	40-44	45-49	
Residence						
Urban	22.6	22.2	21.1	20.8	20.8	21.7
Rural	21.6	21.3	20.5	19.9	20.4	20.7
Region						
West	22.7	22.4	21.5	20.9	20.8	21.8
South	23.0	21.9	21.6	20.4	19.9	21.5
Central	22.0	21.4	20.5	20.4	20.9	21.1
North	22.2	22.2	20.7	20.4	20.9	21.3
East	20.7	20.5	20.2	19.7	19.8	20.2
Education						
No educ./Pri. incomp.	20.1	19.5	19.9	19.5	19.7	19.7
Pri. comp./Sec. incomp.	21.6	21.5	20.3	20.7	20.7	21.1
Sec. comp./+	-	25.9	24.9	23.8	24.7	-
Total	22.2	21.9	20.9	20.5	20.6	21.3

Note: The medians for cohorts 15-19 and 20-24 could not be determined because some women may still have a birth before reaching age 20 or 25, respectively.

The median age at first birth is over 21 years (21.3) among all women 25-49. It varies considerably according to background variables. Women living in urban areas had their first birth one year later than women living in rural areas. When they first become mothers, women living in the East region were 1.6 years younger than women living in the West region. The level of education shows the biggest differentials among the background variables considered in this table. For example, women age 30-34 with no education became mothers on average at the age of 19.5, while women in the same cohort with at least a secondary level of education waited an additional six years before they had the first birth.

3.5 Teenage Pregnancy and Motherhood

Table 3.9 shows the percentage of women age 15-19 who are mothers or pregnant with their first child. About one in twelve (9 percent) of women age 17 have become mothers or are pregnant with their first child. The proportion increases steeply to more than one in six (16 percent) among women age 18 and close to one in four (23 percent) among women age 19. Higher proportions of teenagers living in rural areas have begun childbearing than teenagers living in urban areas (11 percent and 9 percent, respectively). Although fertility is highest in the East region, the proportion of teenagers who have begun childbearing is virtually the same in the Central and South regions as in the East. Education appears to have the strongest association with teenage fertility, not only because the time spent in school results in later marriage and postponed births, but also because of changes in childbearing attitudes.

Table 3.9 Teenage pregnancy and motherhood

Percentage of teenagers 15-19 who are mothers or pregnant with their first child, by selected background characteristics, Turkey 1998

Background characteristic	Percentage who are:		Percentage who have begun child-bearing	Number of teenagers
	Mothers	Pregnant with first child		
Age				
15	1.3	0.5	1.8	305
16	1.6	0.3	1.9	372
17	4.9	3.8	8.8	347
18	12.2	3.7	16.0	396
19	20.1	3.0	23.1	301
Residence				
Urban	7.4	2.0	9.4	1,034
Rural	8.6	2.8	11.4	686
Region				
West	7.2	2.2	9.3	539
South	8.4	2.6	11.0	261
Central	8.8	2.4	11.2	380
North	4.3	2.5	6.7	136
East	8.8	2.2	11.0	404
Education				
No educ./Pri. incomp.	15.3	6.7	22.0	183
Pri. comp./Sec. incomp	8.3	2.1	10.4	1,239
Sec. comp./+	1.6	0.4	2.0	297
Total	7.9	2.3	10.2	1,720

Note: The sum of the absolute values does not add up to the total value in the last three categories due to the ever-married factors used.

CHAPTER 4

FAMILY PLANNING

Turgay Ünalán and İsmet Koç

This chapter presents the TDHS-98 results regarding various aspects of contraceptive knowledge, attitudes and behaviour. While the focus is on women, some results from the husband survey will also be presented, since men play an important role in the realisation of reproductive goals. To get an indication of interspousal communication and agreement (perceived) in the attitudes and knowledge of couples regarding family planning, the responses of men were, where possible, paired with responses obtained from their wives in the same household.

To obtain the data on knowledge and use of family planning, respondents were first asked to name all of the methods of family planning that they knew or had heard about. For methods which were not mentioned spontaneously, a description of the method was read and the respondents were asked if they had heard of the method. Respondents were then asked if they were currently using a method and, if so, from where they had obtained the currently used method.

4.1 Knowledge of Family Planning Methods

Table 4.1 presents the levels knowledge of contraceptive methods for female respondents by marital status and for husbands. Knowledge of modern methods is almost universal. Knowledge of traditional methods is also high among married respondents (91 percent among currently married women and 88 percent among husbands). However, knowledge of traditional methods is much lower among never-married women, with only about three in five recognizing any traditional method. The IUD and

Contraceptive method	All women	Never-married women	Currently married women	Husbands
Any method	98.2	96.5	98.9	97.9
Any modern method	98.0	96.5	98.7	97.1
Pill	94.4	91.0	95.8	92.5
IUD	94.9	89.7	97.0	87.0
Injections	72.8	64.9	76.2	61.7
Diaphragm/Foam/Jelly	45.0	30.5	50.5	26.0
Condom	79.9	66.9	85.1	84.6
Female sterilisation	79.1	67.9	83.4	68.8
Male sterilisation	41.4	32.2	45.0	49.4
Implant	22.2	15.1	24.8	13.1
Any traditional method	81.7	59.0	90.6	87.8
Periodic abstinence	43.6	38.3	45.4	61.0
Withdrawal	77.4	46.5	89.4	83.9
Other methods	6.0	2.7	7.2	2.9
Any traditional/folk method	82.0	59.3	90.8	87.8
Total	8,576	2,380	5,921	1,971
Mean	6.6	5.5	7.0	6.3

pill are the most widely known family planning methods among women and husbands followed by the condom and female sterilisation. Withdrawal, a traditional method of avoiding pregnancy, is known to 89 percent of currently married women and 84 percent of husbands, but to only 47 percent of never-married women. The mean number of methods known is a rough indicator of the breadth of family planning methods. On average, 7 methods are known by currently married women, 6.3 methods are known by husbands, and 5.5 methods by never-married women.

Table 4.2 shows the correspondence between the contraceptive knowledge of husbands and wives for the 1,896 couples interviewed in the TDHS-98. The proportion of couples where both spouses know at least one method of contraception is 97 percent. This proportion is 96 percent for modern methods and 81 percent for traditional methods. Eighty-six percent of couples know the IUD and 89 percent know the pill. The proportion of couples with withdrawal knowledge is also high (78 percent). For couples where only one partner knows of a method, wives are more likely to know the method than their husbands; the exceptions are male sterilisation and periodic abstinence.

Table 4.2 Couples' knowledge of contraceptive methods
Percent distribution of couples by knowledge of specific contraceptive methods, Turkey 1998

Contraceptive method	Both know method	Husband knows method, not wife	Wife knows method, not husband	Neither know	Total	Number of couples
Any method	97.0	1.0	1.8	0.2	100.0	1,896
Any modern method	95.9	1.4	2.5	0.2	100.0	1,896
Pill	88.9	3.9	6.1	1.1	100.0	1,896
IUD	85.9	1.3	10.7	2.1	100.0	1,896
Injections	49.6	12.5	26.4	11.5	100.0	1,896
Diaphragm/Foam/Jelly	16.3	9.2	34.4	40.1	100.0	1,896
Condom	75.7	8.9	9.6	5.9	100.0	1,896
Female sterilisation	59.4	9.6	22.8	8.2	100.0	1,896
Male sterilisation	28.3	21.2	16.3	34.2	100.0	1,896
Implant	6.4	6.7	17.8	69.1	100.0	1,896
Any traditional method	81.4	6.4	9.3	2.9	100.0	1,896
Periodic abstinence	34.3	26.5	12.1	27.0	100.0	1,896
Withdrawal	77.6	6.4	11.8	4.2	100.0	1,896
Any folk method	0.3	2.7	7.4	89.6	100.0	1,896
Any traditional/folk method	81.8	6.1	9.3	2.9	100.0	1,896

4.2 Ever Use of Family Planning Methods

Ever-married women and husbands interviewed in the TDHS-98 who reported that they had heard of a method of family planning were asked if they had ever used that method. Table 4.3 looks at the extent to which women and husbands report having had experience with the use of contraceptive methods. The proportion of currently married women who have ever used any contraceptive method is 84 percent. Overall, modern methods are much more frequently adopted than traditional methods; 67 percent of all currently married women and 68 percent of husbands have used a modern method while 60 percent of currently married women and 56 percent of husbands have used a traditional method.

Table 4.3 Ever use of contraception

Among currently married women and husbands, the percentage who have ever used a contraceptive method, by specific method, according to age, Turkey 1998

Age	Modern methods										Traditional methods					Number of women/husbands
	Any modern method	Pill	IUD	Injection	Diaphragm/ Foam/ Jelly	Condom	Female sterilisation	Male sterilisation	Implants	Any trad. method	Periodic abstinence	Withdrawal	Other methods	Any trad./folk method		
CURRENTLY MARRIED WOMEN																
15-19	54.9	26.4	7.6	9.6	0.8	1.0	16.5	0.0	0.0	0.0	44.7	3.6	44.2	0.0	44.7	262
20-24	74.8	49.7	17.5	23.4	1.5	3.0	29.5	0.3	0.0	0.0	56.2	3.8	55.2	0.7	56.6	924
25-29	85.6	70.8	29.8	40.9	3.5	4.1	35.4	1.8	0.0	0.1	61.4	8.5	59.2	0.7	61.8	1,196
30-34	90.2	77.3	39.7	46.4	3.3	7.1	36.9	5.9	0.0	0.1	63.6	9.0	61.1	1.3	63.9	1,090
35-39	89.2	75.7	41.3	52.8	2.5	10.5	32.8	7.1	0.0	0.2	62.0	9.7	59.8	3.0	63.3	1,014
40-44	87.7	71.9	43.3	42.3	3.5	12.8	29.2	7.0	0.0	0.0	63.2	9.7	61.0	4.2	65.2	789
45-49	84.5	66.9	45.2	30.5	3.9	13.1	20.3	5.7	0.2	0.0	56.6	9.7	54.4	4.2	58.3	645
Total	84.2	67.3	34.2	38.9	2.9	7.6	31.0	4.2	0.0	0.1	60.1	8.1	58.1	2.0	61.0	5,921
HUSBANDS																
15-24	60.4	30.9	11.7	8.6	1.7	0.8	15.2	1.2	0.0	0.0	46.1	7.4	41.6	0.0	46.1	112
25-29	78.6	60.2	21.6	24.4	5.1	0.9	38.2	0.2	0.0	0.0	56.8	16.5	51.3	0.5	57.1	342
30-34	84.8	70.5	28.0	38.2	1.6	3.2	45.5	2.8	0.4	0.2	60.9	20.1	56.4	0.9	60.9	364
35-39	90.1	78.4	36.8	45.7	4.7	3.2	38.3	5.2	0.4	0.4	60.0	20.4	54.3	1.4	60.4	352
40-44	85.5	75.6	37.9	45.0	6.1	2.6	34.8	6.8	0.0	0.0	51.9	17.0	48.5	1.8	51.9	335
45-49	85.5	70.1	47.0	35.3	9.9	8.7	32.0	6.9	0.0	0.0	59.4	21.3	53.1	0.8	59.5	240
50+	72.7	59.8	34.0	31.5	6.3	6.6	26.4	3.3	0.0	0.0	46.2	14.0	42.7	0.4	46.5	226
Total	82.1	67.5	32.2	35.5	5.1	3.6	35.6	3.9	0.1	0.1	55.8	17.7	51.0	0.9	56.0	1,971

With regard to the ever use of specific methods of contraception, slight differences are observed between the levels reported by women and husbands. The proportion of those who reported ever use of the pill, IUD and the diaphragm, foam and jelly, were slightly higher among women while the proportions ever reporting use of condom and injections were slightly higher among husbands. With respect to traditional methods, women were somewhat more likely to report ever use of withdrawal than men, while men were more likely to report ever use of periodic abstinence.

4.3 Current Use of Contraceptive Methods

The level of current use of family planning is one of the indicators most frequently used to assess the success of the family planning programme activities. It is widely used as a measure in the analysis of the determinants of fertility. This section focuses on the levels and differentials in family planning use, with particular emphasis on the method mix among users.

Table 4.4 presents the level of current use of contraceptives for currently married women and husbands by age group. Overall, 64 percent of currently married women in Turkey are currently using a method of contraception. Of the users, the majority rely on a modern method (Figure 4.1). One in five currently married women are using the IUD. The condom, which is the second most popular modern method, is used by approximately 8 percent of married women and 11 percent of the husbands. Withdrawal, a traditional method, is the most popular method among currently married women in Turkey, with 24 percent currently using the method.

Table 4.4 Current use of contraception

Percent distribution of all women, currently married women, and husbands, by contraceptive method currently used, according to age, Turkey 1998

Age	Modern methods								Traditional methods					Not currently using	Number of women/husbands	
	Any method	Any modern method	Pill	IUD	Injection	Dia-phragm/ Foam/ Jelly	Con-dom	Female steri-lisa-tion	Male steri-lisa-tion	Any trad. method	Peri-odic absti-nence	With-draw-al	Other meth-ods			
CURRENTLY MARRIED WOMEN																
15-19	33.6	15.7	1.9	7.4	0.5	0.0	6.0	0.0	0.0	17.8	0.5	17.3	0.0	66.4	100.0	262
20-24	52.9	30.8	4.8	16.3	0.3	0.3	8.8	0.3	0.0	21.9	0.3	21.6	0.2	47.1	100.0	924
25-29	67.0	43.3	6.1	22.9	1.1	0.5	10.9	1.8	0.0	23.4	0.5	22.9	0.3	33.0	100.0	1,196
30-34	74.3	47.3	5.2	25.6	0.3	0.5	9.7	5.9	0.0	26.7	0.8	25.8	0.4	25.7	100.0	1,090
35-39	76.3	46.6	3.9	27.4	0.7	0.8	6.8	7.1	0.0	28.8	1.3	27.5	0.9	23.7	100.0	1,014
40-44	70.0	36.6	3.4	16.6	0.0	1.0	8.6	7.0	0.0	31.9	1.9	30.0	1.5	30.0	100.0	789
45-49	41.4	17.6	2.1	6.4	0.0	0.8	2.5	5.7	0.2	22.6	2.6	20.1	1.1	58.6	100.0	645
Total	63.9	37.7	4.4	19.8	0.5	0.6	8.2	4.2	0.0	25.5	1.1	24.4	0.6	36.1	100.0	5,921
HUSBANDS																
15-24	33.9	17.7	5.8	7.9	0.0	0.0	2.9	1.2	0.0	16.2	0.0	16.2	0.0	66.1	100.0	112
25-29	60.4	39.4	6.8	14.9	1.4	0.2	15.9	0.2	0.0	20.8	1.8	19.0	0.2	39.6	100.0	342
30-34	69.6	50.0	8.8	21.7	0.2	0.4	15.8	2.8	0.4	19.2	1.2	18.0	0.4	30.4	100.0	364
35-39	75.4	50.0	7.7	24.3	0.2	0.4	12.3	4.8	0.4	24.5	1.9	22.7	0.9	24.6	100.0	352
40-44	68.9	50.6	7.0	26.2	1.1	0.5	9.0	6.8	0.0	17.9	1.2	16.7	0.4	31.1	100.0	335
45-49	61.8	40.2	7.1	15.7	0.0	1.6	8.8	6.9	0.0	21.5	2.3	19.2	0.1	38.2	100.0	240
50+	40.7	23.8	1.8	9.5	0.0	2.7	6.6	3.3	0.0	16.7	2.4	14.3	0.2	59.3	100.0	226
Total	62.6	42.2	6.8	18.8	0.5	0.7	11.4	3.9	0.1	20.0	1.6	18.4	0.4	37.4	100.0	1,971

Figure 4.1
Current Use of Family Planning Methods, Turkey 1993 and 1998




Table 4.4 also shows the variation in current use levels by age. Younger and older women are much less likely to be using contraception than women 25-44. As Table 4.5 shows, contraceptive use also increases rapidly with the number of living children, peaking at 78 percent among women with two children, after which it declines to 59 percent among women with four or more children. There appears to be only limited effort to delay the first birth; 18 percent of the currently married women with no children are using a method. The proportions of women currently using correlates positively with educational level, with the largest proportion observed among women with a secondary or higher education. Women with secondary or higher education are also more likely to be using modern contraceptive methods, especially the IUD and the condom, than less educated women. More than half of all women in this education group are users of a modern method, with almost a quarter using the IUD.


There are marked differences by residence in the proportion of women currently using a modern contraceptive method. Urban women are considerably more likely to be using a modern method than rural women. While 71 percent of currently married women in the West region are using some kind of contraception, the proportion is 42 percent in the East (Figure 4.2). The proportions using a modern method are highest in the Central and West regions (43 and 41 percent, respectively) followed by the South, North and the East regions. The use of withdrawal seems to be popular across the country but principally in the North (31 percent) and the West (28 percent).

Table 4.5 Current use of contraception by background characteristics

Percent distribution of currently married women and husbands by contraceptive method currently used, according to selected background characteristics, Turkey 1998

Background characteristic	Modern methods									Traditional methods				Not currently using	Total	Number of women/husbands
	Any modern method	Any modern method	Pill	IUD	Injection	Dia-phragm/ Foam/ Jelly	Con-dom	Female steri-lisa-tion	Male steri-lisa-tion	Any trad. method	Peri-odic absti-nence	With-draw-al	Other meth-ods			
CURRENTLY MARRIED WOMEN																
Number of living children																
None	17.6	9.2	3.3	0.1	0.0	0.2	5.5	0.0	0.0	8.4	0.5	8.0	0.0	82.4	100.0	613
1	59.6	33.1	3.9	17.9	0.7	0.1	10.4	0.1	0.0	26.2	1.5	24.7	0.3	40.4	100.0	1,132
2	78.1	49.3	5.1	27.3	0.6	0.8	11.1	4.3	0.1	28.4	1.5	26.9	0.4	21.9	100.0	1,802
3	76.4	43.8	5.6	23.5	0.3	0.7	6.8	6.9	0.0	31.3	0.6	30.6	1.3	23.6	100.0	1,096
4+	59.0	34.0	3.1	17.1	0.5	1.0	4.8	7.5	0.0	23.9	0.7	23.1	1.1	41.0	100.0	1,279
Residence																
Urban	66.7	40.8	4.6	21.0	0.5	0.6	9.3	4.7	0.0	25.2	1.4	23.8	0.6	33.3	100.0	3,978
Rural	58.1	31.4	3.8	17.3	0.3	0.6	6.0	3.3	0.0	26.0	0.5	25.5	0.7	41.9	100.0	1,943
Region																
West	70.5	40.5	5.4	20.5	0.3	0.7	9.4	4.2	0.1	29.2	1.6	27.6	0.9	29.5	100.0	2,261
South	60.3	35.1	2.3	20.9	0.5	0.7	6.2	4.5	0.0	24.6	0.4	24.2	0.6	39.7	100.0	851
Central	68.3	42.8	4.9	24.2	0.5	0.6	9.2	3.4	0.0	24.7	1.1	23.7	0.8	31.7	100.0	1,426
North	67.0	35.2	4.5	12.4	0.0	0.7	9.2	8.4	0.0	31.5	0.7	30.9	0.3	33.0	100.0	474
East	42.0	26.7	2.9	14.0	1.1	0.2	5.2	3.3	0.0	15.2	0.7	14.4	0.1	58.0	100.0	909
Education																
No educ./																
Pri. incmp.	50.4	27.9	3.1	14.0	0.5	0.8	4.0	5.6	0.0	21.5	0.4	21.1	0.9	49.6	100.0	1,546
Pri. comp./																
Sec. incmp.	67.1	38.6	4.4	21.4	0.5	0.5	8.1	3.6	0.0	27.9	0.6	27.3	0.7	32.9	100.0	3,570
Sec. comp.+	75.3	52.7	6.4	23.8	0.2	0.8	17.0	4.5	0.0	22.5	4.4	18.1	0.1	24.7	100.0	804
Total	63.9	37.7	4.4	19.8	0.5	0.6	8.2	4.2	0.0	25.5	1.1	24.4	0.6	36.1	100.0	5,921
HUSBANDS																
Number of living children																
None	26.0	14.7	6.1	0.5	0.0	0.0	8.0	0.0	0.0	11.3	2.9	8.4	0.0	74.0	100.0	187
1	66.2	42.8	7.2	18.6	0.4	0.4	15.8	0.4	0.0	23.1	2.5	20.6	0.4	33.8	100.0	372
2	77.4	53.3	6.7	25.9	0.5	0.8	15.5	3.5	0.5	24.0	1.5	22.5	0.2	22.6	100.0	566
3	68.3	46.3	7.6	21.3	1.0	1.2	9.3	5.9	0.0	21.5	1.2	20.3	0.5	31.7	100.0	334
4+	53.3	37.1	6.3	16.3	0.4	1.0	6.2	6.9	0.0	15.7	0.9	14.7	0.6	46.7	100.0	512
Residence																
Urban	66.0	45.9	7.4	20.0	0.5	0.7	13.2	3.8	0.2	19.7	2.0	17.8	0.4	34.0	100.0	1,347
Rural	55.4	34.4	5.4	16.2	0.4	0.8	7.5	4.1	0.0	20.7	0.9	19.7	0.3	44.6	100.0	624
Region																
West	70.0	43.8	7.8	19.4	0.2	0.9	12.6	2.6	0.3	25.8	2.5	23.3	0.4	30.0	100.0	767
South	60.6	41.7	3.0	21.4	0.7	1.6	9.7	5.3	0.0	19.0	0.9	18.0	0.0	39.4	100.0	285
Central	67.9	49.9	8.5	22.4	0.9	0.6	12.7	4.8	0.0	17.4	1.6	15.8	0.6	32.1	100.0	481
North	58.5	36.9	7.9	11.2	0.0	0.6	11.5	5.7	0.0	21.1	0.4	20.7	0.5	41.5	100.0	150
East	38.1	28.5	4.3	12.9	0.7	0.0	7.4	3.2	0.0	9.4	0.5	8.9	0.2	61.9	100.0	287
Education																
No education/																
Pri. incmp.	32.7	17.9	0.8	11.3	0.5	0.0	2.2	3.2	0.0	14.8	0.3	14.4	0.0	67.3	100.0	204
Pri. comp./																
Sec. incmp.	62.4	41.5	7.5	18.6	0.3	0.9	10.0	4.3	0.0	20.5	0.8	19.7	0.4	37.6	100.0	1,253
Sec. comp.+	75.1	53.7	7.5	22.5	0.8	0.8	18.6	3.1	0.5	21.0	4.1	16.9	0.4	24.9	100.0	513
Total	62.6	42.2	6.8	18.8	0.5	0.7	11.4	3.9	0.1	20.0	1.6	18.4	0.4	37.4	100.0	1,971

Figure 4.2
Current Use of Family Planning by Region and Method



4.4 Trend in Contraceptive Use

Table 4.6 uses data from the 1988 TPHS, 1993 TDHS and the 1998 TDHS to provide the background to recent trends in current use of contraception in Turkey. A plateauing in contraceptive use at around 63 percent is apparent. However, there are a number of changes in the level of use of specific methods that are noteworthy. The proportions of women using IUD and female sterilisation in the TDHS-98 are higher than the levels reported in 1988 and 1993. In 1988, 14 percent of the currently married women were using the IUD, in 1993, 19 percent were IUD users and in 1998, 20 percent were using the IUD. In contrast, there has been a continuous decline in the proportion of women using the pill, from 6 percent in 1988 to 4 percent in 1998. Use levels for most other modern and traditional methods were almost unchanged in 1998 from the levels reported in 1988 and 1993. Thus, the overall pattern of change in contraceptive use in Turkey between 1988 and 1998 was an increasing shift to the use of modern methods.

Table 4.7 shows the recent trend in contraceptive use by urban-rural residence and region. Both urban and rural areas in Turkey have shared the increase in use of modern methods. The use of traditional methods appears to have declined among both urban and rural women; albeit in a somewhat more pronounced way in rural areas. Overall, the change in the period from 1993 to 1998 seems to have been characterized by an increase in the overall contraceptive use in rural areas, and a decline in the use of traditional methods, especially in the urban areas.

Table 4.6 Trends in current use of contraception

Percent distribution of currently married women by contraceptive method currently used, 1988 TPHS, 1993 TDHS and 1998 TDHS

Contraceptive method	TPHS-88	TDHS-93	TDHS-98
Any method	63.4	62.6	63.9
Any modern method	31.0	34.5	37.7
Pill	6.2	4.9	4.4
IUD	14.0	18.8	19.8
Condom	7.2	6.6	8.2
Female sterilisation	1.7	2.9	4.2
Other modern methods	2.0	1.3	1.1
Any traditional method	32.3	28.1	25.5
Periodic abstinence	3.5	1.0	1.1
Withdrawal	25.7	26.2	24.4
Other methods	3.1	0.9	0.6
Not currently using	36.6	37.4	36.1

Table 4.7 Trends in current use of contraception by residence and region

Percentage of currently married women 15-49 currently using any method, a modern method, or a traditional method, by residence and region, 1993 TDHS and 1998 TDHS

Residence/ Region	TDHS-93			TDHS-98		
	Any method	Any modern method	Any traditional method	Any method	Any modern method	Any traditional method
Residence						
Urban	66.2	38.9	27.3	66.7	40.8	25.2
Rural	56.1	26.8	29.3	58.1	31.4	26.0
Region						
West	71.5	37.3	34.2	70.5	40.5	29.2
South	62.8	36.7	26.0	60.3	35.1	24.6
Central	62.7	36.6	26.1	68.3	42.8	24.7
North	64.2	29.8	34.4	67.0	35.2	31.5
East	42.3	26.3	16.0	42.0	26.7	15.2
Total	62.6	34.5	28.1	63.9	37.7	25.5

The trends are mixed by region. Overall contraceptive use is found to have increased between 1993 and 1998 in the Central region (from 63 to 68 percent) and in the North (from 64 to 67 percent). Most of this increase can be attributed to the increases in the use of modern methods. Contraceptive use decreased slightly in the West and South between 1993 and 1998. In the East, both the proportion of currently married women using contraception and the level of use of modern methods remained unchanged between 1993 and 1998.

4.5 Number of Children at First Use of Contraception

Family planning methods may be used by couples for either spacing births or limiting family size. To explore the possible motivation for use of contraceptives, a question was asked on the number of

children the respondent had when contraception was first used. These data enable an examination of the cohort changes in the timing of adopting contraceptive use. Table 4.8 shows the distribution of ever-married women by age and the number of children the woman had when she first used contraception. More than one-third of women started using contraception after they had one child. Younger cohorts of women reported first use at lower parities than older cohorts of women. For example, the oldest cohort (age 45-49) of ever-married women first used after having 3.4 births on average, while younger cohorts began use on average before having one child. From another perspective, 10 percent of the age cohort 25-29 cohort of ever-married women started contracepting before the birth of their first child, compared with three percent of the age 45-49 cohort.

Table 4.8 Number of children at first use of contraception

Percent distribution of ever-married women by number of living children at the time of first use of contraception, and median number of children at first use, according to current age, Turkey 1998

Current age	Never used contraception	Number of children at time of first use of contraception						Total	Number of women	Median number of children at first use ¹
		0	1	2	3	4+	Missing			
15-19	63.9	20.2	13.5	2.5	0.0	0.0	0.0	100.0	309	0.0
20-24	40.6	18.3	29.1	9.1	1.9	1.1	0.0	100.0	1,007	0.4
25-29	31.5	10.1	28.3	17.8	6.9	5.3	0.1	100.0	1,197	0.9
30-34	29.6	6.0	21.3	14.7	11.2	17.1	0.0	100.0	926	1.5
35-39	30.9	3.1	14.4	10.4	9.8	31.1	0.3	100.0	963	2.7
40-44	38.8	2.1	8.3	6.7	8.2	35.4	0.5	100.0	618	3.7
45-49	47.1	3.0	7.7	7.6	5.9	28.6	0.0	100.0	489	3.4
Total	36.8	8.7	19.9	11.4	6.9	16.2	0.1	100.0	5,509	1.3

¹ Among ever-married women who have ever used contraception

4.6 Knowledge of the Fertile Period

A basic knowledge of reproductive physiology is necessary for successful practice of coital-related methods, such as withdrawal, condom, and vaginal methods (diaphragm, foam or jelly). Knowledge is particularly critical in the case of periodic abstinence or the rhythm method. The successful practice of periodic abstinence depends on an understanding of when during the ovulatory cycle a woman is most likely to conceive. Table 4.9 presents the percent distribution of all respondents and those who have ever used periodic abstinence and withdrawal by reported knowledge of the fertile period in the ovulatory cycle.

Overall, women in Turkey do not have adequate knowledge of the timing of the ovulation. Only 18 percent of all women know the correct time of ovulation, 44 percent have no idea as to the time, and 38 percent have incorrect knowledge (Figure 4.3). Women who have ever used the rhythm method have better knowledge than all women; 55 percent know the correct time of ovulation, nine percent report that they do not know about the time of ovulation, and 36 percent have incorrect knowledge. Ever users of withdrawal are less likely to be able to identify the time in the cycle when a woman is most likely to conceive; 21 percent know the correct time of ovulation, 36 percent are unsure when a woman is fertile, and 43 have incorrect knowledge.


Table 4.9 Knowledge of fertile period

Percent distribution of all women, women who have ever used periodic abstinence, and women who have ever used withdrawal, by knowledge of the fertile period during the ovulatory cycle, Turkey 1998

Perceived fertile period	All women	Ever users of periodic abstinence	Ever users of withdrawal
During period	0.3	0.3	0.3
After period ends	9.3	12.7	12.6
Middle of the cycle	17.6	54.8	21.3
Before period begins	1.5	1.3	1.3
At any time	22.0	9.1	21.3
Other	5.5	13.2	7.2
Don't know/Missing	43.8	8.5	36.0
Total	100.0	100.0	100.0
Number	8,576	503	3,552

Figure 4.3

Knowledge of Fertile Period Among All Women



TDHS 1998

4.7 Timing of Female Sterilisation

Table 4.10 shows the distribution of sterilised women by their age at the time of sterilisation. These findings should, however, be treated with caution since the number of women sterilized is small and misreporting of ages can distort the results. The results indicate that around two-thirds of women who are sterilised had the operation between age 25 and 34. The median reported age at sterilisation was 32 years, as was found in 1993 TDHS.

Age at time of sterilisation	Percentage of sterilised women
<25	6.2
25-29	27.7
30-34	36.4
35-39	22.1
40-44	6.1
45-49	1.5
Total	100.0
Number of women	257
Median age ¹	31.7

4.8 Sources for Family Planning Methods

Information on sources of modern contraceptives is useful for family planning programme managers and implementors. In the TDHS, women who reported using a modern method of contraception at the time of the survey were asked where they obtained the method last time (Table 4.11). In interpreting the results, it is important to note that some of the women may misreport the type of the place where they obtained the method, since the distinction between hospitals and clinics, or between public and private sources may not be clear to them.

In general, the dominance of the public sector sources in the provision of modern contraceptive services is evident. Overall, 56 percent of users of modern methods said that they relied on a public sector provider. Specifically, health centres (primary health care units) are the most frequent source from where women obtain methods (Figure 4.4). In the case of the pill and the condom, the majority of users go to the private sector for supplies, in particular to pharmacies. In the case of the IUD, most users obtained the IUD from the public institutions, but private doctors are also important providers of the IUD; more than one in four IUD users obtained the method from a private sector provider.

Table 4.12 compares the distribution of users of selected methods by the service provider reported in the TDHS-98 with the distribution reported in the TDHS-93. Looking at all methods, the major change is a somewhat decreased reliance upon public sector in the case of female sterilisation. The percentage of women obtaining sterilisation from a private sector provider has increased from 16 in 1993 to 21 percent in 1998.

Among non-users of contraception, when they were asked about whether they know a source for obtaining modern contraceptives, 56 percent knew about a public sector source, 20 percent cited a private sector source, and 24 percent stated did not know about any source (data not shown).

Table 4.12 Source of supply for selected modern methods, 1993 and 1998

Percent distribution of current users of the pill, IUD, condom, and female sterilisation, by source of supply, 1993 TDHS and 1998 TDHS

Source of supply	Pill		IUD		Condom		Female sterilisation	
	TDHS-93	TDHS-98	TDHS-93	TDHS-98	TDHS-93	TDHS-98	TDHS-93	TDHS-98
Public sector	24.2	26.0	70.9	71.8	28.7	27.7	83.4	76.9
Private sector	75.3	73.6	28.1	27.5	66.2	66.8	15.5	20.8
Other	0.5	0.4	1.0	0.7	5.1	5.5	1.1	2.3

4.9 Discontinuation of Contraceptive Use

Couples can realise their reproductive goals only when they use contraceptive methods consistently. Therefore, a particular concern for family planning programmes is the rate at which users discontinue use of contraception and the reasons for such discontinuation. In the TDHS calendar, all segments of contraceptive use between January 1993 and the date of interview were recorded along with reasons for any discontinuation of use during the period. The discontinuation rates presented here refer only to episodes of contraceptive use that begun during the period of time covered by the calendar, not all episodes that occurred during this period. Specifically, the rates presented in Table 4.13 refer to the 60 month period, 3-63 months prior to the survey. The month of interview and the 2 prior months are ignored in order to avoid the bias that may be introduced by unrecognized pregnancies.

One-year contraceptive discontinuation rates based on the information collected in the TDHS calendar are presented in Table 4.13, according to specific methods. The results indicate that one in three family planning users in Turkey stops using a contraceptive method within 12 months of starting use. The one-year discontinuation rate is as low as nine percent for IUD, while it is 52 percent for injections, 56 percent for the pill, and 38 percent for withdrawal.

Five percent of users stopped using because they want to become pregnant, 7 percent stopped as a result of method failure, another five percent stopped due to side effects or health concerns, and the remaining 17 percent stopped due to other reasons. Side effects or health concerns account for a large portion of the relatively high discontinuation rates for the pill and injection (21 and 29 percent, respectively). On the other hand, method failure accounted for a substantial portion of the discontinuation rates of withdrawal (12 percent).

Table 4.13 Contraceptive discontinuation rates

First-year contraceptive discontinuation rates by reason for discontinuation, according to method, Turkey 1998

Contraceptive method	Reason for discontinuation				
	Method failure	To become pregnant	Side effects/ health concerns	Other reason	All reasons
Pill	6.6	4.8	21.4	23.6	56.4
IUD	0.9	0.4	6.0	2.0	9.4
Condom	6.0	8.1	0.8	28.6	43.3
Withdrawal	12.0	7.2	0.3	18.7	38.2
Total	7.4	5.2	5.0	17.2	34.9

Table 4.14 presents the distribution of all discontinuations during the five-year period before the survey according to the reason for discontinuation and the method used. The desire to become pregnant accounted for one-fifth of all discontinuations. Side effects and health concerns were frequently mentioned as reasons for discontinuation of modern methods. Accidental pregnancy was more common among withdrawal users (33 percent) than modern method users. However, 16 percent of condom discontinuations and 13 percent of pill discontinuations also were due to method failure.

Table 4.14 Reasons for discontinuation of contraception

Percent distribution of contraceptive method discontinuations in the five years preceding the survey by main reason for discontinuation, according to specific methods, Turkey 1998

Reason for discontinuation	Contraceptive method					
	Pill	IUD	Diaphragm/ Foam/ Jelly	Condom	With- drawal	All methods ¹
Became pregnant	12.7	5.6	15.9	15.6	33.1	21.2
To become pregnant	13.3	21.4	9.2	22.5	21.4	19.9
Husband disapproved	0.3	0.1	3.8	12.6	2.6	3.0
Side effects	29.5	24.7	7.1	1.0	0.3	10.7
Health concerns	9.1	16.4	7.0	1.5	0.4	5.5
Access/Availability	0.7	0.0	2.3	2.2	0.0	0.5
More effective method	2.3	0.5	12.3	10.7	13.0	8.3
Inconvenient to use	1.2	0.6	1.8	4.7	1.0	1.4
Infrequent sex	6.4	1.7	0.0	2.1	3.1	3.1
Cost	0.5	0.0	0.8	0.5	0.0	0.2
Fatalistic	0.0	0.1	0.0	0.3	0.0	0.0
Menopause	1.9	1.7	2.3	0.9	3.0	2.3
Marital dissolution	0.7	2.8	1.1	1.2	0.8	1.3
Other	11.4	17.0	25.4	10.6	4.1	9.2
Don't know	0.1	0.0	0.0	0.4	0.2	0.1
Missing	10.0	7.2	11.1	13.3	17.1	13.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Number	729	836	89	566	1,871	4,229

¹ Includes discontinuations of injection, periodic abstinence, and folk methods

4.10 Future Use of Family Planning

An important indicator of the demand for family planning is the extent to which nonusers plan to use family planning in the future. Women who were not using contraception at the time of the survey were asked about their intention to use family planning in the future. The results are presented in Table 4.15. Around half of the currently married women who are not using any contraception report that they intend to use family planning in the future and 36 percent plan to use in the next 12 months (Figure 4.5).


Table 4.15 Future use of contraception						
Percent distribution of currently married women and husbands who are not using a contraceptive method by intention to use in the future, according to number of living children, Turkey 1998						
Future use of contraception	Number of living children ¹					Total
	0	1	2	3	4+	
CURRENTLY MARRIED WOMEN						
Intend use in next 12 months	17.1	48.1	45.9	38.5	25.0	35.3
Intend to use later	43.0	18.0	8.7	4.8	4.5	14.7
Unsure as to timing	1.3	3.1	1.7	0.2	0.2	1.4
Unsure as to intention	5.5	4.1	3.4	2.6	2.7	3.6
Do not intend to use	32.6	25.3	39.3	52.6	65.7	43.7
Don't know/Missing	0.4	1.4	1.0	1.3	1.8	1.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Number	345	488	460	291	556	2,139
HUSBANDS						
Intend use in next 12 months	26.9	37.4	23.2	22.6	22.3	26.0
Intend to use later	27.8	22.6	7.1	3.8	4.1	12.2
Unsure as to timing	3.8	0.2	1.8	0.0	0.6	1.2
Unsure as to intention	4.7	5.9	5.1	1.4	2.7	3.9
Do not intend to use	35.4	31.6	56.5	67.0	67.8	53.4
Don't know/Missing	1.4	2.3	6.2	5.2	2.5	3.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Number	139	126	128	106	239	737

¹ Includes current pregnancy

The proportion intending to use, and especially the timing of the intended use, varies with the number of living children. While the proportion who intend to use within the next 12 months is 17 percent among childless nonusers, almost half of the women with 1 or 2 children plan to use soon. Among husbands, the proportion who intend to use family planning is 39 percent. The proportion intending to use among husbands also varies with number of living children, peaking at 60 percent among those with one child.

Figure 4.5

Future Use of Contraception Among Non-User Women



TNSA 1998

Table 4.16 presents the main reasons for not using family planning reported by currently married nonusers who do not intend to use a contraceptive method in the future. A desire for more children accounts for one-third of nonuse among women less than 30 years of age. The reasons cited most frequently by nonusers age 30 years and over were related to perceived subfecundity and infecundity or being menopausal. Spouse's opposition or concern about religious prohibitions were mentioned as reasons for nonuse by less than 10 percent of women.

Table 4.16 Reasons for not using contraception

Percent distribution of women who are not using a contraceptive method and who do not intend to use in the future by main reason for not using, according to age, and percent distribution of husbands who are not using contraception and who do not intend to use by main reason for not using, Turkey 1998

Reason for not using contraception	Age		Total women	Total husbands
	<30	30+		
Infrequent sex	2.1	2.6	2.5	2.9
Menopausal/hysterectomy	0.0	35.9	30.7	24.2
Subfecund/infecund	17.5	30.8	28.9	11.6
Wants more children	36.4	5.1	9.6	9.6
Respondent opposed	7.0	1.6	2.4	3.7
Husband opposed	8.6	2.3	3.2	0.1
Others opposed	0.0	0.1	0.1	0.0
Religious prohibition	5.8	3.1	3.5	5.2
Knows no method	1.0	1.0	1.0	2.4
Knows no source	0.4	0.4	0.4	0.0
Health concerns	4.5	1.7	2.1	4.1
Fear side effects	0.4	0.3	0.3	1.2
Costs too much	1.7	0.3	0.5	0.6
Inconvenient to use	0.0	0.1	0.1	0.0
Interferes with body processes	0.0	0.2	0.2	0.0
Other	9.7	12.2	11.8	22.9
Don't know	3.5	1.7	2.0	7.8
Missing	1.3	0.4	0.5	0.5
Total	100.0	100.0	100.0	100.0
Number	135	800	935	394

One-fourth of all husbands stated that they were not using contraception because their wives were menopausal or had had a hysterectomy. A further 12 percent were not using because their wives were subfecund or infecund.

Future demand for specific methods of family planning was assessed in the TDHS by asking nonusers who planned to use a family planning method in the future which method they intended to use (Table 4.17). More than one-third of currently married women who are not using a contraceptive method intend to use IUD, down from 51 percent in the TDHS-93. Proportions intending to use other modern methods generally increased. Women who intend to use family planning in the next 12 months and those planning to use later have similar preferences (data not shown).

Table 4.17 Preferred method of contraception for future use

Percent distribution of currently married women who are not using a contraceptive method but who intend to use in the future by preferred method, Turkey 1998

Preferred method of contraception	Total
Pill	10.1
IUD	36.5
Injection	8.5
Diaphragm/Foam/Jelly	0.5
Condom	5.7
Female sterilisation	6.4
Male sterilisation	0.4
Implants	1.1
Periodic abstinence	0.3
Withdrawal	9.8
Folk method	5.1
Don't know/Missing	15.6
Total	100.0
Number	1,101

4.11 Exposure to Family Planning Messages in the Electronic Media

Information on the level of public exposure to a particular type of media allows policy-makers to ensure the use of the most effective media for reaching various target groups. Radio and television are major potential sources of information about family planning. To assess the effectiveness of these electronic media for the dissemination of family planning information, all women and husbands in the survey were asked if they had heard messages about family planning on radio or seen them on television during the last few months preceding the survey.

Table 4.18 shows that around half of all women and husbands reported that they had heard or seen a family planning message on radio *or* television. On the other hand, 10 percent of women and 14 percent of husbands reported that they had heard or seen a family planning message on radio *and* television. Overall, family planning messages are seen more often on television rather than heard on radio. The differences in the percentages who had seen a message by type of place of residence, region, and education are greater for women than for men. For both women and men, sharp contrasts in access to media messages are observed between less educated and educated respondents.

Table 4.18 Heard about family planning on radio and television

Percent distribution of all women and of husbands by whether they heard a radio and/or television message about family planning in the few months prior to the interview, according to selected background characteristics, Turkey 1998

Background characteristic	Heard on both radio and TV	Radio only	Television only	Heard on neither	Missing	Total	Number of women/husbands
CURRENTLY MARRIED WOMEN							
Age							
15-19	8.0	1.1	34.6	56.3	0.0	100.0	1,720
20-24	9.1	1.3	43.0	46.3	0.2	100.0	1,558
25-29	9.4	0.6	42.9	47.1	0.0	100.0	1,397
30-34	10.4	0.5	41.4	47.7	0.0	100.0	1,202
35-39	11.5	0.7	34.9	52.8	0.0	100.0	1,081
40-44	10.9	0.4	34.5	54.2	0.0	100.0	885
45-49	12.4	0.8	29.2	57.5	0.1	100.0	733
Residence							
Urban	10.0	0.7	41.1	48.1	0.1	100.0	5,704
Rural	9.8	1.0	31.8	57.3	0.1	100.0	2,872
Region							
West	9.2	0.9	41.7	48.1	0.1	100.0	3,204
South	10.0	0.4	34.9	54.6	0.0	100.0	1,258
Central	10.0	0.8	40.9	48.1	0.1	100.0	1,985
North	7.8	0.9	40.6	50.6	0.1	100.0	692
East	12.0	0.9	27.2	59.9	0.0	100.0	1,437
Education							
No educ./Pri. incomp.	10.0	0.5	24.2	65.1	0.2	100.0	1,861
Pri. comp./Sec. incomp.	9.4	1.0	38.7	51.0	0.0	100.0	5,158
Sec. comp./+	11.5	0.7	52.4	35.3	0.1	100.0	1,556
All women	9.9	0.8	38.0	51.2	0.1	100.0	8,576
HUSBANDS							
Age							
15-24	12.3	4.2	51.1	32.6	0.0	100.0	113
25-29	9.7	1.2	41.4	47.4	0.4	100.0	342
30-34	9.5	0.1	42.4	47.8	0.2	100.0	364
35-39	14.2	0.8	33.3	51.1	0.8	100.0	352
40-44	15.3	0.6	33.7	50.4	0.0	100.0	335
45-49	16.3	1.3	33.8	48.6	0.0	100.0	240
50+	22.4	0.0	26.9	47.9	2.6	100.0	226
Residence							
Urban	13.3	1.1	38.1	47.0	0.4	100.0	1,347
Rural	14.8	0.2	34.0	50.9	0.0	100.0	624
Region							
West	12.9	1.4	38.2	47.4	0.2	100.0	767
South	17.4	1.1	33.5	47.8	0.2	100.0	285
Central	11.8	0.1	39.6	47.9	0.6	100.0	481
North	7.6	0.9	38.8	52.4	0.2	100.0	150
East	19.5	0.3	30.8	49.4	0.0	100.0	287
Education							
No educ./Pri. incomp.	18.6	1.1	25.9	54.0	0.3	100.0	204
Pri. comp./Sec. incomp.	14.4	0.6	34.1	50.5	0.3	100.0	1,253
Sec. comp./+	10.5	1.3	47.8	40.4	0.0	100.0	513
All men	13.8	0.8	36.8	48.3	0.3	100.0	1,971

4.12 Acceptability of Use of Electronic Media to Disseminate Family Planning Messages

To determine the level of acceptance of the dissemination of family planning information through the media, the TDHS asked women and husbands whether it was acceptable to disseminate family planning information on radio and television. Overall, the majority of women (89 percent) and men (87 percent) interviewed reported that it was acceptable to use radio or television to air family planning messages (Table 4.19). Acceptability of radio and television as a source of information is highest among women and husbands in urban areas, in the West, and among those with high levels of education.

Table 4.19 Acceptability of family planning messages on radio and television

Percentage of women and husbands who think it is acceptable to have messages about family planning on radio and television, by selected background characteristics, Turkey 1998

Background characteristic	Women		Husbands	
	Percent	Number	Percent	Number
Age				
15-19	86.2	1,720	*	4
20-24	90.1	1,558	80.9	109
25-29	91.2	1,397	88.1	342
30-34	91.2	1,202	87.0	364
35-39	89.8	1,081	90.0	352
40-44	88.6	885	89.0	335
45-49	85.0	733	87.1	240
50+	NA	NA	76.7	226
Residence				
Urban	91.5	5,704	88.9	1,347
Rural	84.1	2,872	81.8	624
Region				
West	92.6	3,204	92.8	767
South	90.2	1,258	84.4	285
Central	92.1	1,985	85.3	481
North	87.0	692	85.8	150
East	76.7	1,437	75.0	287
Education				
No educ./Pri. incomp.	74.8	1,861	67.1	204
Pri. comp./Sec. incomp.	91.8	5,158	86.6	1,253
Sec. comp./+	97.0	1,556	94.5	513
Total	89.0	8,576	86.6	1,971

NA = Not applicable
 Note: An asterisk indicates a figure is based on fewer than 25 respondents and has been suppressed.

4.13 Exposure to Family Planning Messages in Print Media

Women were asked if they had been exposed to a family planning message through a newspaper/magazine article, a poster, or a leaflet/brochure (i.e., through print media) during the few months prior to the interview. The results are presented in Table 4.20. Twenty-seven percent of the women interviewed reported that they had exposed to at least one of these media that contained family planning information. The most commonly reported source was newspaper/magazine (19 percent). Women in older age groups, living in the East, or who were less educated were less likely to have been exposed to print media on family planning than those in younger age cohorts, living in urban areas or in the West and more educated women.

Table 4.20. Heard about of family planning through print media

Percentage of women and husbands who had seen a message about family planning through print media in the few months prior to the interview, by source of message and selected background characteristics, Turkey 1998

Background characteristic	Print media				Number of women/husbands
	Any source	Newspaper/Magazine	Poster	Leaflet/Brochure	
CURRENTLY MARRIED WOMEN					
Age					
15-19	27.2	19.7	8.5	10.3	1,720
20-24	31.6	22.6	13.6	12.4	1,558
25-29	29.7	21.0	13.2	10.8	1,397
30-34	27.2	19.7	12.1	9.6	1,202
35-39	25.0	17.3	10.2	9.1	1,081
40-44	21.5	15.1	10.4	9.2	885
45-49	17.1	13.5	6.1	6.2	733
Residence					
Urban	31.7	23.1	13.0	12.2	5,704
Rural	16.7	11.2	6.8	5.8	2,872
Region					
West	33.6	23.9	15.3	12.9	3,204
South	22.5	15.8	8.2	8.9	1,258
Central	25.8	18.1	9.8	9.4	1,985
North	24.9	19.0	9.5	9.1	692
East	17.1	12.9	5.7	6.2	1,437
Education					
No educ./Pri. incomp.	7.2	4.4	3.2	2.8	1,861
Pri.comp./Sec. incomp.	25.4	17.5	10.1	8.7	5,158
Sec. comp./+	54.1	42.0	22.9	23.4	1,556
Total	26.7	19.1	10.9	10.1	8,576
HUSBANDS					
Age					
20-24	53.1	42.6	11.7	20.5	112
25-29	54.6	40.0	17.6	21.6	342
30-34	47.4	37.4	20.6	15.8	364
35-39	45.7	40.3	15.4	18.5	352
40-44	45.3	40.4	12.7	17.3	335
45-49	44.5	39.1	10.5	20.0	240
50+	33.8	28.8	5.5	10.0	226
Residence					
Urban	52.1	43.1	15.8	19.6	1,347
Rural	34.2	28.3	11.3	13.5	624
Region					
West	53.2	44.3	16.5	19.8	767
South	41.4	34.5	15.0	20.1	285
Central	47.6	37.8	15.0	17.3	481
North	42.7	36.3	13.4	11.4	150
East	33.5	28.6	7.3	13.6	287
Education					
No education/ Primary incomplete	19.1	14.9	4.2	7.9	204
Complete primary/ Secondary incomplete	43.8	37.0	12.7	15.6	1,357
Secondary complete+	65.7	52.6	23.3	27.8	513
Total	46.4	38.4	14.4	17.7	1,971

Overall, husbands were more likely than women to have seen a family planning message through the print media. Nevertheless, even among husbands, fewer than half had seen a message in any of the print media. Differentials in the likelihood of having seen a message parallel those described for women.

4.14 Attitudes of Couples Toward Family Planning

Contraceptive use is facilitated when couples have a positive attitude towards family planning. Attitudinal data were collected in the TDHS by asking women whether they approve of couples using family planning and what they perceived as their husband's attitude towards family planning. This information is useful in the formulation of family planning policies, since it indicates the extent to which further education and publicity are needed to gain or increase acceptance of family planning. Widespread disapproval of contraception acts as a barrier to adoption of contraceptive methods.

The results presented in Table 4.21 are confined to currently married, nonsterilised women and exclude those who had never heard of a contraceptive method. In general, 91 percent of married women who know of a contraceptive method approve of family planning, and 83 percent believe that their husband approves. Overall, 80 percent of women said that both they and their husbands approve of family planning while only three percent of them reported that both they and their husbands disapprove (Figure 4.6).

Table 4.21 Wives' perceptions of couple's attitude toward family planning


Percent distribution of currently married, nonsterilised women who know of a contraceptive method by wife's attitude toward family planning and wife's perception of her husband's attitude toward family planning, according to selected background characteristics, Turkey 1998

Background characteristic	Woman approves			Woman disapproves				Missing	Total	Wife approves	Husband approves ¹	Number of women
	Both approve	Husband disapproves	Husband's attitude unknown	Both disapprove	Husband approves	Husband's attitude unknown	Wife unsure					
Age												
15-19	71.8	7.8	6.2	2.1	4.1	1.4	5.9	0.7	100.0	86.1	74.8	258
20-24	80.7	6.2	4.1	0.9	2.7	0.5	4.8	0.1	100.0	91.0	82.8	913
25-29	84.1	5.0	4.1	0.7	1.9	0.7	3.4	0.1	100.0	93.3	86.0	1,165
30-34	82.4	5.6	3.4	1.5	2.6	0.7	3.6	0.1	100.0	91.6	85.7	1,019
35-39	80.1	5.7	5.2	0.9	4.1	0.3	3.6	0.1	100.0	91.1	82.2	932
40-44	77.2	5.5	5.8	1.6	3.7	0.8	5.5	0.0	100.0	88.5	80.4	725
45-49	73.2	5.6	7.2	0.5	5.1	1.7	6.6	0.1	100.0	86.1	76.0	593
Residence												
Urban	83.6	4.7	4.2	0.8	2.6	0.4	3.6	0.1	100.0	92.6	85.8	3,762
Rural	72.6	7.7	6.1	1.7	4.4	1.6	6.0	0.1	100.0	86.4	75.8	1,843
Region												
West	88.6	3.8	3.4	0.4	1.2	0.4	2.1	0.0	100.0	95.8	90.0	2,148
South	75.6	5.7	5.8	1.4	3.3	0.9	6.8	0.3	100.0	87.5	79.3	806
Central	80.5	6.7	5.8	1.1	1.6	0.5	3.5	0.2	100.0	93.1	82.7	1,378
North	74.4	7.0	6.4	1.4	2.9	1.2	6.6	0.0	100.0	87.9	78.5	434
East	64.1	7.9	4.9	2.0	10.9	1.7	8.4	0.1	100.0	77.0	67.9	839
Education												
No educ./ Pri. comp.	61.6	9.0	6.4	2.1	9.2	1.9	9.5	0.1	100.0	77.1	66.4	1,410
Pri. comp./Sec.incomp.	83.9	5.2	4.9	0.8	1.4	0.4	3.2	0.1	100.0	94.1	86.0	3,426
Sec. comp./+	96.3	1.8	1.3	0.0	0.0	0.0	0.3	0.2	100.0	99.4	96.4	768
Total	80.0	5.7	4.8	1.1	3.2	0.8	4.4	0.1	100.0	90.5	82.5	5,605

¹ Includes women who are unsure about their own attitude, but know their husband's attitude

Figure 4.6

Attitudes Toward Family Planning as Reported by Couples



TNSA 1998

The likelihood that a woman will report that both she and her husband approve of family planning is higher among women age 25-34 and declines slightly with increasing age. Urban women were more likely to report that both they and their husbands approve of family planning than those in rural areas (84 and 73 percent, respectively). Approval was highest in the West region (89 percent) while it was lowest in the East (64 percent). More educated women are also more likely to say that their spouses approve family planning.

The fact that both women and husbands in the same household were interviewed provided an opportunity to link responses obtained from currently married women with those obtained independently from their husbands. A total of 1,896 couples were linked in this way. Table 4.22 shows the percent distribution of these couples by both spouses' approval of family planning, according to the difference in age and educational level between husbands and wives. In the case of 78 percent of couples, both spouses reported that they approve of family planning. In the case of only 2 percent of the couples did both disapprove. When only one spouse disapproved, it was more likely to be the husband than wife (9 percent and 2 percent, respectively). Generally, as the age difference between husband and wife increases so does the likelihood that one spouse or the other disapproves family planning. Couples are less likely to disapprove family planning when both spouses are educated.

Table 4.22 Attitudes of couples toward family planning

Percent distribution of couples by approval of family planning, according to age and education differences between spouses, Turkey 1998

Background characteristic	Both approve	Both disapprove	Wife approves, husband disapproves	Husband approves, wife disapproves	Unsure/ Missing	Total	Percentage of couples in agreement	Number of couples
Age difference								
Husband younger than wife	68.7	1.2	12.4	4.9	12.8	100.0	69.9	186
Husband older by:								
0-4 years	81.7	1.6	7.5	1.7	7.5	100.0	83.3	848
5-9 years	80.2	1.6	5.6	1.7	10.8	100.0	81.8	625
10-14 years	71.1	4.7	12.7	2.4	9.1	100.0	75.8	183
15 years +	53.9	6.6	29.8	1.2	8.5	100.0	60.5	55
Education difference								
Husband and wife not educated	33.8	10.6	17.1	9.7	28.8	100.0	44.4	93
Wife educated, husband not	(43.5)	(5.3)	(24.3)	(5.5)	(21.3)	100.0	48.9	32
Husband educated, wife not	59.5	6.5	10.5	3.2	20.3	100.0	66.0	262
Husband and wife educated	84.8	0.6	7.3	1.3	6.0	100.0	85.4	1,509
Total	78.1	2.0	8.5	2.1	9.3	100.0	80.1	1,896

Note: Figures in parentheses are based on 25-49 cases.

CHAPTER 5

ABORTIONS AND STILLBIRTHS

Banu Akadlı Ergöçmen and Turgay Ünalán

This chapter presents information on induced abortions, spontaneous abortions, and stillbirths. Stillbirths and spontaneous abortions are important maternal health indicators. Induced abortions have additional significance for family planning services since one important goal of family planning is to eliminate unwanted pregnancies.

Women resort to induced abortions when they lack access to contraceptive services or other psychosocial barriers keep them from using family planning to avoid unwanted pregnancies. They may also decide to abort a pregnancy that results from contraceptive failure.

Legalising abortion provides safe conditions to terminate unwanted pregnancies. In Turkey, the practice of abortion was legalised in 1983 with the enactment of the new population planning law. This law provided safe abortion on request during the first ten weeks of gestation for every woman who needed the service. Since the law was enacted, induced abortion has been available to women at government hospitals for a nominal fee and from the private sector for a fee.

In the TDHS-98, questions were included to determine the total number of induced and spontaneous abortions and stillbirths. Information on the duration, the month and year in which the pregnancy was terminated and the person who assisted the woman at the time of abortion was collected. In addition, information was collected on the reason for the last abortion.

5.1 Life-time Experience of Women

Table 5.1 shows the distribution of ever-married women by the total number of abortions (induced and spontaneous) and stillbirths that they reported ever having had during their reproductive lives. Overall, the table shows that relatively few women (5 percent) have had a still birth. Slightly less than one-quarter of ever-married women reported ever having had a spontaneous abortion. The proportion of reporting that they had ever had an induced abortion was somewhat higher; a total of 27 percent of ever-married women have ever had an induced abortion. Among the women who had had an induced abortion, around half had had only one induced abortion.

Both age and the number of living children are strongly associated with the likelihood that a woman will have had an induced abortion. Table 5.2 shows that the percentage of ever-married women ever having had an induced abortion varies from less than 10 percent among women under age 25 to more than 40 percent among women age 45-49. The percentage of ever-married women who have had an induced abortion also increases steadily with the number of living children, from five percent of women with no living children to a peak of 41 percent among women with four children.

Urban women are more likely to report having had an induced abortion than rural women. Considering regional patterns, women in the East region are the least likely to have ever had an induced abortion and women in the West region are the most likely to report an induced abortion. There is little variation by education in the proportions of ever-married women who have had an induced abortion.

Table 5.1 Number of abortions and stillbirths

Percent distribution of ever-married women by number of abortions (spontaneous and induced) and stillbirths, Turkey 1998

Number of pregnancies/terminations	Abortions		Stillbirth
	Spontaneous	Induced	
None	78.6	73.3	95.0
1	15.2	14.2	4.3
2	3.8	7.2	0.5
3	1.4	3.0	0.1
4	0.5	1.2	0.1
5 or more	0.4	1.1	0.0
Missing	0.1	0.0	0.0
Total	100.0	100.0	100.0
Mean number	0.3	0.5	0.1
Number of ever-married women	6,196	6,196	6,196

Table 5.2 Induced abortions by background characteristics

Percentage of ever-married women ever having an induced abortion, by selected background characteristics, Turkey 1998

Background characteristic	Ever had induced abortion	Number of women
Age		
15-19	5.5	266
20-24	7.3	945
25-29	19.9	1,216
30-34	26.6	1,124
35-39	37.3	1,055
40-44	37.9	869
45-49	42.4	721
Number of living children		
None	5.4	638
1	14.8	1,189
2	30.1	1,880
3	34.8	1,141
4	41.4	595
5 or more	31.0	754
Residence		
Urban	29.6	4,182
Rural	20.5	2,015
Region		
West	30.7	2,385
South	24.0	892
Central	27.3	1,483
North	26.5	494
East	18.2	942
Education		
No educ./Pri. incomp	26.7	1,635
Pri. comp./Sec. incomp.	26.4	3,698
Sec. comp./+	27.8	863
Total	26.7	6,196

5.2 Current Levels and Trends

The data collected in the TDHS-98 can be used to look at the current levels of abortions and stillbirths. Table 5.3 shows the rates of abortions (spontaneous and induced) and still births per 100 pregnancies as reported by ever-married women in the TDHS-98. The table shows that, during the five-year period before the survey, almost one in four pregnancies terminated in other than a live birth. As expected, most non-live terminations were abortions. Only about 2 out of every 100 pregnancies ended in a still birth. There were 23 abortions per 100 pregnancies, of which 15 were induced.

Outcome	Number per 100 pregnancies
Abortions	23.2
Spontaneous	8.7
Induced	14.5
Stillbirths	1.5

Table 5.4 looks at the trends in the levels of induced abortions during the period between the TDHS-93 and the TDHS-98. Overall, the results suggest that there was a slight decline in induced abortions during the period, from 18 abortions to 15 abortions per 100 pregnancies. The level of induced abortions also declined for most subgroups. Among urban women, for example, there were 21 induced abortions per 100 pregnancies during the five-year period prior to the TDHS-93. At the time of the TDHS-98, the level of induced abortions among urban women had dropped to 16 per 100 pregnancies.

Background characteristic	1993 DHS	1998 TDHS
Age		
15-19	3.8	5.8
20-24	8.3	7.7
25-29	20.4	12.6
30-34	27.9	23.3
35-39	36.2	33.4
40-44	47.1	42.5
45-49	47.6	66.2
Residence		
Urban	21.3	16.1
Rural	12.4	11.6
Region		
West	24.9	18.0
South	16.3	13.7
Central	19.8	16.7
North	17.0	15.6
East	8.7	7.6
Education		
No educ./Pri. incomp.	13.9	11.8
Pri. comp./Sec. incomp.	19.4	15.1
Sec. comp./+	22.6	17.3
Total	18.0	14.5

5.3 Contraceptive Use Before and After Induced Abortions

One of the issues that the TDHS calendar can be used to examine is the use of contraception of women before and after an abortion. An examination of the patterns of contraceptive use before a woman has an abortion is important because pregnancies that end in abortions are often result from the use of ineffective methods or from the failure to use methods effectively as well as from the failure to use contraception at all. According to the TDHS-98 results, all of these factors are related to abortion in Turkey. As Table 5.5 shows, in around one-third of the abortions during the five-year period before the survey, the woman was not using any form of contraception in the month before she became pregnant. Among abortions in which contraception was used immediately prior to the pregnancy, women were more likely to be using a traditional than a modern method. More than four in ten abortions occurred following a period of use of withdrawal, while 22 percent occurred after the woman became pregnant while using a modern method, primarily the condom (9 percent) or the pill (7 percent).

Method	Percentage using method before last abortion
Pill	7.0
IUD	5.2
Injection	0.2
Diaphragm/Foam/Jelly	1.1
Condom	8.5
Periodic abstinence	2.2
Withdrawal	41.5
Other	1.0
No method	33.4
Total	100.0
Number	608

Information on the use of contraception in the month following an abortion is presented in Table 5.6. Women did not use any contraceptive method following abortion in the case of almost one-third of the abortions in the five-year period before the survey, and they initiated use of a traditional method, principally withdrawal, following a similar percentage of abortions. Modern family planning methods were adopted in the month following the pregnancy termination in the case of 35 percent of abortions during the five-year period before the survey. In comparison with the mix of method used before abortion, this represented a substantial increase in the use of modern methods. The IUD was the most popular method among women adopting a modern method in the month after an abortion, followed by the pill and condom.

Table 5.6 Method used after abortion

Method used within one month after pregnancy for the last abortion reported in the five years preceding the survey, Turkey 1998

Method	Percentage using method after last abortion
Pill	9.4
IUD	14.9
Injection	0.4
Diaphragm/Foam/Jelly	2.3
Condom	8.4
Female sterilization	1.1
Periodic abstinence	0.9
Withdrawal	27.4
Other	1.7
No method	32.1
Total	100.0
Number	608

5.4 Reasons for Induced Abortion

The reasons women gave for having their last abortion are presented in Table 5.7. Six of every ten women had the abortion because they did not want to have another child. An additional seven percent ended the pregnancy because it followed a previous pregnancy too closely. Thus, a total of 69 percent of women who had an abortion wanted to space or limit their births at the time of the abortion. Among women citing other reasons for the abortion, the majority indicated that they ended the pregnancy because of concerns about their health.

As the age of women increases, the proportion of women had an abortion because they did not want another child also increases. The desire to delay (space) a wanted birth or concerns about health were cited somewhat more frequently as the reasons for ending the pregnancy among women under age 35 than among older women.

There is little variation between urban and rural areas in the reasons women gave for having an abortion, with nearly two-thirds of both urban and rural women saying that they had an abortion because they did not want another child. In all of the regions, the predominant reason for having an abortion was also a desire to limit births. The West Region has the highest percentage of women reporting they had the abortion because they did not want another child (68 percent), and the East region had the lowest percentage (52 percent). In the East, one-fifth of women had the abortion because of concerns related to their own health, which is more than twice the percentage expressing this concern in the West.

As expected, not wanting another child is more frequently reported by the less-educated women. However, even among more highly educated women, a majority of women reported the last abortion was the result of a desire to limit births. Women with secondary or higher education were somewhat more likely than women in the other education categories to say that they had terminated the pregnancy because it was too close to a prior pregnancy. On the other hand, the likelihood that a women had an abortion because of concern about her own health was somewhat greater for women with with no education (17 percent) than for women with a secondary or higher education (11 percent).

Table 5.7. Reasons for induced abortion

Percent distribution of women who ever had an induced abortion by reason for last induced abortion, according to background characteristics, Turkey 1998

Background characteristic	Reason for induced abortion						Total	Number of women
	Health related to:			Didn't want another	Previous pregnancy just ended	Other		
	Mother	Child	Both					
Age								
15-19	*	*	*	*	*	*	100.0	15
20-24	14.7	10.5	6.6	39.9	13.8	13.3	100.0	65
25-29	18.2	9.4	4.4	43.1	13.3	11.7	100.0	226
30-34	14.4	6.0	1.7	55.6	10.1	11.9	100.0	271
35-39	12.8	1.1	2.0	69.8	4.3	9.6	100.0	368
40-44	13.2	1.2	1.9	70.2	4.1	9.2	100.0	299
45-49	12.1	3.4	0.5	71.0	5.1	8.0	100.0	274
Residence								
Urban	13.2	4.4	1.8	62.0	7.9	10.4	100.0	1,129
Rural	16.1	3.5	3.2	62.0	5.7	9.3	100.0	389
Region								
West	9.6	3.1	1.7	67.9	7.9	9.5	100.0	680
South	17.4	3.3	3.0	56.0	7.7	12.2	100.0	189
Central	16.4	6.1	1.9	57.6	6.1	11.9	100.0	372
North	17.9	2.9	1.7	64.1	7.2	6.3	100.0	125
East	20.0	6.2	4.7	52.2	7.5	8.6	100.0	153
Education								
No educ./Pri. incomp.	17.4	2.5	2.3	66.9	3.6	6.4	100.0	383
Pri. comp./Sec. incomp.	13.6	4.3	2.6	61.7	7.8	9.8	100.0	810
Sec. comp./+	10.8	5.6	1.0	56.9	10.5	15.1	100.0	324
Total	14.0	4.1	2.2	62.0	7.3	10.1	100.0	1,518

Note: An asterisk indicates a figure is based on fewer than 25 respondents and has been suppressed.

5.5 Timing of Induced Abortions

Although abortions are legal for up to 10 weeks of pregnancy (2.5 months), it is safer for a woman to have an abortion as early as possible. Table 5.8 shows the distribution of induced abortions in the five-year period prior to the TDHS by the number of months that the woman was pregnant at the time of the abortion. Overall, 68 percent of the abortions took place in the first month of pregnancy and 23 percent in the second month of pregnancy. Thus, only nine percent of induced abortions were performed outside of the recommended time limit. Women in the South and East regions were more likely to report having an induced abortion when they were three or more months pregnant than women in the other regions.

Table 5.8 Timing of induced abortion

Percent distribution of women with recent induced abortions by number of months pregnant at time of abortion, according to place of residence, Turkey 1998

Residence/ Region	Number of months pregnant			Total	Number of women
	1	2	3+		
Residence					
Urban	68.9	23.0	8.1	100.0	437
Rural	66.4	23.4	10.2	100.0	171
Region					
West	69.2	23.4	7.4	100.0	638
South	62.7	20.1	17.2	100.0	77
Central	70.6	23.4	6.0	100.0	155
North	77.3	17.8	4.8	100.0	53
East	59.5	28.2	12.3	100.0	77
Total	68.2	23.1	8.7	100.0	608

5.6 Provider

Table 5.9 provides information on the abortion provider. Nearly three in four women who had an induced abortion in the five-year period preceding the survey reported the abortion took place at a private doctor's office (59 percent) or at a private hospital or clinic (15 percent). Women seeking abortion services from public sector providers were more likely to report that they had obtained the abortion at a hospital than from a health center.

Rural women were somewhat more likely than urban women to have had the abortion performed at private provider (78 percent and 72 percent, respectively). The proportion obtaining abortion services at a private doctor's office or a private clinic or hospital also varied by region, from 68 percent in the North to 77 percent in the Central region.

Table 5.9 Abortion providers

Percent distribution of women who used induced abortion to terminate pregnancies during the last five years, by place of provision, according to place of residence, Turkey 1998

Residence/ Region	Abortion provider									Total	Number of women
	Government hospital	Maternity hospital	Health center	SSK hospital	Private hospital/ clinic	Doctor's office	Other private	Uni- versity hospital	Missing		
Residence											
Urban	10.4	6.5	0.0	6.3	17.0	53.3	0.0	2.0	2.7	100.0	467
Rural	14.1	3.0	0.5	2.7	8.6	69.6	0.6	0.0	0.9	100.0	189
Region											
West	9.0	4.6	0.0	9.7	24.6	48.4	0.0	1.1	2.7	100.0	259
South	15.3	3.3	0.0	0.8	7.3	66.9	0.0	1.6	4.8	100.0	86
Central	9.4	8.3	0.0	2.5	10.2	66.7	0.0	2.3	0.6	100.0	176
North	12.0	10.7	1.5	6.7	4.2	63.4	0.0	0.6	0.7	100.0	56
East	19.7	0.8	0.0	0.9	6.1	68.4	1.4	0.7	2.1	100.0	80
Total	11.5	5.5	0.1	5.3	14.5	59.4	0.2	1.4	2.2	100.0	657

CHAPTER 6

OTHER PROXIMATE DETERMINANTS OF FERTILITY

Banu Akadlı Ergöçmen and İsmet Koç

This chapter addresses a number of factors other than contraception that affect a woman's risk of becoming pregnant, namely, nuptiality, postpartum amenorrhoea, abstinence from sexual relations, and secondary infertility. The nuptiality data collection procedure in the TDHS-98 differed in a number of ways from the standard DHS questionnaire. In the TDHS-98, the nuptiality questions for ever-married women followed the fertility section rather than being included in a separate section at the end of the questionnaire as is typical in the DHS. For ever-married women, there also were additional questions about family formation, religious marriages, and consanguinity. Husbands but not ever-married women were asked about the age at which they initiated sexual activity while, for cultural reasons, neither the husbands nor ever-married women were asked about recent sexual activity. Finally, in the never-married women questionnaire, the marriage section is composed only of attitude questions.

Although it is by no means always true, marriage is an indicator of exposure of women to the risk of pregnancy; therefore it is important for the understanding of fertility. Populations in which age at marriage is low also tend to experience early childbearing and high fertility. Trends in the age at which women marry can help to explain the trends in fertility levels. Measures of other proximate determinants of fertility including the duration of postpartum amenorrhoea and postpartum abstinence, and the level of secondary infertility are also important in understanding fertility patterns.

6.1 Current Marital Status


Table 6.1 and Figure 6.1 show the distribution of all women age 15-49 by their marital status at the time of the survey. Overall, 69 percent are currently married¹, 2 percent are widowed, 1 percent are divorced, and 28 percent are never married. In Turkey, marriage is still almost universal. By the end of the reproductive years, less than 2 percent of women have never married. The universality of marriage is also evident from the

Age	Marital status					Total	Number of women
	Never married	Married	Widowed	Divorced	Not living together		
15-19	84.5	15.2	0.1	0.0	0.2	100.0	1,720
20-24	39.3	59.3	0.2	0.9	0.3	100.0	1,558
25-29	12.9	85.6	0.4	0.8	0.2	100.0	1,397
30-34	6.5	90.6	1.3	1.3	0.3	100.0	1,202
35-39	2.4	93.8	1.1	1.7	0.9	100.0	1,081
40-44	1.8	89.2	6.1	2.1	0.8	100.0	885
45-49	1.7	87.9	7.7	2.3	0.4	100.0	733
Total	27.7	69.0	1.7	1.1	0.4	100.0	8,576

¹The term married refers both to "currently married" and "currently in union."

fact that among women age 30 and over, 94 percent or more are, or have been, married. The percentage of never-married women declines rapidly with age, decreasing almost by half, from 85 percent among teenagers to 39 percent among women in their early twenties.

As expected, the proportion widowed increases with age, from less than 1 percent of women under age 30 to 8 percent among women age 45-49. The percentage of divorced women is low, and very few women report that they are not living with their husbands.



6.2 Age at First Marriage

In Turkey, marriage is almost universal and almost all births occur within marriage. Therefore, age at first marriage is an important demographic indicator since it represents the beginning of exposure to the risk of pregnancy.

As shown in Table 6.2, the median age at first marriage is 19.5 among women 25-49, indicating that half the women in those age groups married before that age. A steady increase is observed in the median age at first marriage, from 18.4 years for the 45-49 age group to 20.4 years for the 25-29 age group. There has been an especially marked decline in getting married at very young ages. For example, the percentage of women getting married by age 15 has dropped from 12 percent among women age 45-49 to only 4 percent among women 20-24.

Although the median is a convenient summary measure, not all changes in age at marriage are necessarily reflected in the median. Cohort trends in age at marriage can be more thoroughly examined by comparing the percentages who first marry at specific ages for successive 5-year age groups. These percentages reflect a change in age at marriage in Turkey over the past several decades. The percentages of women married at each specific ages are all lower for the younger cohorts than for the older cohorts. For example, among women age 45-49, 66 percent married by age 20, whereas only 46 percent of women age 25-29 married by age 20.

Table 6.2 Age at first marriage

Percentage of women and husbands who were first married by specific exact ages and median age at first marriage, according to current age, Turkey 1998

WOMEN								
Current age	Percentage of women who were first married by exact age:					Percentage who had never married	Number of women	Median age at first marriage
	15	18	20	22	25			
15-19	2.3	NA	NA	NA	NA	84.5	1,720	a
20-24	4.2	23.0	42.8	NA	NA	39.3	1,558	a
25-29	5.8	27.8	45.6	63.0	80.3	12.9	1,397	20.4
30-34	7.2	27.9	47.5	65.7	82.4	6.5	1,202	20.3
35-39	9.4	37.9	60.6	73.6	85.3	2.4	1,081	19.0
40-44	10.8	43.0	66.2	81.9	89.3	1.8	885	18.5
45-49	12.3	45.0	66.4	80.5	91.2	1.7	733	18.4
Women 20-49	7.6	32.1	52.5	67.7	79.3	13.5	6,856	19.7
Women 25-49	8.6	34.8	55.4	71.4	84.8	5.9	5,299	19.5
HUSBANDS								
Current age	Percentage of husbands who were first married by exact age:					Percentage who had never married	Number of husbands	Median age at first marriage
	20	22	25	28	30			
25-29	15.0	28.2	70.1	NA	NA	NA	342	23.5
30-34	12.8	24.0	63.1	85.8	92.6	NA	364	24.0
35-39	21.8	30.3	60.9	80.5	90.0	NA	352	24.0
40-44	25.8	36.7	72.2	87.9	94.5	NA	335	22.9
45-49	28.4	38.0	71.9	88.8	92.1	NA	240	22.9
50-64	16.8	26.1	59.5	79.0	90.0	NA	226	24.2
Husbands 25-64	19.8	30.3	66.3	87.0	93.4	NA	1,859	23.6

NA = Not applicable
^a Omitted because less than 50 percent of the women in the age group x to $x+4$ were first married by age x

Comparison with the data from the husband survey shows that husbands enter into first marriage at a much later age than women (Table 6.2). The median age at first marriage for husbands is almost 24 years, compared with 19.5 years for women. Only 20 percent of husbands were married by age 20, compared with 55 percent of women. The percentages of husbands married at each specific marriage age are lower for the younger age groups and increase up to age 50. For example, among husbands in age group 25-29, 15 percent married by age 20, whereas for the husbands in age group 45-49, 28 percent had married by age 20.

Table 6.3 examines the differences in the median age at first marriage among women age 20-49 and husbands age 25-64 years by residence, region, and education. The overall median ages at first marriage observed for urban and rural women age 25-49 indicate that urban women marry slightly later than their rural counterparts (19.8 and 18.9, respectively). For husbands, the difference in the median age at first marriage is more than one year between urban and rural areas (24.0 and 22.6, respectively).

The greatest variation between the regions is observed between the East and the remaining regions. The lowest median age, 18.1, is found in the East and the highest, 19.9, in the West, indicating that women in the East marry nearly two years earlier than women in the West (Table 6.3). In the other regions, the median age at marriage among women varies from 19.3 in the Central region to 19.6 in the South. The regional pattern

Table 6.3 Median age at first marriage

Median age at first marriage among women age 25-49 years, and among husbands age 25-64, by current age and selected background characteristics, Turkey 1998

WOMEN							
Background characteristic	Current age					Women age 25-49	
	25-29	30-34	35-39	40-44	45-49		
Residence							
Urban	20.6	20.5	19.2	18.8	18.6		19.8
Rural	19.8	19.9	18.7	17.8	18.3		18.9
Region							
West	20.6	20.7	19.4	19.0	18.8		19.9
South	21.0	20.5	19.4	18.1	17.8		19.6
Central	20.4	19.9	18.5	18.5	18.8		19.3
North	20.6	20.5	19.2	18.3	18.7		19.5
East	18.7	18.7	18.1	17.3	16.8		18.1
Education							
No educ./Pri. incomp.	18.1	18.0	17.9	17.2	17.3		17.6
Pri. comp./Sec. incomp.	19.9	19.8	18.6	18.9	18.8		19.3
Sec. comp./+	24.1	23.8	23.0	21.9	22.9		23.5
All women	20.4	20.3	19.0	18.5	18.4		19.5
HUSBANDS							
Background characteristic	Current age					Husbands age 50-64	Husbands age 25-64
	25-29	30-34	35-39	40-44	45-49		
Residence							
Urban	23.8	24.4	24.8	23.5	23.2	24.5	24.0
Rural	23.0	22.6	22.7	21.8	21.7	23.7	22.6
Region							
West	24.0	24.5	24.7	23.3	22.9	(24.1)	24.0
South	23.2	24.8	23.7	23.3	(23.6)	24.8	24.0
Central	22.9	23.4	23.8	22.5	22.5	(24.4)	23.1
North	(22.5)	24.0	22.6	22.8	(22.3)	(23.6)	22.9
East	24.0	(22.6)	(23.1)	22.1	(23.6)	23.2	23.0
Education							
No educ./Pri. incomp.	*	(22.7)	(23.1)	(22.3)	(22.9)	22.6	22.7
Pri. comp./Sec.	23.1	23.6	23.4	22.4	22.6	24.6	23.2
Sec. comp./+	a	24.9	26.0	25.4	23.9	27.1	a
All husbands	23.5	24.0	24.0	22.9	22.9	24.2	23.6


Note: Parentheses indicate that a figure is based on 25-49 respondents. An asterisk indicates a figure is based on fewer than 25 respondents and has been suppressed.

a Less than 50 percent of respondents in age group x to $x+4$ were first married by age x .

in the median age at first marriage for the husbands is different in certain aspects from that of the women. The median age at marriage among husbands in the Central, North, and East regions is around 23 years, approximately one year lower than median age at marriage among husbands in the West and South regions (Figure 6.2).

As expected, there is positive association between the median age at first marriage and educational level of respondents. The differentials between women who have completed at least secondary school and other women are especially pronounced. The median age at first marriage among women with at least secondary education is 23.5 years, almost 6 years higher than the median age among women with less than primary

education (Table 6.3). For women with a primary or higher education, there is a one-year increase in the age at marriage from older cohorts to younger ones. The increase in the median age at first marriage across cohorts observed for women who have no education is not as great as for the other education groups. There also are differences in the median age at first marriage across education categories for husbands; however, these differences are not as marked as for women. Moreover, the upward trend in the median age at first marriage from older cohorts to younger ones observed among women, is not evident for the husbands.



6.3 Age at First Sexual Intercourse

Age at first marriage is often used as a proxy for the onset of exposure to the risk of pregnancy. However, people can become sexually active before marriage, and the age at which sexual intercourse is initiated marks the beginning of reproductive risk. In the 1998 TDHS only husbands are asked to state the age at which they first had sexual intercourse.

The percentage of husbands who have ever had intercourse by specific ages is given in Table 6.4. Almost one-third of husbands age 25-29 had begun to have intercourse by age 18. The percentages who became sexually active at each specific age are slightly higher for younger cohorts than for older cohorts. Overall, the median age at first intercourse is 19.2, indicating that men initiate sexual relations on average 4 years earlier than they marry (23.6 years).

Table 6.4 Age at first sexual intercourse

Percentage of husbands age 25-64 who had first sexual intercourse by exact ages and selected background characteristics, Turkey 1998

Age	Exact age at first sexual intercourse					Number of husbands	Median
	15	18	20	22	25		
25-29	6.4	29.1	56.5	70.9	88.9	342	19.2
30-34	9.1	36.7	58.7	69.8	86.7	364	18.9
35-39	4.1	36.2	64.1	70.9	89.1	352	18.8
40-44	4.1	29.0	59.8	73.5	90.2	335	19.0
45-49	2.4	27.4	61.0	68.9	88.9	240	19.3
50-64	2.0	21.7	39.0	55.9	80.1	226	20.9
Men age 25-64	5.0	30.8	57.4	69.1	87.7	1,859	19.2

Table 6.5 presents differentials in the median age at first sexual intercourse by background characteristics for the husbands age 25-64. With respect to place of residence, husbands living in urban settlements start sexual relations earlier than their rural counterparts. The regional variation is also noteworthy. The East has the highest median age at first sexual intercourse (20.7) while the West and South have the lowest (18.7). The median age at first intercourse declines as the level of education increases. Husbands with no formal education initiate sexual relations nearly two years later than those with formal education.

Table 6.5 Median age at first intercourse: husbands

Median age at first sexual intercourse among husbands age 25-64 years, by current age and selected background characteristics, Turkey 1998

Background characteristic	Current age						Husbands 25-64
	25-29	30-34	35-39	40-44	45-49	50-64	
Residence							
Urban	18.8	18.7	18.7	18.9	19.0	21.2	19.0
Rural	20.7	19.5	19.2	19.3	19.7	20.6	19.8
Region							
West	18.6	18.0	18.5	18.8	18.6	20.9	18.7
South	19.0	18.3	18.6	18.4	18.9	20.5	18.7
Central	20.1	21.1	19.1	19.3	19.8	21.3	20.0
North	20.1	20.6	19.2	20.1	19.4	20.9	19.9
East	21.0	19.3	21.3	20.6	21.6	20.8	20.7
Education							
No educ./Pri. incomp.	19.1	20.6	22.4	22.1	21.8	20.7	20.8
Pri. comp./Sec. incomp.	19.3	18.5	18.9	18.8	19.1	21.8	19.0
Sec. comp./+	19.1	20.3	18.6	19.1	19.5	18.6	19.0
All men	19.2	18.9	18.8	19.0	19.3	20.9	19.2

6.4 Postpartum Amenorrhoea, Postpartum Abstinence, and Insusceptibility

The period of postpartum amenorrhoea is the interval between childbirth and the return of menstruation. Postpartum protection from conception can be prolonged by two factors: breastfeeding and sexual abstinence. Breastfeeding lengthens the duration of amenorrhoea and postpartum abstinence delays the resumption of sexual relations. Women are considered as insusceptible if they are not exposed to the risk of pregnancy, either because they are amenorrhoeic or are still abstaining from sex following a birth.

The percentage of births whose mothers are postpartum amenorrhoeic, abstaining, and postpartum insusceptible is presented in Table 6.6 by the number of months since the birth. The estimates of the median and mean durations shown in the table are calculated from these current status proportions. The data are grouped by two-month intervals to minimize fluctuations in the estimates.

The results in Table 6.6 indicate the median duration of postpartum amenorrhoea is about 3 months. Eighty-nine percent of women are amenorrhoeic immediately following the delivery, but this value decreases to 53 percent starting from the second month after birth.

Table 6.6 Postpartum amenorrhoea, abstinence and insusceptibility


Percentage of births whose mothers are postpartum amenorrhoeic, abstaining and insusceptible, by number of months since birth, and median and mean durations, Turkey 1998

Months since birth	Amenorrhoeic	Abstaining	Insusceptible	Number of births
< 2	88.9	79.3	95.8	103
2-3	53.1	21.1	61.2	147
4-5	35.7	10.5	40.7	118
6-7	24.6	11.5	34.2	124
8-9	13.6	8.3	20.1	129
10-11	3.2	6.0	9.1	102
12-13	7.9	4.3	11.5	139
14-15	5.4	6.5	9.3	131
16-17	4.3	2.3	6.7	137
18-19	0.5	7.5	8.0	102
20-21	3.2	3.0	6.2	105
22-23	0.0	2.2	2.2	98
24-25	1.6	6.8	8.3	116
26-27	0.0	0.9	0.9	109
28-29	0.0	2.4	2.4	106
30-31	1.5	1.2	2.8	126
32-33	0.0	0.4	0.4	79
34-35	1.9	1.1	3.0	108
Total	14.3	9.7	18.7	2,078
Median	3.3	1.9	4.1	-
Mean	5.2	3.9	6.7	-
Prevalence/ Incidence mean ¹	5.1	3.4	6.6	-

¹ The prevalence/incidence mean is borrowed from epidemiology and is defined as the number of children whose mothers are amenorrhoeic (prevalence) divided by the average number of births per month (incidence).

In Turkey, traditionally there is a period of sexual abstinence after birth that lasts 40 days. The estimates of postpartum abstinence in Table 6.6 are in accordance with this tradition. Of all mothers, 79 percent abstain from sexual relations immediately following a birth. However, starting from the second month after a birth, the contribution of abstinence to the period of insusceptibility is greatly reduced since more women resume sexual relations. At 2-3 months following a birth, the percentage of abstaining mothers decreases to 21 percent and by 6-7 months to 12 percent (Figure 6.3).

Figure 6.3
**Percentage of Births Whose Mothers are Amenorrhoeic,
 Abstaining, or Insusceptible**



TDHS 1998

Table 6.7 shows the median duration of postpartum amenorrhoea, abstinence, and insusceptibility by background characteristics of mothers. In the absence of contraception, variations in the lengths of the periods of postpartum amenorrhoea and abstinence are the most important determinants of the interval between births and, ultimately, of completed fertility. In some populations, differentials across subgroups in the duration of postpartum amenorrhoea and abstinence also may indicate incipient changes in traditional postpartum practices. The average duration of postpartum abstinence does not vary greatly according to the background characteristics of women. However, some variation is observed in the duration of postpartum amenorrhoea by age, region, and level of education. Older women, women living in the South, and women with less than primary education have the longest median duration for postpartum amenorrhoea. It is noteworthy that the shortest duration for postpartum amenorrhoea, 2.8 months, is found in the Central region.

Differentials in the median duration of postpartum insusceptibility reflect the combined effects of amenorrhoea and abstinence. They exhibit a pattern similar to those for amenorrhoea. In general, women over 30, women living in the East and women with no education are insusceptible for relatively longer periods.

Table 6.7 Median duration of postpartum insusceptibility by background characteristics

Median number of months of postpartum amenorrhoea, postpartum abstinence, and postpartum insusceptibility, by selected background characteristics, Turkey 1998

Background characteristic	Postpartum amenorrhoea	Postpartum abstinence	Postpartum insusceptibility	Number of births
Age				
<30	3.0	1.7	3.8	1,460
30+	4.0	2.3	4.8	618
Residence				
Urban	3.2	1.9	4.1	1,290
Rural	3.4	1.8	4.2	788
Region				
West	3.2	2.0	3.3	617
South	3.8	1.9	3.9	288
Central	2.8	1.7	3.7	486
North	3.0	1.8	3.2	159
East	3.6	2.2	6.4	528
Education				
No educ./Pri. incomp.	4.2	2.0	6.1	525
Pri. comp./Sec. incomp.	3.1	1.9	3.6	1,300
Sec. comp./+	3.3	1.8	3.4	254
Total	3.3	1.9	4.1	2,078

Note: Medians are based on current status information.

6.5 Termination of Exposure to Pregnancy

Above age 30, the risk of pregnancy declines with age as increasing proportions of women become menopausal. Table 6.8 presents data on the proportion of women who are menopausal among non-pregnant, non-amenorrhoeic currently married women whose last menstrual period occurred six or more months preceding the survey. The percentage of women in menopause increases gradually with age, rising rapidly after age 45. At age 48-49, 43 percent of women are menopausal.

Table 6.8 Termination of exposure to the risk of pregnancy

Menopause indicator among currently married women age 30-49, by age, Turkey 1998

Age	Menopause ¹	
	Percentage	Number
30-34	0.7	1,202
35-39	2.7	1,081
40-41	3.9	354
42-43	11.2	364
44-45	19.8	350
46-47	29.3	269
48-49	42.5	282
Total	9.2	3,901

¹ Percentage of non-pregnant, non-amenorrhoeic currently married women whose last menstrual period occurred six or more months preceding the survey or who report that they are menopausal.

CHAPTER 7

FERTILITY PREFERENCES


Turgay Ünalan and Elif Kurtuluş

This chapter addresses the following questions which allow an assessment of the need for contraception. Does the respondent want more children? If so, how long would he/she prefer to wait before the next child? If he/she could start afresh, how many children in all would he/she want? To what extent do unwanted or mistimed pregnancies occur? What effect would the prevention of such pregnancies have on the fertility rates?

7.1 Desire for More Children

Currently married women were asked to state their preferences on whether they want to have a(nother) child in the future, and if they so desired, the timing of the next birth. For pregnant women, the question on desire for more children was rephrased to refer to desire for another child after the one that they were expecting. In some cases, the answers of pregnant women with respect to preferred waiting time before the next birth may have included the remaining gestation period of the current pregnancy and, thus, may not be strictly comparable with the answers of non-pregnant women. Also, women who have been sterilised for contraceptive purposes were not asked about their desire for another child. However, for purposes of the fertility preference analysis, these women are classified as wanting no more children.

Two-thirds of currently married women said that they do not want to have a birth in the future or were already sterilised for contraceptive purposes (Figure 7.1). Another 14 percent wanted another child only after waiting two years or more. Thus, overall, four out of every five currently married women can be



regarded as in need of family planning either to avoid or to postpone childbearing. Among the remaining women, most want another birth soon (within two years). The proportion of currently married women who were undecided about having another child was only 3 percent while 5 percent were infecund.

Table 7.1 shows the variation in fertility preferences of both currently married women and husbands with the number of living children.¹ The proportion of currently married women who do not want to have additional births increases sharply with the number of children, from 2 percent for women with no children to 75 percent for women with two living children. In parallel with this increase, the proportion of currently married women who want to have more children decreases sharply. Among women with one child, the majority wants either to delay having another birth (45 percent) or to have no more children (26 percent). Husbands' fertility preferences are very similar to those of currently married women except that more husbands than women want to postpone their first birth (36 percent versus 14 percent, respectively).

Table 7.1 Fertility preference by number of living children						
Percent distribution of currently married women and husbands by desire for more children, according to number of living children, Turkey 1998						
Desire for more children	Number of living children ¹					Total
	0	1	2	3	4+	
CURRENTLY MARRIED WOMEN						
Have another soon ²	69.7	18.0	3.2	1.8	1.9	10.6
Have another later ³	13.6	45.4	9.0	3.5	0.8	13.6
Have another, undecided when	4.7	2.6	1.6	0.2	1.1	1.7
Undecided	1.1	5.3	4.0	1.3	0.6	2.8
Want no more	2.2	25.9	74.5	81.3	80.6	62.1
Sterilised	0.0	0.1	4.2	6.7	7.3	4.3
Declared infecund	8.7	2.5	3.5	5.2	7.7	4.9
Missing	0.0	0.1	0.0	0.0	0.1	0.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Number	453	1,162	1,867	1,128	1,310	5,921
HUSBANDS						
Have another soon ²	47.6	19.5	3.0	3.9	2.4	10.3
Have another later ³	36.4	42.0	11.1	3.2	2.4	15.8
Have another, undecided when	1.4	2.7	1.7	0.4	1.6	1.6
Undecided	0.8	2.6	4.1	1.8	2.4	2.7
Wants no more	7.2	30.3	74.3	81.9	79.8	62.3
Husband/Wife sterilised	0.0	0.4	3.9	5.9	6.9	4.0
Husband/Wife declared infecund	6.5	2.2	1.9	3.0	4.4	3.2
Missing	0.2	0.1	0.0	0.0	0.1	0.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Number	187	372	566	334	512	1,971
¹ Includes current pregnancy ² Wants next birth within 2 years ³ Wants to delay next birth for 2 or more years						

¹ For pregnant women, the current pregnancy is included in the total number of children.

As Table 7.2 shows, the proportion wanting to limit childbearing (including those who are sterilised) increases rapidly with age. Among women age 20-24, one-third desired no more children and, among those age 25-29, 57 percent said they did not want another birth. The proportion wanting no more children peaks at nearly 90 percent among women age 35-39 before falling off as more women classify themselves as infecund.

Only about one-fourth of husbands age under age 30 want to have no more children. However, this proportion increases to 60 percent among husbands age 30-34 and to more than 80 percent among those age 40 and older. The intention to have a child soon or to postpone declines quickly with increasing age to below 10 percent among husbands age 40 and over.

Overall, for both women and husbands, education is negatively associated with the desire to stop childbearing. This is largely because those who are better educated are younger and have fewer children. Among those with two or more births, the percentage wanting no more children declines with increasing education.

Table 7.2 Fertility preference by age

Percent distribution of currently married women and husbands by desire for more children, according to age, Turkey 1998

Desire for more children	Age of woman							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
Have another soon ¹	30.2	19.2	16.0	9.7	4.8	2.3	1.1	10.6
Have another later ²	47.3	41.4	19.7	5.4	0.4	0.2	0.1	13.6
Have another, undecided when	4.1	2.4	2.4	2.2	0.8	0.6	0.0	1.7
Undecided	3.3	4.4	4.5	3.4	1.5	0.7	0.4	2.8
Want no more	15.1	32.1	55.0	71.9	80.7	79.7	69.5	62.1
Sterilised	0.0	0.3	1.8	5.9	7.1	7.0	5.9	4.3
Declared infecund	0.0	0.3	0.4	1.4	4.6	9.5	23.0	4.9
Missing	0.0	0.0	0.2	0.0	0.1	0.1	0.0	0.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Number	262	924	1,196	1,090	1,014	789	645	5,921

Desire for more children	Age of husband							Total
	15-24	25-29	30-34	35-39	40-44	45-49	50+	
Have another soon ¹	20.1	24.4	10.2	7.8	5.3	2.1	4.4	10.3
Have another later ²	53.8	39.2	21.9	7.8	1.8	0.9	0.3	15.8
Have another, undecided when	1.2	3.7	2.1	0.9	1.5	0.8	0.0	1.6
Undecided	2.1	5.1	3.7	3.2	1.0	1.1	0.8	2.7
Want no more	21.6	26.9	57.0	74.2	80.5	81.9	78.8	62.3
Husband/Wife sterilised	1.2	0.2	3.2	5.2	6.8	6.9	3.3	4.0
Husband/Wife declared infecund	0.0	0.4	1.7	0.8	3.1	6.1	12.5	3.2
Missing	0.0	0.0	0.2	0.2	0.0	0.1	0.0	0.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Number	112	342	364	352	335	240	226	1,971

¹ Wants next birth within 2 years

² Wants to delay next birth for 2 or more years

Table 7.3 shows that the desire to stop childbearing does not vary greatly by type of place of residence or region. The largest difference with respect to residential characteristics is for women with two living children, in particular when the regional differences are considered. More women living in urban areas and in the West region desire to stop childbearing compared to those in rural areas and in the East region. As regards the background characteristics very similar trends are observed for husbands. The largest differences in the proportions of husbands who want to stop childbearing are observed for those with two children. More husbands living in the urban areas and in the West desire to stop childbearing compared to those living in rural areas and in the South and Central.

Table 7.3 Desire to limit (stop) childbearing						
Percentage of currently married women and husbands who want no more children, by number of living children and selected background characteristics, Turkey 1998						
Background characteristic	Number of living children ¹					Total
	0	1	2	3	4+	
CURRENTLY MARRIED WOMEN						
Residence						
Urban	2.2	28.2	81.0	88.5	87.7	65.8
Rural	2.1	20.1	73.3	87.2	88.2	67.5
Region						
West	0.7	28.7	84.2	89.9	88.2	65.2
South	2.0	18.5	76.4	82.2	88.7	63.9
Central	5.5	27.5	78.9	93.1	91.0	69.7
North	(1.3)	24.1	74.6	85.2	91.5	69.8
East	2.3	21.8	59.8	80.7	83.9	64.5
Education						
No educ./Pri. incomp.	4.6	31.2	70.2	82.3	86.0	74.5
Prim. comp./Sec. incomp.	1.0	20.0	78.1	90.0	91.4	64.9
Sec. comp./+	3.1	39.5	86.8	(92.1)	*	57.2
Total	2.2	26.0	78.7	88.0	87.9	66.3
HUSBANDS						
Residence						
Urban	10.0	31.4	80.1	88.8	86.5	65.7
Rural	0.0	28.7	72.8	85.7	86.9	67.8
Region						
West	10.8	31.9	84.3	90.1	86.5	64.4
South	(5.3)	(27.4)	68.4	80.2	90.0	65.7
Central	(6.8)	32.8	78.9	95.0	86.8	68.9
North	*	*	(75.0)	(85.2)	90.0	69.1
East	(0.0)	*	(58.9)	(78.9)	83.9	66.5
Education						
No educ./Pri. incomp.	*	*	*	(93.7)	79.7	75.1
Pri. comp./Sec. incomp.	6.3	27.3	78.6	88.0	89.0	68.2
Sec. comp./+	5.9	35.0	78.5	84.9	89.9	58.2
Total	7.2	30.7	78.3	87.8	86.7	66.3

Note: Women who have been sterilised are considered to want no more children.
¹ Includes current pregnancy
 () Figure in parentheses is based on 25-49 cases
 * Less than 25 cases

7.2 Need for Family Planning Services

Information on fertility preferences alone is not sufficient to assess the need for family planning services. Many women who do not want to have another child or who want to space the next birth are already using contraception or are not exposed to the risk of pregnancy because they are menopausal or infecund. In general, women who are currently married and who declare either that they do not want to have any more children (to limit their childbearing) or that they want to wait two or more years before having another child (to space their births), but are not currently using contraception, have an unmet need for family planning. The calculation of unmet need is further refined by excluding women who are currently amenorrhoeic and, therefore, not in need of family planning at present. Women with unmet need and those currently using contraception constitute the total demand for family planning.

The data in Table 7.4 indicate that 10 percent of currently married women in Turkey are in need of a family planning method, either for spacing (4 percent) or for limiting (6 percent). An additional 64 percent of currently married women are using contraception, (14 percent to delay the next birth and 50

Table 7.4 Need for family planning services

Percentage of currently married women with unmet need for family planning, met need for family planning, and the total demand for family planning services, by selected background characteristics, Turkey 1998

Background characteristic	Unmet need for family planning ¹			Met need for family planning ² (currently using)			Total demand for family planning ³			Percentage of demand satisfied	Number of women
	For spacing	For limiting	Total	For spacing	For limiting	Total	For spacing	For limiting	Total		
Age											
15-19	16.6	3.4	20.0	30.3	3.2	33.6	51.1	6.6	57.7	65.4	262
20-24	10.9	5.1	16.0	33.9	19.0	52.9	47.3	25.2	72.5	77.9	924
25-29	4.6	5.9	10.5	24.4	42.6	67.0	30.4	49.4	79.8	86.8	1,196
30-34	2.0	7.5	9.6	11.3	63.0	74.3	13.4	71.8	85.3	88.8	1,090
35-39	0.3	6.2	6.6	2.6	73.7	76.3	3.0	80.5	83.5	92.2	1,014
40-44	0.1	7.1	7.2	1.2	68.8	70.0	1.3	76.1	77.4	90.7	789
45-49	0.3	7.0	7.3	0.0	41.4	41.4	0.3	48.5	48.8	85.1	645
Residence											
Urban	3.3	5.2	8.5	15.6	51.1	66.7	19.8	56.9	76.7	88.9	3,978
Rural	5.0	8.5	13.5	11.5	46.5	58.1	17.4	56.1	73.5	81.6	1,943
Region											
West	2.3	3.4	5.7	17.2	53.4	70.5	20.4	57.1	77.4	92.7	2,261
South	4.1	5.9	10.0	13.2	47.1	60.3	18.3	53.9	72.2	86.1	851
Central	3.2	5.9	9.2	14.2	54.1	68.3	18.3	60.7	79.0	88.4	1,426
North	3.0	5.4	8.4	12.1	54.9	67.0	16.0	61.3	77.3	89.1	474
East	8.9	14.9	23.8	9.1	32.9	42.0	18.8	49.3	68.1	65.0	909
Education											
No educ./Pri. incomp.	3.8	11.9	15.7	4.0	46.4	50.4	8.1	58.9	67.0	76.7	1,546
Pri. comp./Sec. incomp.	3.9	4.8	8.7	16.2	51.0	67.1	21.1	56.6	77.7	88.8	3,570
Sec. comp./+	3.7	2.3	6.1	25.5	49.8	75.3	30.5	52.5	82.9	92.7	804
Total	3.8	6.3	10.1	14.3	49.6	63.9	19.0	56.6	75.6	86.6	5,921

¹ Unmet need for spacing includes pregnant women whose pregnancy was mistimed, amenorrhoeic women whose last birth was mistimed, and women who are neither pregnant nor amenorrhoeic and who are not using any method of family planning but say they want to wait two or more years for their next birth. Also included in unmet need for spacing are women who are unsure whether they want another child or who want another child but are unsure when to have the birth. Unmet need for limiting refers to pregnant women whose pregnancy was unwanted, amenorrhoeic women whose last child was unwanted, and women who are neither pregnant nor amenorrhoeic and who are not using any method of family planning but want no more children. Excluded from the unmet need category are menopausal or infecund women.

² Using for spacing is defined as women who are using some method of family planning and say they want to have another child or are undecided whether to have another. Using for limiting is defined as women who are using and who want no more children. Note that the specific methods used are not taken into account here.

³ Total demand for family planning includes pregnant or amenorrhoeic women who became pregnant while using a method (method failure).

percent to stop childbearing). Another 2 percent of women had become pregnant while using a method and, thus, are considered to be in a need of a better method (not shown separately in table). Thus, the total demand for family planning among currently married women in Turkey is 76 percent, 19 percent for spacing purposes and 57 percent for limiting purposes. As was in the TDHS-93, 87 percent of the total demand has been satisfied.

The overall unmet need for family planning declines with increasing age. Unmet need for spacing purposes is higher among the younger women, while unmet need for limiting childbearing is higher among the older women. The level of unmet need is also more pronounced among rural women, among women in the East, and among uneducated women. The percentage of the total demand that is satisfied is higher for urban women, women living in the West, and for educated women.

7.3 Ideal and Actual Number of Children

The tables examined so far in this chapter take into account the number of sons and daughters that women and husbands already have. In order to ascertain what the respondents consider to be the ideal number of children (i.e., their desired family size), TDHS respondents were asked "If you could go back to the time you did not have any children and could choose exactly the number of children to have in your whole life, how many would that be?" Table 7.5 presents the ideal number of children according to the number of living children. A strong desire for a two-child family is evident. Fifty-seven percent of women and 52 percent of husbands consider two children to be the ideal family size. Three-child families are preferred by 19 percent of women and 22 percent of husbands. The mean ideal number of children is 2.5 among currently married women and 2.7 among husbands. (The comparable figure from the TDHS-93 was 2.4 for currently married women.)

The mean ideal number of children increases with the number of living children for both women and husbands (from 2.2 to 3.0 among currently married women and from 2.4 to 3.6 among husbands). The reason is twofold. First, to the extent that women or husbands implement their preferences, those who want larger families tend to achieve larger families. Second, women or husbands may adjust upwards their ideal family size, as the actual number of children increases (i.e., as a result of rationalisation).

Table 7.5. Ideal and actual number of children

Percent distribution of currently married women and husbands by ideal number of children and mean ideal number of children for all women and currently married women and men, according to number of living children, Turkey 1998

Ideal number of children	Number of living children ¹					Total
	0	1	2	3	4+	
WOMEN						
0	0.0	0.0	0.4	0.1	0.8	0.3
1	11.4	10.8	5.2	7.1	2.6	6.6
2	60.7	66.5	67.9	46.2	41.0	57.0
3	16.9	16.8	16.2	29.2	16.8	19.0
4	5.6	3.2	6.6	12.2	22.8	10.5
5	0.5	0.7	0.9	1.0	3.7	1.5
6+	0.5	0.3	0.5	0.8	4.4	1.4
Non-numeric response	4.3	1.9	2.3	3.3	7.8	3.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Number of women	453	1,162	1,867	1,128	1,310	5,921
All women						
Mean ideal number	2.1	2.2	2.3	2.6	3.0	2.4
Number of women	2,716	1,195	1,901	1,132	1,271	8,215
Currently married women						
Mean ideal number	2.2	2.2	2.3	2.6	3.0	2.5
Number of women	434	1,140	1,824	1,091	1,208	5,696
HUSBANDS						
0	0.0	1.2	0.8	0.4	1.0	0.8
1	7.5	12.3	5.3	6.2	2.6	6.3
2	54.3	60.8	66.6	40.7	34.3	51.6
3	24.1	19.0	15.7	37.2	19.7	21.8
4	5.9	3.3	6.6	8.1	18.8	9.3
5	1.9	0.4	1.6	0.6	3.2	1.6
6+	0.4	1.1	1.3	2.9	9.9	3.7
Non-numeric response	5.9	2.1	2.0	3.9	10.6	4.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Number of husbands	187	372	566	334	512	1,971
Mean ideal number	2.4	2.2	2.4	2.7	3.6	2.7
Number of husbands	176	365	555	321	458	1,874

Note: The means exclude women/husbands who gave non-numeric responses.
¹ Includes current pregnancy.

Table 7.6 presents the mean ideal number of children for currently married women and husbands by age and selected background characteristics. Mean ideal family size increases with age; from 2.3 children among women age 15-19 to 2.7 children among women age 45-49 and from 2.6 children among husbands age 15-19 to 3.3 among husbands age 50 or over. The mean ideal number of children is higher among women and husbands living in rural areas, in the East, and among less educated women and husbands. Husbands tend to have a somewhat higher ideal family size than women regardless of the background characteristic examined.

Table 7.6 Mean ideal number of children by background characteristics								
Mean ideal number of children for currently married women and husbands, by age and selected background characteristics, Turkey 1998								
Background characteristic	Age of woman							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
Residence								
Urban	2.3	2.3	2.3	2.4	2.5	2.6	2.5	2.4
Rural	2.3	2.6	2.5	2.5	2.6	2.8	2.8	2.6
Region								
West	2.1	2.1	2.2	2.3	2.4	2.3	2.3	2.3
South	2.6	2.5	2.4	2.6	2.6	2.8	3.0	2.6
Central	2.1	2.3	2.2	2.4	2.4	2.6	2.7	2.4
North	2.2	2.2	2.2	2.4	2.3	2.5	2.8	2.4
East	2.5	2.8	3.0	3.1	3.1	3.8	3.4	3.1
Education								
No educ./Pri. incomp.	2.6	2.8	3.1	2.9	2.8	3.0	3.0	2.9
Pri. comp./Sec. incomp.	2.2	2.3	2.3	2.4	2.5	2.4	2.4	2.4
Sec. comp./+	2.4	2.1	2.0	2.2	2.2	2.2	1.9	2.1
Total women	2.3	2.4	2.4	2.5	2.5	2.6	2.7	2.5
	Age of husband							Total
	15-24	25-29	30-34	35-39	40-44	45-49	50+	
Residence								
Urban	2.6	2.4	2.6	2.6	2.9	2.6	3.1	2.7
Rural	(2.6)	2.6	2.7	2.6	2.7	2.8	3.6	2.8
Region								
West	(2.4)	2.3	2.4	2.3	2.3	2.2	2.6	2.3
South	*	2.8	3.1	3.4	2.6	2.5	3.8	3.0
Central	(2.6)	2.4	2.5	2.3	2.5	2.7	2.3	2.5
North	*	2.8	2.5	2.9	2.9	2.6	2.4	2.7
East	(3.2)	3.0	3.6	3.4	5.2	4.4	4.6	4.0
Education								
No educ./Pri. incomp.	*	*	*	(3.3)	(5.2)	(4.4)	4.0	4.1
Pri. comp./Sec. incomp.	2.7	2.6	2.6	2.7	2.7	2.4	3.0	2.6
Sec. comp./+	2.4	2.2	2.6	2.4	2.4	2.4	2.6	2.4
Total men	2.6	2.5	2.6	2.6	2.8	2.6	3.3	2.7
Note: Parentheses indicate that a figure is based on 25-49 respondents. An asterisk indicates a figure is based on fewer than 25 respondents and has been suppressed.								

7.4 Fertility Planning

Women were asked a series of questions for each child born in the preceding five years and any current pregnancy to determine whether the particular pregnancy was planned (wanted then), unplanned (wanted later), or not wanted at all (wanted no more). These questions form a potentially powerful indicator of the degree to which couples successfully control childbearing. However, the questions are extremely demanding. The respondent is required to recall accurately her wishes at one or more points in the last five years and to report them honestly. The danger of rationalisation is present; an unwanted conception may well become a cherished child. Therefore, the values presented here are likely to be underestimates of unwanted fertility.

Table 7.7 is a birth-based rather than a woman-based table and provides perhaps the single most useful indicator of the degree of successful reproductive control exercised by couples in the recent past. Overall, 69 percent of births in the past five years were wanted at the time they were conceived, whereas 11 percent were wanted later and 19 percent were not wanted at all. The proportion of births that are reported as not wanted increases with birth order and mother's age at birth. Nearly half of the women who have four or more children or are 35 years or older reported that their last child was not wanted. The proportion of mistimed births is highest among women with two children and those under age 19.

Birth order and mother's age at birth	Planning status of birth				Total	Number of births
	Wanted then	Wanted later	Not wanted	Missing		
Birth order						
1	87.0	10.0	2.1	0.9	100.0	1,343
2	70.8	19.1	9.3	0.9	100.0	1,093
3	60.5	10.2	29.0	0.3	100.0	591
4+	45.7	4.1	49.6	0.7	100.0	863
Age at birth						
<19	74.8	18.8	6.1	0.3	100.0	554
20-24	74.7	14.3	10.3	0.8	100.0	1,421
25-29	68.9	9.6	20.6	0.9	100.0	1,042
30-34	61.3	4.9	32.8	1.1	100.0	590
35-39	49.9	0.5	49.3	0.3	100.0	220
40-44	47.7	1.1	51.2	0.0	100.0	61
45-49	*	*	*	*	100.0	3
Total	69.2	11.2	18.8	0.7	100.0	3,891

Note: Birth order includes current pregnancy.
* Less than 25 cases

Another way of measuring the extent of unwanted fertility is to estimate what the fertility rate would be if all unwanted births were avoided. This rate, known as the *wanted fertility rate*, is calculated in a similar manner as the total fertility rates presented in Chapter 3. The Lightbourne method of calculating a "wanted" birth is used for Table 7.8. Under the Lightbourne method, a birth is considered wanted if the number of living children at the time of conception was less than the current ideal number of children, as reported by the respondent. Births classified as unwanted according to above definition are omitted from the numerator and the remainder is cumulated to form a wanted total fertility rate which is analogous to the conventional total fertility rate. This rate represents the level of fertility that

theoretically would result if all unwanted births were prevented. Comparison of actual rates with wanted rates indicates the potential demographic impact of the elimination of unwanted births.

Table 7.8 presents the total wanted fertility rate and the total fertility rate by selected background characteristics. The total wanted fertility rate is 1.9 births per women compared to the total fertility rate of 2.6 (1.8 and 2.7 in 1993 TDHS, respectively). The differences between the wanted and the actual fertility rates are greatest for women living in rural areas, in the East, and among women who have no or less than a primary education.

Table 7.8 Wanted fertility rates		
Total wanted fertility rates and total fertility rates for the three years preceding the survey, by selected background characteristics, Turkey 1998		
Background characteristic	Total wanted fertility rate	Total fertility rate
Residence		
Urban	1.9	2.4
Rural	2.1	3.1
Region		
West	1.7	2.0
South	1.9	2.6
Central	2.0	2.6
North	1.9	2.7
East	2.7	4.2
Education		
No educ./Pri. incomp.	2.4	3.9
Pri. comp./Sec. incomp.	2.0	2.6
Sec. comp./+	1.5	1.6
Total	1.9	2.6
Note: The total fertility rates are the same as those presented in Table 3.2.		

CHAPTER 8

INFANT AND CHILD MORTALITY

Attila Hancıoğlu

The level of early childhood mortality is an important indicator of the level of development of a society, particularly of the prevailing health conditions. Thus, infant mortality rates and under-five mortality rates are used to assess the level of development, commonly as part of composite indexes. The use of these rates is not limited to such assessments; they are also important in making informed decisions about health and population policies and programs. They can be used to identify population groups facing higher mortality risks, who should be priority targets in efforts to increase child survival. They also are used in making population projections.

This section focuses on the findings of the TDHS-98 with respect to mortality during infancy and early childhood. In the absence of alternative estimates of such rates from other sources, (e.g. a vital registration system), sample surveys in Turkey serve as the main data source for obtaining direct estimates of infant and child mortality. Hence, the results of the TDHS-98 are critical for the reassessment of policies and strategies for the improvement of survival chances of children in Turkey.

8.1 Definitions of Infant and Child Mortality

All ever-married female respondents in the TDHS-98 were asked to provide a complete birth history, including the sex, birth date, survival status, and current age or age at death for each of their live births. The data were used to calculate five measures of infant and child mortality, namely:

- Neonatal mortality, the probability of dying in the first month of life
- Post-neonatal mortality, the probability of dying after the first month of life but before the first birthday
- Infant mortality (${}_1q_0$), the probability of dying in the first year of life
- Child mortality (${}_4q_1$), the probability of dying between the first and fifth birthday
- Under-five mortality (${}_5q_0$), the probability of dying before the fifth birthday.

8.2 Assessment of Data Quality

Like all other demographic or health indicators from the TDHS-98, infant and child mortality rates are subject to both sampling and non-sampling errors. While the measurement of sampling errors is relatively easy, non-sampling errors are difficult to detect and correct for. Non-sampling errors cover a wide range of errors, involving under-reporting of births and deaths, errors by the interviewers in recording responses, and so forth. International research has shown that infant and child mortality estimates from sample surveys are particularly susceptible to such errors. The first step in the consideration of the TDHS mortality data is to look for evidence of non-sampling errors in order to assess whether information collected in the TDHS-98 on infant and child mortality are of acceptable quality.

One of the most powerful interviewing tools for collecting information on births and deaths is the birth history. However, birth histories are subject to respondent recall errors, and these errors may result in biased rates and trends over time. Therefore, a preliminary assessment of the quality of birth history data is made in this section with respect to completeness and accuracy of date reporting, heaping of age at death, and sex-selective omission of births.

A commonly encountered problem in birth history data is unreported birth dates and ages at death. The TDHS-98 interviewers were required to obtain full information on birth dates (i.e., month and year of birth) for births in the 5 years immediately preceding the survey. Table D.4 shows that complete information on birth dates were indeed collected for almost all births in this period. There is somewhat greater deterioration in the completeness of birth date information the further back one goes from the survey date. Overall, the percentage of live births in the 15 years preceding the survey for which information on month of birth was missing is 9 percent; this figure is relatively high, especially when compared to the results of the 1993 TDHS (2 percent). Both month and year of birth were missing for less than one percent of all live births in the 15-year period before the TDHS. Unreported ages at death also were not common in the TDHS data; less than one percent of deaths recorded in the birth histories lacked an age at death. Thus, with respect to the completeness of the information collected on dates of birth and ages at death, the TDHS appears to be of good quality.

A further assessment of the data suggests there is some evidence of heaping. In particular, Table D.4 shows that there is a deficit of births in the TDHS in the calendar year 1993 and an excess of births in calendar year 1992. This pattern is one found in the 1993 TDHS, as well as DHS surveys in other countries. It is thought to result, at least partly, from the transference of births by interviewers out of the period for which health and calendar data were collected (January 1993 through the date of the survey) in order to reduce their workload.

A problem common to most retrospective surveys is heaping of age at death on "convenient" digits, for example, 6, 12, and 18 months. This phenomenon may lead to the calculation of biased rates, especially if, as a net result, deaths are shifted from one age segment to another. In this regard, one critical shift would be to record infant deaths as child deaths, by respondents heaping the age at death on 12 months, or by interviewers recording ages of death as "1 year". Heaping of age at death at 12 months in the TDHS-98 was minimal, especially in the last 5-year period (see Table D.6). A simulation model was applied to the data to see if the heaping of age at death on 12 months that was evident would bias estimates of infant mortality. The results indicated that any bias in the infant mortality rate from heaping would be less than 5 percent. The rates presented here are therefore unadjusted; that is, all deaths reported at 12 months or "1 year" are assigned to the post-infant age period. Similar results were obtained in the 1993 TDHS; however, the amount of heaping on 12 months (indicating a possible shift of age at death from infancy to childhood) was slightly lower in the 1993 survey.

Another check that can be performed to assess the reliability of birth history data is to calculate sex ratios at birth for all live births. These ratios are expected to fluctuate around 105 male births per 100 female births. Table D.4 shows that the overall sex ratio for all births in the birth history is 105.6, which is in line with expectations. However, the sex ratio of live births during the 5-year period before the TDHS-98 is somewhat higher than expected (108.4), and those for the prior two five-year periods also are in excess of 105. While higher than expected, nevertheless, the ratios for these periods are within acceptable limits for sample surveys and are not likely to bias mortality rates considerably.

8.3 Levels and Trends in Infant and Child Mortality

The first panel of Table 8.1 presents infant and child mortality rates for periods 0-4, 5-9, and 10-14 years preceding the 1998 TDHS. These periods refer approximately to calendar periods of 1993-1998, 1988-1993, and 1983-1988, respectively. The estimated infant mortality rate for the most recent period (0-4 years preceding the survey) is 43 per 1,000 live births. Sixty percent of infant deaths occurred during the neonatal period (i.e.: during the first four weeks of life). Child mortality is found to be approximately 10 per 1,000. The results also show that the probability of dying before the fifth birthday is around 52 per 1,000.

Table 8.1 Infant and child mortality

Infant and child mortality rates by five-year periods preceding the 1998 TDHS and 1993 TDHS


Years preceding survey	Approximate reference period	Neonatal mortality (NN)	Postneonatal mortality (PNN)	Infant mortality (IQ0)	Child mortality (4Q1)	Under-five mortality (5Q0)
1998 TDHS						
0-4	1993-1998	25.8	16.9	42.7	9.8	52.1
5-9	1988-1993	30.1	23.9	54.0	14.0	67.2
10-14	1983-1988	32.4	37.2	69.7	17.0	85.5
1993 TDHS						
0-4	1988-1993	29.2	23.4	52.6	8.8	60.9
5-9	1983-1988	44.6	36.9	81.5	16.8	96.9

The TDHS-98 confirms a major finding of the 1993 TDHS that most of the deaths before the first birthday in Turkey now occur before the first month. In other words, accompanying the general improvements of mortality, a pattern has evolved in which neonatal mortality rates have begun to exceed post-neonatal mortality rates. An examination of the variation in mortality rates by age segments also reveals that a large proportion of under-five deaths occurs before the first birthday (82 percent). This finding is also consistent with previous information on the pattern of under-five mortality in Turkey, including the results of the 1993 TDHS. It is also characteristic of other countries in which child mortality levels have declined sharply.

The figures in Table 8.1 confirm the relatively fast pace of decline in infant and child mortality rates in Turkey. For the two most recent periods, major declines in post-neonatal mortality (29 percent) and child mortality (30 percent) have taken place. The rate of decline for the neonatal mortality rate has been somewhat slower (14 percent). The infant mortality rate has declined by 21 percent.

Table 8.1 also presents comparable mortality estimates from the 1993 TDHS. Such comparisons are useful for further assessing the quality of data, as well as confirming the observed trends. Since the two surveys are exactly five years apart, rates can be calculated for almost exactly the same periods from the two surveys, as indicated in the table. Comparing the estimates of the two surveys for the 1988-93 period, the consistency is impressive; neonatal, post-neonatal and infant mortality rates differ by only less than 1.5 per thousand, while child mortality and under-five mortality estimates differ by 5-6 per thousand. Such differences are negligible in the context of comparisons based on sample survey data, and are statistically not significant. Comparing the estimates from the two surveys for the 1983-88 period, however, there are more evident differences, not only in terms of the overall levels, but also in the indicated underlying age patterns. For this period, the 1993 TDHS found higher neonatal mortality than post-neonatal mortality, while, in the TDHS-98, the opposite pattern was observed. The results of an independent data source, the 1988 TPHS, are closer to those of the 1993 TDHS (shown in Figure 8.1). Given the truncated nature of the results of the TDHS-98 for this period, these results are not surprising and they suggest the need for caution in using the mortality data from TDHS 98 for the period 10-14 years preceding the survey.

Figure 8.1
Trends in Infant and Child Mortality
 Estimates for 5-Year Periods Preceding 1988 TPHS, 1993 TDHS, and 1998 TDHS



TDHS 1998

8.4 Differentials in Infant and Child Mortality

Table 8.2 presents the mortality rates by urban-rural residence, region, level of mother's education, and use of medical maternity care (antenatal care and delivery care) for the five-year period preceding the survey. The figures should be interpreted with caution, since they are based on a small number of observations and are, therefore, statistically unstable. The infant mortality rate in the rural areas is about 56 percent higher than in urban areas (55 and 35 per 1,000, respectively). While neonatal rates for urban and rural areas are quite close, rural post-neonatal and child mortality rates are 2.2 and 1.8 times higher than urban rates, respectively. In both rural and urban areas, neonatal mortality rates are higher than post-neonatal mortality rates.

Infant and under-five mortality rates are lower than the national average in the West and South regions, close to the national average in the Central and North regions, and more than 40 percent higher than the national average in the less-developed East region. Post-neonatal mortality is lowest in the West and highest in the East. In all regions, neonatal mortality is either higher than or very close to post-neonatal mortality.

The relationship between survival chances of children and the level of education of their mothers is well known. The 1998 TDHS results confirm this relationship. For all measures of mortality, probabilities of dying are lower for children of mothers with higher educational levels. For children whose mothers have less than primary education completed, the under-5 mortality rate is 73 per thousand, 67 percent higher than children whose mothers are at least primary school graduates. Differences of similar magnitudes are also observed in other measures. The mortality advantage in favour of children of more educated mothers especially after the first month of life, once again exhibits the importance of the impact education has on child care.

Table 8.2 Neonatal, post-neonatal, infant, child, and under-five mortality by socioeconomic characteristics

Neonatal, post-neonatal, infant, child, and under-five mortality for the five-year period preceding the survey, by socioeconomic characteristics, Turkey 1998

Socioeconomic characteristic	Neonatal mortality (NN)	Post-neonatal mortality (PNN)	Infant mortality (IQ0)	Child mortality (4Q1)	Under-five mortality (5Q0)
Residence					
Urban	23.5	11.7	35.2	7.5	42.4
Rural	29.6	25.4	55.0	13.7	68.0
Region					
West	25.3	7.5	32.8	5.6	38.3
South	16.8	15.9	32.7	10.7	43.0
Central	26.8	14.5	41.3	8.7	49.6
North	19.7	22.4	42.0	8.9	50.5
East	32.6	28.9	61.5	15.4	75.9
Education					
No educ./Pri. incomp.	34.9	25.5	60.5	13.7	73.4
Pri. comp./+	22.4	13.6	36.1	8.1	43.9
Medical maternity care					
Received no ANC or DS	37.3	57.9	95.2	-	-
Received either ANC or DS	26.2	23.9	50.2	-	-
Received both ANC and DS	23.2	5.4	28.6	-	-
Total	25.8	16.9	42.7	9.8	52.1

ANC = Antenatal care
DS = Delivery services

Medical maternity care is also an important factor in the reduction of mortality rates. Infant and child mortality appears to be especially low if the mother has both received antenatal care and delivery care from trained health professionals. In cases when neither care has been provided, infant mortality is as high as 95 per thousand. When either antenatal care or delivery care is obtained, mortality rates decline considerably. Obtaining medical care appears to be especially significant in the improvement of survival chances after the first month of life. Lack of medical care does not seem to significantly elevate mortality risks during the first month of life; however, selection factors are clearly operating in this case (women with problematic pregnancies receiving medical care) and the differentials should be interpreted with caution.

Shown in Table 8.3 are differentials in infant and child mortality by various biodemographic characteristics. In order to base the calculations on sufficient numbers of deaths and exposure, the rates are calculated for the 10-year period before the survey.

The influence of various biodemographic characteristics on survival chances of infants is well documented, and the TDHS-98 results confirm most of the expected relationships. Male mortality rates are higher than those for females during the neonatal period, but higher for females after the neonatal period through age 5. This finding is consistent with the findings of the 1993 TDHS, as well as previous studies on the sex differentials of infant and child mortality in Turkey (Ministry of Health et al, 1994). Table 8.3 also shows that children of teenage mothers, high-birth-order children, and children born following a short birth interval are at greater risk of dying than those in other subgroups. The differentials are especially pronounced in the case of short birth intervals.

Table 8.3 Neonatal, post-neonatal, infant, child, and under-five mortality by biodemographic characteristics

Neonatal, post-neonatal, infant, child, and under-five mortality for the ten-year period preceding the survey, by selected biodemographic characteristics, Turkey 1998

Biodemographic characteristic	Neonatal mortality (NN)	Post-neonatal mortality (PNN)	Infant mortality (IQ0)	Child mortality (4Q1)	Under-five mortality (5Q0)
Sex of child					
Male	32.0	19.1	51.0	10.4	60.9
Female	23.6	21.9	45.5	13.4	58.3
Age of mother at birth					
Less than 20	28.3	25.3	53.5	14.6	67.4
20-29	28.7	18.3	47.0	11.1	57.6
30-39	21.9	22.8	44.6	12.0	56.1
40-49	85.8	25.5	111.4	9.7	120.0
Birth order					
1	27.9	11.1	39.1	8.2	46.9
2-3	26.9	18.9	45.8	8.7	54.1
4-6	35.1	34.6	69.7	18.7	87.0
7+	18.2	36.0	54.3	26.4	79.2
Previous birth interval					
< 2 years	43.8	47.5	91.3	24.1	113.2
2-3 years	14.8	19.2	34.1	12.7	46.3
4 years or more	26.0	7.2	33.1	2.9	35.9
Size at birth¹					
Small or very small	37.4	24.8	62.2	-	-
Average or larger	17.4	12.2	29.6	-	-

¹ Refers to births in the five years before the survey

Children's weight at birth is closely associated with their chances of survival, especially during the neonatal period. Children reported by mothers as "small or very small" at birth were at more than twice the risk of dying compared with children whose birth weight was reported as being "average or larger than average."

8.5 High-risk Fertility Behaviour

As the findings in the previous section indicate, a strong relationship exists between a mother's pattern of fertility behavior and her children's survival chances. Infants and young children born to very young mothers or to older mothers, born after a short birth interval, or born after their mothers have already had many children face higher mortality risks. In the following analysis, mothers are classified as "too young" if they were less than 18 years old at the time of the birth, and "too old" if they were 34 or older at the time of the birth. A "short birth interval" is defined as less than 24 months, and a "high birth order" as one occurring after three or more previous births. Additionally, deaths may be more likely among first order births than higher order births, but such births are not avoidable. For this reason, first births are included in Table 8.4 to show elevated risks associated with them, but excluded from the high-risk categories.

Column 1 in Table 8.4 shows the percentage of children born in the five years preceding the survey who were in specific risk categories. Risk ratios are also presented for each of the risk categories (see column 2, Table 8.4). Here, a risk ratio here is defined as the ratio of the proportion dead among children in a risk category, to the proportion dead among children *not in any high-risk category*. The latter group includes children whose mothers were age 18-34 at delivery, who were born after an interval of 24 or more months after the previous birth, and who are parity of 2 or 3.

Table 8.4 High-risk fertility behaviour

Percent distribution of children born in the five years preceding the survey by category of elevated risk of dying, and the percent distribution of currently married women at risk of conceiving a child with an elevated risk of dying, by category of increased risk, Turkey 1998

Risk category	Births in 5 years preceding the survey		Percentage of currently married women ^a
	Percentage of births	Risk ratio	
Not in any high-risk category	29.9	1.00	27.1 ^b
Unavoidable risk category (First births)	29.9	1.14	8.2
Single high-risk category			
Mother's age <18	4.4	(1.77)	0.6
Mother's age >34	1.9	(0.28)	13.6
Birth interval <24 months	10.9	2.15	9.3
Birth order >3	12.4	1.35	9.4
Subtotal	29.6	1.64	33.0
Multiple high-risk category^c			
Age <18 & birth interval <24 months	0.5	1.93	0.2
Age >34 & birth interval <24 months	0.2	4.42	0.3
Age >34 & birth order >3	4.5	1.31	26.1
Age >34 & birth interval <24 months & birth order >3	0.7	4.51	1.2
Birth interval <24 months & birth order >3	4.7	4.21	3.9
Subtotal	10.6	2.91	31.8
In any high-risk category	40.2	1.97	64.7
Total	100.0	-	100.0
Number of births	3,459	-	5,921

Note: Risk ratio is the ratio of the proportion dead of births in a specific high-risk category to the proportion dead of births *not in any high-risk category*.

^a Women were assigned to risk categories according to the status they would have at the birth of a child, if the child were conceived at the time of the survey: age less than 17 years and 3 months, age older than 34 years and 2 months, latest birth less than 15 months ago, and latest birth of order 3 or higher.

() Figures in parentheses are risk ratios based on less than 22 observations.

^b Includes sterilised women

^c Includes the combined categories *Age <18 and birth order >3*.

Forty percent of children born in the five years preceding the survey were at elevated risk of dying at the time of their birth. First births are not included in this figure. Among all children, 30 percent had an increased risk due to a single risk category, and 11 percent had an increased risk due to multiple factors. It is evident from the table that high birth order and short birth intervals are major factors contributing to elevated risks of mortality. More than a fifth of children born in the last five years were of high birth orders, while 17 percent were born after a short interval.

Children whose mothers were in a single high-risk category faced mortality risks 64 percent more than those whose mothers were in none of the risk categories. For those in a multiple high-risk category, relative mortality risks were up to 2.9 times the risks faced by children not in any risk category (Column 2, Table 8.4). The table shows that children born after a short birth interval faced more than twice the risk of dying compared to those in the no-risk category.

The final column of Table 8.4 includes the distribution of currently married women according to category of increased risk they would have been in if they had conceived at the time of the survey. A woman's current age, time elapsed since last birth, and parity are used to determine into which category her next birth would have fallen if she had conceived at the time of the survey. For example, if a woman age 37 who had five children and had had her last birth three years before the survey were to have become pregnant, she would have fallen into the multiple risk category of being too old (35 or older) and at too high a parity (4 or more children). Women who have been sterilised are categorised as not being in a high-risk category.

Sixty-five percent of women who were married at the time of the TDHS-98 were found to be at risk of conceiving a child with an increased risk of dying. Only 27 percent of women fell into none of the risk categories. Thirty-two percent of women fell into a multiple risk category. As indicated in the second column of the table, if a woman in this category were to conceive, the survival chances of a child would be considerably lower than those of births to women not in the risk categories. The largest group of women fell into the multiple risk category where the child to be born would have had, at the time of birth, a mother who was older than 34 and who already had at least three births.

CHAPTER 9

MATERNAL AND CHILD HEALTH

Banu Akadlı Ergöçmen

This chapter presents findings concerning maternal and child health in Turkey. Information is presented on maternal care during pregnancy and delivery, vaccinations of children, and diarrhoea and its treatment. Data were obtained for all live births that occurred in the five years preceding the survey.

9.1 Antenatal Care and Delivery Assistance

Aspects of antenatal care (ANC) that are examined include the type of provider, the number of visits made, the stage of pregnancy at the time of the first visit, and the number of the tetanus toxoid (TT) doses received. Similarly, the delivery services are described according to the person assisting and the type and place of delivery.

Source of Antenatal Care

Table 9.1 shows the percent distribution of live births in the five years preceding the survey by source of ANC received during pregnancy, according to the maternal background characteristics and birth order. The interviewers were instructed to record all responses if more than one source of ANC was mentioned for the same pregnancy. However, for this tabulation, only the provider with the highest qualifications is considered if there were more than one response.

As seen in Table 9.1, 68 percent of the mothers received at least one ANC visit from trained health personnel. The share of the doctor is more than 60 percent and it worth noting that proportion of mothers seeing a doctor for ANC has increased by 13 percent since the previous survey conducted in 1993. On the other hand, nearly one-third of the mothers did not receive any ANC.

There are marked differences in ANC by background characteristics. Younger mothers are more likely to seek ANC from trained health personnel than women over age 35 (Figure 9.1). In the case of almost half of the births to women age 35 and over, the mother did not receive any ANC. The differences in the proportions of live births with ANC according to birth order are also striking. Receiving ANC declines as the birth order increases. Children of birth order three or lower are around twice as likely to have received ANC than births of order six or higher.

There are substantial residential and regional variations in ANC (Figure 9.2). Mothers living in urban settlements are more likely to have ANC from trained health personnel than those living in rural areas (78 percent and 51 percent, respectively). For almost half of rural births in the five years preceding the survey, the mother did not receive any ANC. Antenatal care coverage exceeds 67 percent in all regions except the East, where it was received by mothers for 38 percent of the births in the five years prior to the survey. The use of antenatal care services is strongly associated with mother's education. Almost all births to women with at least secondary level education received antenatal care, while only one-third of women with less than primary education got any antenatal care.


Table 9.1 Antenatal care

Percent distribution of births in the five years preceding the survey by source of antenatal care (ANC) during pregnancy, according to selected background characteristics, Turkey 1998

Background characteristic	Antenatal care provider ¹					Total
	Doctor	Trained nurse/ Midwife	Traditional birth attendant	No one	Missing	
Age at birth						
<20	54.4	9.1	1.0	35.5	0.1	100.0
20-34	62.6	7.2	0.5	29.3	0.5	100.0
35+	46.3	5.3	0.5	47.6	0.3	100.0
Birth order						
1	72.1	7.1	0.4	20.1	0.3	100.0
2-3	61.7	8.3	0.6	29.0	0.4	100.0
4-5	43.4	7.2	0.8	47.6	0.9	100.0
6+	33.1	3.5	0.5	62.4	0.5	100.0
Residence						
Urban	71.1	6.6	0.4	21.7	0.3	100.0
Rural	42.1	8.5	0.9	47.9	0.6	100.0
Region						
West	79.5	6.0	0.1	13.9	0.4	100.0
South	63.4	8.8	2.0	25.4	0.4	100.0
Central	61.0	12.7	0.2	25.8	0.2	100.0
North	61.5	5.2	0.0	32.6	0.6	100.0
East	34.4	3.7	0.7	60.6	0.6	100.0
Education						
No educ./Pri. incomp.	32.0	5.3	0.5	61.4	0.8	100.0
Pri. comp./Sec. incomp.	66.3	9.1	0.6	23.6	0.3	100.0
Sec. comp./+	93.4	2.5	0.2	3.9	0.0	100.0
Total	60.2	7.3	0.6	31.5	0.4	100.0


¹ If the respondent mentioned more than one provider, only the most qualified provider is considered.

Figure 9.1
Source of Antenatal Care
by Maternal Age and Birth Order



TDHS 1998

Figure 9.2
Antenatal Care
by Region and Residence



Number and Timing of Antenatal Care Visits

Antenatal care is most beneficial when it is sought early in pregnancy and is continued throughout a pregnancy. The first antenatal visit should take place before the third month of pregnancy. The advantage of early detection of pregnancy is that a woman’s normal baseline health status can be assessed; knowledge of a woman’s baseline health will make early diagnosis of any abnormalities easier. The total number of antenatal visits also is important in assessing the adequacy of ANC. According to the recommended schedule, three visits should be made during the first 28 weeks (7th month), with subsequent visits in the 32nd, 36th, and 39th weeks. Regular visits allow proper monitoring of the mother and child throughout pregnancy.

Table 9.2 presents information on the timing and number of visits made to health providers in the five years preceding the survey. The table shows that, although 32 percent of women received no ANC, 42 percent made four or more visits. The median number of ANC visits is 4.2. In nearly 60 percent of the births, ANC was sought before the sixth month of pregnancy. The median number of months of pregnancy for the first antenatal visit is 3 months. It is clear that in Turkey, when mothers seek antenatal care, they go to a provider early in the pregnancy.

Table 9.2. Number of antenatal care visits and stage of pregnancy

Percent distribution of live births in the five years preceding the survey by number of antenatal care (ANC) visits, and by the stage of pregnancy at the time of the first visit, Turkey 1998

Number of visits and stage of pregnancy	Total
Antenatal visits during pregnancy	
None	31.5
1	7.2
2-3 visits	17.6
4+ visits	42.0
Don't know/Missing	1.7
Total	100.0
Median	4.2
Number of months pregnant at the time of first visit	
No antenatal care	31.5
Less than 6 months	57.9
6-7 months	7.2
8+ months	2.2
Don't know/Missing	1.1
Total	100.0
Median	3.0
Number of births	3,459

Tetanus Toxoid Coverage

Tetanus toxoid (TT) vaccination is one of the important preventive measures for neonatal tetanus. According to the Turkish vaccination schedule, two doses of TT are necessary during pregnancy for the full immunisation of unvaccinated woman. However, if a woman has been vaccinated during a previous pregnancy, she may be given one dose for the current pregnancy.

Table 9.3 presents TT coverage for all births in the five years preceding the survey. Among these births, 15 percent had one dose, and 29 percent had two or more doses. These results are similar to those of the 1993 TDHS, with the exception that there has been a 3 percentage point increase in the percentage of births for which the mother received two or more doses.

The variations in TT vaccination coverage according to background characteristics are similar those observed for ANC coverage. As was seen with use of antenatal care, TT coverage is related to mother's age and birth order of the child. Younger women and women of low parity are more likely to have received two doses of tetanus toxoid vaccination.

There are marked differences in TT vaccination coverage among regions, while not much difference is observed between the urban and rural settlements. As it was in TDHS-93, the South had both the highest overall TT coverage and the greatest proportion receiving the second dose.

Table 9.3 Tetanus toxoid vaccinations

Percent distribution of births in the five years preceding the survey by number of tetanus toxoid injections mother received during pregnancy, according to selected background characteristics. Turkey 1998

Background characteristic	Number of tetanus toxoid injections				Total	Number of births
	No injection	One dose	Two doses or more	Don't know/ Missing		
Mother's age at birth						
< 20	52.7	15.3	30.1	1.9	100.0	501
20-34	52.0	14.8	30.4	2.8	100.0	2,706
35+	71.7	9.6	15.2	3.4	100.0	252
Birth order						
1	45.6	14.5	37.2	2.7	100.0	1,192
2-3	50.7	17.4	29.3	2.6	100.0	1,495
4-5	66.1	9.5	21.5	2.9	100.0	435
6+	77.4	8.1	10.9	3.6	100.0	337
Residence						
Urban	53.3	14.8	28.7	3.3	100.0	2,162
Rural	53.9	14.0	30.3	1.8	100.0	1,297
Region						
West	59.8	14.0	22.5	3.6	100.0	1,031
South	31.3	16.9	47.2	4.5	100.0	490
Central	43.2	18.5	36.4	1.9	100.0	795
North	37.0	16.4	43.6	3.1	100.0	271
East	73.1	9.4	16.1	1.4	100.0	871
Education						
No educ./Pri. incomp.	72.4	8.3	15.6	3.6	100.0	936
Pri. comp./Sec. incomp.	45.6	17.4	34.3	2.7	100.0	2,118
Sec. comp./+	50.9	13.4	34.7	1.0	100.0	405
Total	53.5	14.5	29.3	2.7	100.0	3,459

Place of Delivery and Assistance during Delivery

Hygienic conditions during delivery and proper medical attention reduce the health risks of mothers and children. The 1998 TDHS collected information on the place of delivery and the person assisting delivery for all children born in the five years preceding the survey. Overall, 73 percent of all births were delivered at a health facility (Table 9.4), which represents an increase in the level since 1993. Younger women are more likely to deliver at a health facility than older women. Likewise, lower birth order is associated with a greater likelihood of delivery at health facility (Figure 9.3). The level of education is strongly related to the utilisation of health institutions for delivery. The proportion of births delivered in a health facility increases from 45 percent among births to women with no education to 96 percent among births to women with secondary and higher level. Receiving ANC is also associated with the use of health facility at delivery. Women who received ANC during pregnancy are much more likely to deliver in a health facility than women who have had no ANC. Ninety-two percent of women with four or more antenatal visits delivered in a health facility.


Regional and residential differences are also apparent. A child born in an urban area is 1.3 times more likely to have been delivered at health facility than a rural child. In all regions, except the East region (where more than half of the births take place at home), the majority of births are delivered in health facilities.

Table 9.4 Place of delivery

Percent distribution of births in the five years preceding the survey by place of delivery, according to selected background characteristics, Turkey 1998

Background characteristic	Place of delivery			Total	Number of births
	At a health facility	At home	Don't know/ Missing		
Mother's age at birth					
< 20	73.6	25.8	0.6	100.0	501
20-34	73.4	25.8	0.8	100.0	2,706
35+	61.3	37.6	1.2	100.0	252
Birth order					
1	86.9	12.7	0.4	100.0	1,192
2-3	73.5	25.8	0.7	100.0	1,495
4-5	57.6	40.7	1.6	100.0	435
6+	36.8	61.7	1.4	100.0	337
Residence					
Urban	80.2	19.1	0.6	100.0	2,162
Rural	59.7	39.2	1.1	100.0	1,297
Region					
West	86.6	12.7	0.7	100.0	1,031
South	69.2	29.8	1.0	100.0	490
Central	83.3	16.3	0.3	100.0	795
North	83.7	15.5	0.8	100.0	271
East	44.4	54.4	1.2	100.0	871
Education					
No educ./Pri. incomp.	44.5	54.4	1.1	100.0	936
Pri. comp./Sec. incomp.	80.4	18.8	0.8	100.0	2,118
Sec. comp./+	96.2	3.8	0.0	100.0	405
Antenatal care visits					
None	45.4	54.1	0.5	100.0	1,090
1-3 visits	74.8	24.8	0.4	100.0	859
4+ visits	91.9	7.9	0.2	100.0	1,451
DK/missing	64.4	7.6	28.0	100.0	59
Total	72.5	26.7	0.8	100.0	3,459

Figure 9.3
Place of Delivery by
Maternal Age and Birth Order



The type of assistance a woman receives during the birth of her child depends to great extent on the place of delivery, with births delivered outside the health facility being much less likely to receive assistance from a doctor or other trained health professional. The proportion of all births delivered with the assistance of a doctor or trained health personnel was 81 percent (Table 9.5). As was the case in the TDHS-93, the likelihood of having a birth assisted by qualified health personnel is greater than the likelihood of receiving ANC from a medical care provider (68 percent). A positive relationship is observed between the percentage assisted by medical personnel and the number of antenatal care visits. Medical personnel assisted ninety-six percent of births to women who had at least four antenatal care visits at delivery.

Maternal age and child's birth order, are associated with type of assistance at delivery. Older women and women who have already had a number of births are less likely to receive assistance from medical personnel. Mother's education is also closely tied to medical supervision at delivery. Almost all women with secondary or more education received medical assistance at delivery.

In addition to the regional and residential variation in assistance during delivery, there are differences in the percentage of deliveries assisted by a doctor. For instance, in the East region, where around half of the deliveries were assisted by medical personnel, doctors assisted with 17 percent of the births. More deliveries are attended by doctors in the West region (64 percent), than other regions. The likelihood of delivery under a doctor's supervision is two times greater for urban women compared to rural women.

Table 9.5 Assistance during delivery

Percent distribution of births in the five years preceding the survey by type of assistance during delivery and place of delivery, according to selected background characteristics, Turkey 1998

Background characteristic	Attendant assisting during delivery						Total	Place of delivery			Total	Number of births
	Doctor	Nurse/Trained midwife	Trained TBA ¹	Relative/Other	No one	Don't know/Missing		Health facility	At home	Don't know/Missing		
Mother's age at birth												
< 20	33.9	46.7	10.6	8.1	0.6	0.1	100.0	73.6	25.8	0.6	100.0	501
20-34	41.4	40.4	9.6	7.2	0.6	0.7	100.0	73.4	25.8	0.8	100.0	2,706
35+3	9.2	28.4	18.4	10.3	3.7	0.0	100.0	61.3	37.6	1.2	100.0	252
Birth order												
15	1.7	40.4	4.5	2.8	0.1	0.5	100.0	86.9	12.7	0.4	100.0	1,192
2-3	40.1	43.2	8.9	6.7	0.6	0.5	100.0	73.5	25.8	0.7	100.0	1,495
4-5	26.5	41.1	16.5	14.0	0.8	1.2	100.0	57.6	40.7	1.6	100.0	435
6+	17.3	27.5	30.4	20.3	3.9	0.5	100.0	36.8	61.7	1.4	100.0	337
Residence												
Urban	49.6	38.1	7.2	4.0	0.5	0.6	100.0	80.2	19.1	0.6	100.0	2,162
Rural	24.3	44.4	15.7	13.6	1.3	0.7	100.0	59.7	39.2	1.1	100.0	1,297
Region												
West	63.9	28.4	3.4	3.2	0.3	0.8	100.0	86.6	12.7	0.7	100.0	1,031
South	34.7	51.5	9.6	3.1	0.7	0.4	100.0	69.2	29.8	1.0	100.0	490
Central	38.6	51.4	4.4	4.8	0.8	0.2	100.0	83.3	16.3	0.3	100.0	795
North	39.7	50.0	7.1	2.2	0.3	0.6	100.0	83.7	15.5	0.8	100.0	271
East	16.7	35.6	25.7	19.6	1.5	0.8	100.0	44.4	54.4	1.2	100.0	871
Education												
No educ./Pri. incomp.	19.0	34.8	24.7	18.7	1.9	0.9	100.0	44.5	54.4	1.1	100.0	936
Pri. comp./Sec. incomp.	43.6	45.4	6.0	4.1	0.5	0.4	100.0	80.4	18.8	0.8	100.0	2,118
Sec. comp./+	70.9	27.7	0.2	0.2	0.0	1.0	100.0	96.2	3.8	0.0	100.0	405
Antenatal care visits												
None	16.3	40.9	22.6	18.4	1.6	0.2	100.0	45.4	54.1	0.5	100.0	1,090
1-3 visits	34.4	50.2	10.4	4.1	0.8	0.1	100.0	74.8	24.8	0.4	100.0	859
4+ visits	61.2	35.1	1.5	1.7	0.2	0.3	100.0	91.9	7.9	0.2	100.0	1,451
DK/missing	46.9	23.2	3.1	3.0	0.0	23.8	100.0	64.4	7.6	28.0	100.0	59
Total	40.1	40.5	10.4	7.6	0.8	0.6	100.0	72.5	26.7	0.8	100.0	3,459

¹ Traditional birth attendant

Characteristics of Delivery

Respondents were asked, whether the delivery was by caesarean section or not. They were also asked, if their children were weighed at the time of birth and, if so, how much each baby weighed. In addition, mothers were asked for a subjective assessment of their baby's size at birth.

Table 9.6 indicates that, according to the mother's reports, 14 percent of babies born in the five years preceding the TDHS-98 were delivered by caesarean section. Caesarean sections are more common among births to older women, women residing in urban areas, and more educated women. It is also noteworthy that nearly one-fifth of first births in the five-year period preceding the survey were delivered by caesarean section.

For 64 percent of babies born in the five years preceding the survey, a birth weight was reported. Among births for which a birth weight was reported, 12 percent (8 percent of all births) were reported to have a weight less than 2.5 kilograms. Classified as low-birth-weight births, these children are considered to have a higher than average risk of early mortality. The mother's subjective evaluation of the baby's size at birth is also shown in Table 9.6. According to the mother's subjective evaluation of birth size, 10 percent of all births were reported to be very small and 16 percent were considered to be smaller than average.

Table 9.6 Delivery characteristics: caesarean section, birth weight and size

Among births in the five years preceding the survey, the percentage of deliveries by caesarean section (C-section), and the percent distribution by birth weight and by the mother's estimate of baby's size at birth, according to selected background characteristics, Turkey 1998

Background characteristic	Delivery by C-section	Birth weight			Size of child at birth				Number of births
		Less than 2.5 kg	2.5 kg or more	Don't know/ Missing	Very small	Smaller than average	Average or larger	Don't know/ Missing	
Age at birth									
<20	7.3	8.9	50.2	40.9	12.1	20.9	65.0	1.9	501
20-34	14.8	7.9	58.6	33.5	9.5	15.4	73.8	1.2	2,706
35+	17.2	6.2	42.1	51.7	11.0	14.7	73.9	0.4	252
Birth order									
1	18.8	11.1	67.9	21.0	12.0	18.9	67.7	1.4	1,192
2-3	13.7	6.7	60.0	33.3	7.6	14.0	77.7	0.8	1,495
4-5	8.9	6.6	38.0	55.4	13.0	16.4	68.4	2.1	435
6+	3.3	3.3	21.8	74.9	9.5	16.1	72.7	1.6	337
Residence									
Urban	17.6	8.2	65.2	26.6	8.8	14.1	76.0	1.1	2,162
Rural	7.7	7.3	41.3	51.4	11.9	19.6	66.9	1.5	1,297
Region									
West	21.9	8.4	78.3	13.3	8.2	13.1	76.9	1.9	1,031
South	14.4	9.2	53.7	37.1	10.6	13.1	75.2	1.1	490
Central	11.8	10.7	65.3	24.0	8.9	15.9	74.2	1.0	795
North	14.3	7.5	73.2	19.3	7.1	13.2	78.9	0.8	271
East	5.7	4.0	18.0	78.0	13.7	22.8	62.5	1.0	871
Education									
No educ./Pri.incomp.	5.8	5.0	23.3	71.7	12.5	19.5	66.3	1.7	936
Pri. comp./Sec. incomp.	13.7	9.2	64.8	26.1	9.6	15.7	73.4	1.3	2,118
Sec. comp./+	33.3	7.9	87.7	4.4	5.9	11.4	82.8	0.0	405
Total	13.9	7.9	56.2	35.9	10.0	16.2	72.6	1.3	3,459

Delivery Complications

Table 9.7 shows that mothers reported some complication at the time of delivery in the case of 56 percent of births in the five-year period preceding the survey. Mothers reported prolonged labour in the case of 37 percent of the births, 25 percent involved convulsions, 21 percent involved vaginal infection, and 16 percent involved excessive bleeding. The table also indicates that nearly half of the births who received either delivery care or both the antenatal and delivery care did not involve complications at delivery. Among those deliveries where there was not medical assistance or ANC, prolonged labour and vaginal infections were the most frequently cited complication. Excessive bleeding, convulsions and vaginal infection were the most often cited complications among women whose offspring succumbed to early neonatal death. The percentages of women who had complications during delivery were lower for the women who delivered by caesarean section.

Table 9.7 Delivery complications

Percent of live births in the last five years with complications of delivery, according to antenatal and delivery care, Turkey 1998

Antenatal/ delivery care	Complications at delivery					Number of births
	Prolonged labor	Excessive bleeding	Vaginal infection	Convulsions	None	
Medical maternity care						
Both	34.5	13.9	17.2	24.4	45.4	2,163
Antenatal	48.5	24.5	34.9	35.5	29.1	173
Delivery	38.2	15.8	20.4	20.6	46.9	636
None	42.0	25.1	33.7	31.6	39.4	488
Early neonatal death						
No	37.0	16.1	21.0	25.2	44.0	3,383
Yes	35.4	27.9	23.4	27.9	43.2	76
Delivery by caesarean section						
Yes	20.7	9.4	12.8	18.1	61.2	479
No	39.8	17.5	22.4	26.6	40.9	2,954
Missing	19.7	15.4	12.0	6.6	80.3	26
Total	37.0	16.4	21.0	25.2	44.0	3,459

9.2 Immunisation of Children

The World Health Organisation (WHO) guidelines on childhood immunisation call for all children to receive a BCG vaccination against tuberculosis, three doses of DPT vaccine to prevent diphtheria, pertussis, and tetanus; three doses of polio vaccine (OPV); and one dose of measles vaccine before they reach 12 months of age.

In TDHS-98, information on vaccination status was collected for all children born in the five years preceding the survey. However, the data presented here are restricted to children who were alive at the time of the survey fieldwork.

To obtain immunisation data for each eligible child, mothers were asked whether they had a vaccination card for the child, and if so, to show the card to the interviewer. The dates of the vaccinations were copied from the card to the questionnaire. Mothers were also asked whether the child has been given any vaccination not recorded on the card. If the vaccination card was not available for the child, the mother was asked a number of questions in order to determine the vaccination status of the child for each specific vaccine. In case of DPT and polio, the mother was asked to report the number of doses of the vaccine that the child had received. Children who received one dose of BCG, three doses of DPT and OPV, and one dose of measles vaccine were considered to be fully vaccinated.

Coverage of Children Age 12-23 Months

Information on vaccination coverage is presented in Table 9.8 according to the source of information used to determine coverage, i.e., the child's vaccination card or the mother's report. Data are presented for children age 12-23 months, by which age the child should be fully vaccinated.

The information was gathered from a vaccination card in the case of 39 percent of children while mothers supplied the information for the remaining cases. For children whose information was based on the mother's report, the proportion vaccinated during the first year of life is lower than that for children with a written record of vaccination. The OPV coverage rate for children without a written record is somewhat higher than that of the DPT coverage rate.

Taking into account both the card information and the mother's report, Table 9.8 shows that 46 percent of the children had received all of the recommended immunisations at some time before the survey. Only 4 percent had not received any vaccination at all. The remaining 50 percent were partially vaccinated. The percentage of children who were fully immunised by 12 months of age was 40 percent.

Table 9.8 Vaccinations by source of information

Percentage of children 12-23 months who have received specific vaccines at any time before the survey, by source of information, and the percentage vaccinated by 12 months of age, Turkey 1998

Source of information	Percentage of children who received:										Number of children
	BCG	DPT			Polio			Measles	All ¹	None	
		DPT1	DPT2	DPT3	Polio1	Polio2	Polio3				
Vaccinated at any time before the survey											
Vaccination card	36.0	38.0	36.2	34.5	38.3	36.7	35.6	33.3	29.1	0.0	266
Mother's report	52.5	48.9	38.1	24.1	55.8	50.5	28.8	45.2	16.6	3.6	424
Either source	88.5	86.9	74.3	58.7	94.0	87.2	64.4	78.5	45.7	3.6	689
Vaccinated by 12 months of age²	87.4	85.5	72.4	55.8	92.4	85.2	60.8	70.9	40.4	4.8	689

¹ Children who are fully vaccinated (i.e., those who have received BCG, measles and three doses of DPT and polio).
² For children whose information was based on the mother's report, the proportion of vaccinations given during the first year of life was assumed to be the same as for children with a written record of vaccination.

Coverage Rates by Background Characteristics

Vaccination coverage rates for children in the 12-23 month age group are presented in Table 9.9 by background characteristics. There are definite differences in vaccination coverage by urban-rural residence. The percentages of children receiving the first two doses of OPV are similar for both urban and rural children. However the percentages receiving the first two DPT doses are higher for urban children than for rural children. As a result of high drop-out rates, coverage in rural children for the third dose of DPT/OPV falls to 50 and 57 percent, respectively. BCG and measles coverage rates are also lower for rural children than for urban children. Overall, around half of the urban children are fully vaccinated which is much higher than the proportion for rural children. (37 percent).

Considering regional differences, the percentage fully immunised is significantly lower in the East (23 percent), followed by the West and Central regions (50 and 52 percent, respectively). The North Region has the highest vaccination coverage; 59 percent of children 12-23 months in the North are fully immunised. The data in Table 9.9 also verify the fact that drop-out rate is markedly high in the East compared to other regions. The proportion of children whose mothers showed a vaccination card also was lower in the East (21 percent) than other regions.

The mother's level of education is related to the likelihood that a child will be vaccinated. The percentage of children who are fully vaccinated varies from 29 percent among children whose mothers have no education to 64 percent among children whose mothers have at least secondary education. The DPT/OPV drop-out rates are higher for children of mothers with no education than for other children; for example, DPT coverage rates among children of women with no education fall from 72 percent in the case of the first dose to 45 percent for the third dose. Only 58 percent of children of women with no education received a measles vaccination, and only 70 percent received a BCG vaccination.

A child's birth order is related to coverage rates. Children of high birth order tend to have lower coverage than children of lower birth order. The percentage fully immunised among children of birth order 4-5 is 27 percent. There seems to be little difference between the vaccination levels of male and female children.

Table 9.9 Vaccinations by background characteristics

Percentage of children 12-23 months who had received specific vaccines by the time of the survey (according to the vaccination card or the mother's report), and the percentage with a vaccination card, according to background characteristics, Turkey 1998

Background characteristic	Percentage of children who received:									Percentage with vaccination card		Number of children
	BCG	DPT			Polio			Measles	All ¹	None	card	
		DPT1	DPT2	DPT3	Polio1	Polio2	Polio3					
Child's sex												
Male	89.3	88.3	75.5	60.3	94.1	86.9	66.3	78.9	46.2	3.1	37.7	360
Female	87.7	85.3	73.0	56.9	94.0	87.5	62.2	78.2	45.2	4.2	39.5	329
Birth order												
1	91.2	90.0	78.0	61.9	94.8	88.0	68.4	85.8	51.8	2.6	46.6	246
2-3	93.5	90.3	79.8	62.6	95.5	89.1	68.0	83.4	48.9	2.6	40.5	295
4-5	80.5	77.8	57.7	42.9	90.5	79.9	42.6	57.3	27.2	6.3	23.1	84
6+	66.3	71.0	57.1	49.2	89.2	84.8	60.6	56.3	31.8	8.7	18.9	64
Residence												
Urban	91.6	88.8	78.4	63.9	94.4	88.8	69.0	82.2	51.2	2.9	45.5	424
Rural	83.7	83.7	67.8	50.3	93.4	84.7	57.0	72.7	36.8	4.8	27.5	265
Region												
West	95.8	89.6	79.2	61.4	95.3	91.4	72.3	80.6	50.2	1.6	47.8	221
South	92.7	94.9	85.6	67.4	95.6	89.9	68.9	87.1	57.3	1.5	35.6	95
Central	96.0	95.3	81.5	64.2	98.7	88.4	65.5	86.4	51.8	1.3	41.3	157
North	97.5	95.4	81.9	71.4	98.3	93.5	73.8	84.0	58.9	1.7	52.3	52
East	66.4	67.8	52.1	40.6	85.6	76.8	47.0	61.7	22.9	10.3	20.8	165
Education												
No educ./Pri. incomp.	69.7	72.2	58.1	45.4	88.2	81.4	51.1	57.6	28.5	9.4	17.8	160
Pri. comp./Sec. incomp.	93.3	89.4	76.6	60.0	95.1	87.8	65.9	83.1	48.0	2.2	42.9	434
Sec. comp./+	98.6	99.6	91.1	74.7	99.2	94.2	79.6	93.0	64.0	0.0	53.5	95
Total	88.5	86.9	74.3	58.7	94.0	87.2	64.4	78.5	45.7	3.6	38.5	689

¹ Children who are fully vaccinated (i.e., those who have received BCG, measles and three doses of DPT and polio).

Vaccination in First Year of Life by Current Age

Table 9.10 presents information on children 12-59 months and shows the percentage of children who have a vaccination record as well as the percentage who have received each vaccine during the first year of life according to information from the vaccination card or mother's recall. As was the case in earlier tables, the distribution of vaccinations during the first year of life for children whose information was based on the mother's recall was assumed to be the same as that for children for whom a vaccination record was available.

The proportion of children for whom vaccination cards were seen declines with increasing age of child, from 39 percent among children age 12-23 months to 14 percent among children age 48-59 months. This suggests that there has been an increase in immunization levels in the recent past. However the proportion of children who had received each vaccine during the first year of life are higher for children age 48-59 months than for children in the 24-35- and 36-47-month age groups. With regard to individual vaccines, a substantial increase is observed in the proportion of children receiving the measles vaccine during the first year of life, 71 percent for the children age 12-23 months, compared to less than 60 percent or less among the children age 24-47 months.

Table 9.10 Vaccinations in first year of life by current age

Among children age one to four years, the percentage with a vaccination card and the percentage who have received each vaccine before their first birthday, according to current age of the child, Turkey 1998

Vaccine	Current age of child in months				All children 12-59 months
	12-23	24-35	36-47	48-59	
Vaccination card seen by interviewer	38.5	25.0	12.8	13.5	22.7
Percentage vaccinated at 0-11 months					
BCG	87.4	83.7	76.1	83.8	82.9
DPT 1	85.5	79.6	64.3	77.8	77.1
DPT 2	72.4	65.2	53.0	65.1	64.2
DPT 3	55.8	46.9	41.4	48.0	48.3
Polio 1	92.4	85.3	72.5	88.2	84.9
Polio 2	85.2	77.1	65.4	80.7	77.4
Polio 3	60.8	55.9	49.4	58.3	56.3
Measles	70.9	58.8	52.7	58.0	60.4
All vaccinations ²	40.4	32.8	33.7	35.8	35.8
No vaccinations	4.8	11.8	22.1	9.6	11.8
Number of children	689	612	612	674	2,587

¹ Information was obtained either from a vaccination card or from the mother if there was no written record. For children whose information was based on the mother's report, the proportion of vaccinations given during the first year of life was assumed to be the same as that for children with a written vaccination record.

² Children who have received BCG, measles, and three doses each of DPT and polio vaccines

9.3 Prevalence and Treatment of Diarrhoea

Dehydration brought on by severe diarrhoea is an important cause of morbidity and mortality. In Turkey, the National Control of Diarrhoeal Diseases Programme was established in 1986. The main objective of the programme was reduction in deaths by prevention of dehydration. For this reason, Oral Rehydration Therapy (ORT) has been taught actively since the 1980s.

In the TDHS-98, mothers who had children under age five were asked if their children experienced diarrhoea during the two weeks before the survey. Mothers were also asked what treatment they had given to those children who had diarrhoea. The timing of the TDHS-98 fieldwork should be taken into consideration when assessing these findings, since the fieldwork took place mainly between August and October. As the prevalence of diarrhoea varies seasonally, the results do not represent the average prevalence of diarrhoea throughout the year in Turkey.

Table 9.11 and Figure 9.4 show the percentage of children under five years of age with diarrhoea during the two weeks preceding the survey. Overall 30 percent of children had experienced diarrhoea at some time in the two weeks preceding the survey. There has been a slight increase in the prevalence of diarrhoea since the TDHS-93, where the two-week prevalence of diarrhoea for the same period was 25 percent. Bloody stools were reported by only a small fraction of children (3 percent). Children under two years of age, especially those 6-11 and 12-23 months old, are more likely than older children to have had diarrhoea. This is a pattern which is observed in many surveys including the previous quinquennial surveys, and it is believed to be associated with the effects of weaning practices and poor sanitation, especially the use of contaminated water supplies.


Table 9.11 Prevalence of diarrhoea

Percentage of children under five years who had diarrhoea in the two weeks preceding the survey, and the percentage of children who had diarrhoea with blood in the stools, by selected background characteristics, Turkey 1998

Background characteristic	Diarrhoea in the preceding two weeks		Number of children
	All diarrhoea	Diarrhoea with blood	
Child's age			
Under 6 months	26.3	2.4	364
6-11 months	44.1	4.7	348
12-23 months	43.7	4.5	689
24-35 months	31.1	3.9	612
36-47 months	21.7	2.6	612
48-59 months	15.8	1.8	674
Child's sex			
Male	31.9	3.2	1,713
Female	27.3	3.3	1,586
Birth order			
1	29.2	2.6	1,145
2-3	28.0	2.7	1,432
4-5	34.8	6.6	411
6+	32.9	4.1	310
Residence			
Urban	26.1	2.6	2,081
Rural	35.7	4.5	1,217
Region			
West	21.8	1.5	995
South	28.7	3.8	472
Central	31.0	3.2	762
North	26.4	1.7	260
East	39.7	5.7	810
Education			
No educ./Pri. incomp.	36.1	5.7	875
Pri. comp./Sec. incomp.	29.6	2.8	2,026
Sec. comp./+	16.1	0.6	398
Total	29.7	3.3	3,299

Note: Figures are for children born in the period 1-59 months

Figure 9.4
Percentage of Children Under Five Years with Diarrhoea,
by Age, Sex, Birth Order and Residence



TDHS 1998

There is little variation in diarrhoea prevalence by sex and birth order, with the prevalence of diarrhoea being only slightly higher for male children and for children with higher birth orders than for other children. The prevalence of diarrhoea was higher among rural children (36 percent), children in the East (40 percent) and children whose mothers did not attend school (36 percent) than among other children.

Findings related to the treatment of diarrhoea are presented in Table 9.12. The majority of children under five years of age with diarrhoeal episodes received some kind of treatment. Almost 70 percent of the mothers report increasing the fluids given the child during the diarrhoeal episode or using some form of oral rehydration therapy. Mothers reported that in 37 percent of the diarrhoeal episodes children were taken to a health facility. However the proportion of children who were taken to a health facility is higher in urban areas than in rural areas (43 percent and 29 percent, respectively). Urban mothers were also more likely than rural mothers to use some form of oral rehydration therapy (ORT) or to increase fluids. Among the regions, the South has the highest percentages in seeking health care (48 percent). The use of home remedies or other treatments is widespread.

Table 9.12 Treatment of diarrhoea

Among children under five years who had diarrhoea in the two weeks preceding the survey, the percentage taken for treatment to a health facility or provider, the percentage who received oral rehydration therapy (ORT) (either solution prepared from ORS packets, recommended home fluid (RHF), or increased fluids), the percentage who received no ORT, and the percentage given other treatments, according to selected background characteristics, Turkey 1998

Background characteristic	Percentage taken to a health facility or provider ¹	Oral rehydration therapy				Did not receive ORT	Other treatments			Number of children	
		ORS packet	RHF at home	Either ORS or RHF	In-creased fluids		Injec-tion	Home remedy/ Other	No treat-ment		Missing
Child's age											
Under 6 months	31.7	15.8	11.6	23.8	45.9	42.3	2.6	94.7	1.8	0.0	95
6-11 months	43.8	14.9	13.8	26.3	59.9	30.8	5.9	97.8	1.7	0.0	154
12-23 months	44.5	19.0	17.9	34.0	66.7	24.1	3.9	100.0	0.0	0.0	301
24-35 months	33.2	11.9	14.6	25.0	58.1	31.3	4.0	99.3	0.0	0.7	190
36-47 months	22.8	9.3	12.3	21.6	50.9	37.8	1.6	98.3	0.0	0.5	133
48-59 months	30.9	9.1	12.6	20.0	63.3	30.1	4.8	100.0	0.0	0.0	106
Child's sex											
Male	37.9	14.1	13.3	24.9	58.8	31.6	3.9	98.9	0.5	0.0	547
Female	34.9	14.6	16.4	29.4	60.3	29.9	3.9	98.7	0.4	0.5	433
Birth order											
1	39.1	15.6	13.6	27.6	58.1	30.9	2.8	98.4	0.0	0.4	334
2-3	35.6	14.5	16.8	28.3	62.3	28.3	3.5	98.9	0.8	0.0	400
4-5	40.2	14.7	13.5	26.2	59.4	30.5	5.6	98.9	0.7	0.0	143
6+	26.8	8.4	11.6	19.5	52.7	40.9	6.5	99.4	0.0	0.6	102
Residence											
Urban	42.7	16.4	17.0	30.2	62.4	26.9	5.0	99.0	0.3	0.2	544
Rural	28.9	11.6	11.8	22.7	55.7	35.8	2.5	98.5	0.6	0.2	435
Region											
West	37.0	12.8	16.7	27.1	62.8	30.4	3.5	100.0	0.0	0.0	217
South	47.9	10.1	11.8	20.8	57.6	33.8	3.5	97.9	0.5	0.5	136
Central	29.9	16.0	15.1	29.3	57.2	30.4	1.4	97.7	1.1	0.6	236
North	31.6	16.8	18.7	31.5	45.0	37.9	1.0	97.0	0.0	0.0	69
East	37.4	15.3	13.3	26.4	62.7	28.7	6.8	99.5	0.3	0.0	322
Education											
No educ./Pri. incomp.	35.1	12.3	10.9	21.5	54.6	35.9	5.5	98.5	0.3	0.0	316
Pri. comp./Sec. incomp.	35.2	15.5	16.5	29.6	61.8	28.8	3.2	99.1	0.5	0.1	599
Sec comp./+	56.3	12.4	16.1	27.7	61.7	24.7	2.4	97.4	0.0	2.1	64
Total	36.5	14.3	14.7	26.9	59.5	30.8	3.9	98.8	0.4	0.2	979

ORS = Oral rehydration salts

¹ Includes health centre, hospital, clinic, and private doctor

In the TDHS-98, mothers were specifically asked about changes in feeding practices during the diarrhoeal episode. All mothers who had a child with diarrhoea were asked if they had changed the amount of fluids and foods given to the child having the diarrhoeal episode. Among all children with diarrhoea 60 percent received increased fluids and while one-fifth received the same amount as was before the illness (Table 9.13). However, in 17 percent of the cases, the amount of fluid given was reduced.

Table 9.13 Feeding practices during diarrhoea

Percent distribution of children under five years who had diarrhoea in the two weeks preceding the survey, by amount of fluids and solid foods given compared with normal practice, Turkey 1998

Feeding practice	Total
Amount of fluids given	
Same	21.2
Increase	59.5
Decrease	17.1
Don't know/Missing	2.2
Amount of solid foods given	
Same	26.3
Increase	5.3
Decrease	65.6
Don't know/Missing	2.8
Total	100.0
Number	970

CHAPTER 10

INFANT FEEDING, MATERNAL AND CHILDHOOD NUTRITION

Ergül Tunçbilek, Elif Kurtuluş, and Attila Hancıoğlu

This chapter covers three related topics: infant feeding including breastfeeding, supplementary foods and use of a bottle with a nipple while breastfeeding; and anthropometric assessment of the nutritional status of both children under the age of five years and their mothers.

10.1 Breastfeeding and Supplementation

Infant feeding affects both the mother and the child. It affects the child through his/her nutritional status, which in turn has an effect on the risk of dying. The mother is affected through the effect of breastfeeding on the period of postpartum infertility, and, hence, the length of birth interval and fertility levels. These effects are influenced by both the duration and intensity of breastfeeding, and by the age at which the child receives supplemental foods and liquids.

Breastfeeding of infants is surely the most important factor contributing to the maintenance of growth. Breastmilk contains all the nutrients needed by children in the first few months of life. Moreover, breastmilk is clean and always available at just the right temperature, and it promotes a close mother-child relationship. In addition, it provides some immunity to disease through the mother's antibodies and helps in reducing the prevalence of diarrhoea and nutritional deficiencies.

The World Health Organization recommends that children be exclusively breastfed (no other supplementation or plain water) during the first 4 months of life and that children be given solid or mushy supplements beginning with the seventh month of life. Breastfeeding should continue, along with the complementary foods, up to the second birthday or beyond.

As Table 10.1 indicates, breastfeeding is almost universal in Turkey; 95 percent of all children are breastfed for some period of time. Differentials in the proportion of children breastfed are quite small.

Early initiation of breastfeeding is of benefit to both mother and infant. Suckling stimulates production of oxytocin, a hormone that causes the mother's uterus to contract. The first breast milk, colostrum, protects the newborn infant from infections because of its high concentration of antibodies. Information from the TDHS-98 on the timing of initiation of breastfeeding for all children indicates that initiation to breastfeeding is rather late (Table 10.1). Only one-half of ever-breastfed children were started breastfeeding as early as within one hour of birth. A comparison of the 1998 findings with those of the 1993 survey indicates that both percentages of children starting breastfeeding within one hour and one day of life have increased since 1998. While only one-fifth of children were put to breast within one hour in 1993, according to the findings of the TDHS-98, more than half of the children has been put to the breast within one hour. The proportion of children who were put to breast within one day has also increased from 76 percent in 1993 to 85 percent in 1998.

Increases in the early initiation of breastfeeding occurred in all regions and all education groups (data not shown in tables). Despite the increases, marked variations in the timing of initiation of breastfeeding remain between regions and education subgroups. The percentage of children who started breastfeeding within one hour of birth is highest in the North Region (56 percent) and lowest in the East region (45 percent). By education, the percentage initiating breastfeeding within an hour of birth varies

from 45 percent for births to mothers with no education to 65 percent for births to mothers with a secondary education.

The proportion of children who started breastfeeding within one day of birth also varies markedly by region and education. In the East, where mothers are usually less educated and are more likely to give birth without the assistance of a medically trained person, 22 percent of all children were not put to the breast during the first day. Seventy-nine percent of children of mothers with no education started breastfeeding within one day of their birth compared with 89 percent of births to the most highly educated mothers.

Table 10.1 Initial breastfeeding

Percentage of children born in the five years preceding the survey who were ever breastfed, and the percentage of last-born children who started breastfeeding within one hour of birth and within one day of birth, by selected background characteristics, Turkey 1998

Background characteristic	Percentage ever breastfed	Percentage who started breastfeeding:		Number of children
		Within 1 hour of birth	Within 1 day ¹ of birth	
Child's sex				
Male	95.5	51.4	83.7	1,797
Female	94.9	52.2	86.0	1,662
Residence				
Urban	95.8	53.2	85.1	2,162
Rural	94.3	49.4	84.2	1,297
Region				
West	96.5	51.5	86.1	1,031
South	94.9	55.2	86.1	490
Central	95.6	55.7	89.6	795
North	94.5	55.8	82.9	271
East	93.9	45.2	78.5	871
Education				
No educ./Pri. incomp.	94.5	44.5	78.5	936
Pri. comp./ Sec. incomp.	95.3	52.4	86.8	2,118
Sec. comp./+	96.5	65.0	88.7	405
Assistance at delivery				
Medically trained	95.2	53.8	86.7	2,788
Traditional midwife	95.5	45.7	77.8	360
Other/none	95.0	43.0	80.1	290
Missing	*	*	*	21
Place of delivery				
Health facility	95.2	53.5	86.5	2,509
At home	95.5	47.9	81.6	922
Missing	(87.4)	(16.5)	(32.2)	28
Total	95.2	51.8	84.8	3,459

Note: Table is based on all children born in the five years preceding the survey, whether living or dead at the time of the interview. Parentheses indicate that a figure is based on 25-49 respondents. An asterisk indicates a figure is based on fewer than 25 respondents and has been suppressed.

¹ Includes children who started breastfeeding within one hour of birth

The percent distribution of living children by breastfeeding status at the time of the survey is shown in Table 10.2. The child's breastfeeding status is based on information collected in the survey on feeding practices in the last 24 hours before the interview. "Exclusively breastfed" refers to children who receive breastmilk only. "Children who are fully breastfed" includes those who are exclusively breastfed and those who receive only plain water in addition to breastmilk.

Table 10.2 Breastfeeding status
Percent distribution of living children by breastfeeding status, according to child's age in months, Turkey 1998

Age in months	Not breast-feeding	Exclusively breast-fed	Breastfeeding and:		Total	Number of living children
			Plain water only	Supple-ments		
0-1	3.9	14.2	32.6	49.3	100.0	103
2-3	9.8	5.9	29.5	54.8	100.0	141
4-5	17.7	2.0	11.1	69.2	100.0	119
6-7	28.5	0.0	8.4	63.0	100.0	120
8-9	39.7	0.0	1.6	58.7	100.0	129
10-11	49.4	0.0	1.5	49.1	100.0	99
12-13	48.0	0.0	0.0	52.0	100.0	136
14-15	51.0	0.6	0.6	47.9	100.0	128
16-17	64.0	0.0	0.5	35.5	100.0	128
18-19	73.0	0.0	0.0	27.0	100.0	101
20-21	73.2	0.0	0.0	26.8	100.0	102
22-23	86.0	0.0	0.0	14.0	100.0	94
24-25	88.2	0.0	0.0	11.8	100.0	108
26-27	96.5	0.0	0.0	3.5	100.0	103
28-29	96.4	0.0	0.0	3.6	100.0	99
30-31	89.2	0.0	1.7	9.1	100.0	119
32-33	96.3	0.0	0.0	3.7	100.0	77
34-35	97.5	0.0	0.6	1.9	100.0	107
0-3 months	7.3	9.4	30.8	52.5	100.0	245
4-6 months	22.6	1.3	10.3	65.8	100.0	179
7-9 months	35.0	0.0	3.7	61.3	100.0	189

Note: Breastfeeding status refers to preceding 24 hours. Children classified as *breastfeeding and plain water only* receive no supplements.

Table 10.2 indicates that even among children in the first month of life, only 14 percent were exclusively breastfed, a figure lower than that found in the TDHS-93 (19 percent). A substantial proportion of children (47 percent) are fully breastfed (i.e., they receive only water in addition to breastmilk). However, one-half of children (49 percent) are being given other supplements within the first two months of birth. The percentage of children receiving supplements increases to 55 percent among children 2-3 months of age. Early introduction of supplementary food to infant nutrition increases the risk of gastrointestinal infections, which is one of the leading causes of infant mortality in Turkey.

Table 10.3 shows the median duration of any, exclusive and full breastfeeding. The median duration of breastfeeding for all children is 12 months, which is unchanged from the median reported in 1993. There are some differences in breastfeeding durations among subgroups. With respect to the sex of the child, there is a more than a three month difference in favour of males. Women living in the East are breastfeeding their children 16 months, longer than any other region. In the Central and North regions, median durations shorter than 9 months are observed. Women who never attended school are breastfeeding for nearly 17 months, at least 7 months longer on average than more educated women. Median durations for

exclusive breastfeeding are very short, around half a month (0.4) for all subgroups. There is greater variation in the median duration of full breastfeeding. Children living in urban areas, in the Central region, and those who are assisted by traditional midwife at delivery are likely to have a somewhat longer period of full breastfeeding.

Table 10.3 also presents findings on the frequency of breastfeeding, as indicated by the percentage of children under 6 months of age who were breastfed six or more times in the 24 hours preceding the survey. Seventy-nine percent of children under 6 months of age were breastfed 6 times or more in the 24-hour period preceding the survey. The lowest percentage of children breastfed 6 times or more is found in the North region (67 percent) and the highest percentage among children assisted by traditional midwife at delivery (90 percent).

Table 10.3 Median duration and frequency of breastfeeding

Median duration of any breastfeeding, exclusive breastfeeding, and full breastfeeding among children under three years of age, and the percentage of children under six months of age who were breastfed six or more times in the 24 hours preceding the interview, according to background characteristics, Turkey 1998

Background characteristic	Median duration of breastfeeding			Children under six months		
	Any breast-feeding	Exclusive breast-feeding	Full breast-feeding ²	Number of children under 3 years of age	Breastfed 6 or more times in preceding 24 hours	Number of children
Child's sex						
Male	13.9	0.4	0.7	1,103	77.3	174
Female	10.5	0.5	0.7	1,006	79.5	189
Residence						
Urban	10.7	0.5	1.3	1,308	80.0	231
Rural	13.9	0.4	0.5	801	75.7	132
Region						
West	12.2	0.4	0.6	628	72.7	102
South	11.0	0.4	1.1	295	82.5	51
Central	8.1	0.5	1.3	491	81.3	101
North	6.8	0.4	0.6	162	(66.9)	28
East	16.2	0.5	0.7	532	83.4	82
Education						
No educ./Pri.incomp.	16.5	0.4	0.7	531	76.1	82
Pri. comp./Sec.incomp.	9.5	0.5	0.7	1,319	78.5	237
Sec. comp./+	8.3	0.4	0.9	258	82.5	45
Assistance at delivery						
Medically trained	10.8	0.4	0.7	1,740	77.3	314
Traditional midwife	17.9	0.5	3.1	189	(89.6)	28
Other or none	18.6	0.4	0.4	166	*	22
Missing	*	*	*	13	-	-
Total	12.0	0.4	0.7	2,108	78.5	364
Mean	13.7	1.1	2.7	95.2	-	-
Prevalence/incidence mean	13.9	0.4	2.2	-	-	-

Note: Figures in parentheses are based on 25 to 49 children; an asterisk indicates that a figure is based on fewer than 25 children and has been suppressed.

² Medians and means are based on current status and durations are in months.
² Either exclusive breastfeeding or breastfeeding and plain water only

Table 10.4 shows the percentage of breastfeeding and non-breastfeeding children who received different types of supplements. Because children may have received more than one type of supplement, the percentages do not add to 100. Only 10 percent of breastfeeding children, age 0-3 months received no other supplements. One in every five infants in this age group received infant formula. The percentage who were given infant formula peaks at 32 percent for infants age 4-6 months and then decreases to 21 percent for those age 7-9 months who were increasingly being given other fluids. For non-breastfeeding children, the number of observations are small; however, the results for this group also indicate that infant formula is commonly given in the first months of life, with other types of milk being given more often as a baby grows older.

In Table 10.4, the extent to which bottles are used to feed infants is also presented. Using a bottle with a nipple increases the state of being under the risk of gastrointestinal infections for children. More than one-third of breastfeeding children age 0-3 months are fed by bottle with a nipple.

In summary, although breastfeeding is universal in Turkey, exclusive breastfeeding is not widely practiced. Early introduction of infant formula and other liquids is common, and bottle-feeding is a popular feeding practice. Results of the TDHS-98 imply that the efforts of national and international organisations to promote appropriate infant feeding practices must be increased.

Table 10.4 Types of food received by children in the preceding 24 hours

Percentage of children under three years of age who received specific types of food in the 24 hours before the interview, and the percentage using a bottle with a nipple, by breastfeeding status and child's age in months, Turkey 1998

Age (in months)	Breast milk only	Infant formula	Other milk	Other liquids	Other	Use of bottle with a nipple	Number of children
BREASTFEEDING CHILDREN							
0-3	10.2	19.6	13.1	38.8	2.9	36.9	227
4-6	1.7	32.1	20.6	78.4	28.6	46.4	139
7-9	0.0	21.2	35.7	89.3	55.1	46.4	123
10-11	0.0	10.3	42.6	93.7	79.6	38.6	50
12-13	0.0	9.2	27.8	98.9	65.2	15.6	71
14-15	1.1	8.4	42.9	94.9	80.7	25.4	63
16-17	(0.0)	(1.5)	(29.9)	(98.7)	(67.0)	(21.5)	46
18-23	0.0	3.5	27.2	97.5	77.5	28.0	68
24-29	*	*	*	*	*	*	20
30-35	*	*	*	*	*	*	18
Total 0-35	3.2	16.5	25.8	76.4	43.5	34.8	824
NON-BREASTFEEDING CHILDREN							
0-3	*	*	*	*	*	*	18
4-6	(0.0)	(35.7)	(57.7)	(84.7)	(32.1)	(80.6)	41
7-9	0.0	29.0	59.4	95.1	73.9	90.1	66
ALL CHILDREN							
Total 0-35	0.0	10.0	48.2	97.2	81.9	49.8	1,189

Note: Total for non-breastfeeding children includes children age 0-17 months. Parentheses indicate that a figure is based on 25-49 respondents. An asterisk indicates a figure is based on fewer than 25 respondents and has been suppressed.

10.2 Nutritional Status

One of the major contributions of the TDHS to the study of child health status is the anthropometric data collected for all children under five years of age. Both weight and height (length) measurements are obtained for each child. Employing this information, standard indices are used to describe the nutritional status of the children: height-for-age, weight-for-height, and weight-for-age.

In presenting the anthropometric results, the nutritional status of children in the survey population is compared against an international reference population defined by the U.S. National Center for Health Statistics (NCHS) and accepted by the U.S. Centers for Disease Control (CDC) and the World Health Organization (WHO). The use of the international reference population is based on the finding that well-nourished young children of all population groups (for which data exist) follow very similar growth patterns before puberty.

In any large population, there is obviously a natural variation in height and weight. This variation approximates a normal distribution. The reference population serves as a point of comparison, facilitating the examination of differences in the anthropometric status of subgroups in a population and of changes in nutritional status over time.

The height-for-age index presented in Tables 10.5 and 10.6 provides an indicator of linear growth retardation among children. Children who are more than two standard deviations below the median of the reference population in terms of height-for-age may be considered short for their age ("stunted"), or chronically undernourished. Children who are below minus three standard deviations (-3 SD) from the median of the reference population are considered severely stunted. Stunting reflects the outcome of a failure to receive adequate nutrition over a long period of time, and is also affected by recurrent and chronic illness. Thus, height-for-age, represents a measure of the long-term effects of undernutrition in a population and does not vary appreciably according to the season of data collection. Stunted children are not immediately obvious in a population; a stunted three-year-old child could look like a well-fed two-year-old.

The weight-for-height index measures body mass in relation to body length. Children who are more than two standard deviations below the median of the reference population in terms of their weight-for-height may be considered too thin ("wasted") or acutely undernourished. Severe wasting represents the failure to receive adequate nutrition in the period immediately before the survey and may be the result of recent illness episodes, especially diarrhea, or of seasonal variations in food supply.

Weight-for-age takes into account both acute and chronic undernutrition and often is used to monitor nutritional status on a longitudinal basis. It is a useful tool in clinical settings for continuous assessment of nutritional progress and growth. Children whose weight-for-age is below minus two standard deviations from the median of the reference population are classified as "underweight." In the reference population, only 2.3 percent of children fall below minus two (-2 SD) for each of the three indices.

Table 10.5 shows how the percentage of children under five years of age classified as undernourished according to the height-for-age, weight-for-height, and weight-for-age indices varies with the child's age and selected demographic characteristics.

Table 10.5 Nutritional status of children by background characteristics

Percentage of children under five years of age who are classified as malnourished according to three anthropometric indices of nutritional status: height-for-age, weight-for-height, and weight-for-age, by selected background characteristics, Turkey 1998

Background characteristic	Height-for-age		Weight-for-height		Weight-for-age		Number of children
	Percentage below -3 SD	Percentage below -2 SD ¹	Percentage below -3 SD	Percentage below -2 SD ¹	Percentage below -3 SD	Percentage below -2 SD ¹	
Child's age							
Under 6 months	0.4	2.0	0.2	2.1	0.5	1.7	285
6-11 months	1.0	4.6	0.9	3.7	0.9	7.0	298
12-23 months	4.9	16.7	0.6	2.9	2.6	10.5	547
24-35 months	7.1	17.4	0.3	1.4	1.7	8.2	481
36-47 months	10.8	20.1	0.3	1.4	1.8	9.7	496
48-59 months	7.8	23.7	0.0	0.7	0.5	9.0	570
Sex of child							
Male	5.8	16.0	0.6	2.1	1.7	8.4	1,415
Female	6.4	16.0	0.1	1.7	1.0	8.1	1,261
Birth order							
1	4.1	13.5	0.4	1.8	1.0	7.5	929
2-3	5.7	14.2	0.3	2.0	1.2	7.0	1,172
4-5	7.1	17.5	0.8	2.4	2.5	10.2	329
6+	14.1	32.2	0.0	1.2	2.5	14.8	247
Birth interval							
First birth	4.0	13.4	0.4	1.9	1.0	7.6	938
Under 24 months	12.2	24.9	0.5	2.6	2.3	12.3	426
24-47 months	7.4	18.3	0.4	1.6	2.0	8.8	687
48+ months	3.7	11.3	0.1	1.6	0.7	5.9	625
Residence							
Urban	4.7	12.6	0.3	1.7	0.9	6.2	1,696
Rural	8.4	22.0	0.4	2.3	2.3	11.9	981
Region							
West	3.6	9.9	0.2	1.5	0.2	3.8	763
South	4.3	13.5	0.3	2.2	1.7	8.9	407
Central	4.3	11.6	0.1	1.3	1.3	5.4	643
North	2.9	12.8	0.0	1.6	0.6	4.8	211
East	12.9	30.0	0.9	2.9	3.0	17.1	652
Education							
No educ./Pri. incomp.	14.4	31.0	0.4	2.4	3.4	17.2	710
Pri. comp./Sec. incomp.	3.5	11.8	0.3	1.7	0.8	5.6	1,652
Sec. comp./+	1.1	4.0	0.5	1.8	0.2	2.4	315
Total	6.1	16.0	0.4	1.9	1.4	8.3	2,677

Note: Figures are for children born in the period 0-59 months preceding the survey. Each index is expressed in terms of the number of standard deviation (SD) units from the median of the NCHS/CDC/WHO international reference population. Children are classified as malnourished if their z-scores are below minus two or minus three standard deviations (-2 SD or -3 SD) from the median of the reference population.

¹ Includes children who are below -3 SD

Table 10.6 Anthropometric indicators of maternal nutritional status


Percent distribution and mean and standard deviation for women who had a birth in the five years preceding the survey, by selected anthropometric indicators (height, weight, and body mass index (BMI)), Turkey 1998

Indicator	Total	Total plus missing
Height (cm)		
135.0-139.9	0.2	0.2
140.0-144.9	1.9	1.8
145.0-149.9	11.1	10.7
150.0-154.9	28.0	27.0
155.0-159.9	32.4	31.3
160.0-164.9	20.7	19.9
165.0-169.9	4.9	4.8
170.0-174.9	0.7	0.7
>= 180.0	0.1	0.1
Missing	-	3.6
Total	100.0	100.0
Mean	156.2	-
Standard deviation	5.7	-
Number of women	2,524	2,617
Weight (kg)		
35.0-39.9	0.3	0.3
40.0-49.9	12.4	11.9
50.0-59.9	32.5	31.2
60.0-69.9	28.6	27.5
>= 70.0	26.2	25.2
Missing	-	3.9
Total	100.0	100.0
Mean	63.3	-
Standard deviation	12.5	-
Number of women	2,191	2,281
BMI (kg/m²)		
12.0-15.9 (Severe)	0.1	0.1
16.0-16.9 (Moderate)	0.2	0.2
17.0-18.4 (Mild)	2.3	2.2
18.5-20.4 (Normal)	8.7	8.3
20.5-22.9 (Normal)	19.5	18.7
23.0-24.9 (Normal)	17.0	16.3
25.0-26.9 (Overweight)	16.6	15.9
27.0-28.9 (Overweight)	11.6	11.1
29.0-29.9 (Overweight)	5.2	5.0
>= 30.0 (Obese)	18.8	18.0
Missing	-	4.3
Total	100.0	100.0
Mean	26.0	-
Standard deviation	5.0	-
Number of women	2,183	2,281

Note: The weight and BMI measures exclude pregnant women and those who are less than 3 months postpartum.

In the TDHS, all children under five years of age whose mother was interviewed are included in the anthropometric data collection. However, not all eligible children are included in the results presented here; height or weight measurements are missing for 18 percent of eligible children (see Appendix D). In addition, since two of the indices (height-for-age and weight-for-age) are influenced by the accuracy of the reporting of the child's age, a small number of children (1 percent) were excluded from the calculation because the month and year of birth was not known. Hence, height and weight data are shown for only 81 percent of the eligible children.

Figure 10.1 shows the extent to which Turkish children deviate at various ages from the reference population in terms of z-scores¹ for the three nutritional indices. The weight-for-height z-score is close to that of the reference population except for children in the second half of the first year, when the z-scores are negative (i.e., the children are thinner). There is a rapid decline in the height-for-age and weight-for age z-scores after the first 6 months of life. Height-for-age continues to decline until the fourth year of life. In contrast, weight-for-age stabilizes around the second birthday.



As a whole, the youngest children show little evidence of undernutrition (Table 10.5). However, the proportion classified as stunted exhibits a steady increase starting in the first year of life. The deterioration in nutritional status continues through the second and third years of life, and, thereafter, appears to reach a plateau. Among children 24-59 months of age, around 20 percent are classified as stunted. According to the survey (Table 10.5), by age 5, around one-quarter of the children are chronically undernourished, with eight percent considered as severely stunted. These patterns reflect inadequate feeding practices and the presence of recurrent and chronic illness.

¹ A z-score provides a measure of the mean number of standard deviation units above or below the median for the reference population that the values for the index for the children in the TDHS sample fall.

The fact that undernutrition percentages increase with the increasing birth order is important. Nearly one-third of children of birth order six or above and almost one-fifth of children of birth orders four and five are stunted. Birth interval is one of the most important variables affecting the height-for-age index. Children who are born with an interval of less than two years are much more prone to be stunted. Of these children, 25 percent are stunted and 12 percent are severely stunted.

Overall, wasting is not a problem. Only two percent of children have a weight-for-height z-score below -2SD which is very close to the reference population. However, this figure increases to four percent among children between 6-11 months of age and it is three percent among children age 12-23 months.

Weight-for-age is an index reflecting both height-for-age and weight-for-height. According to the survey results, eight percent of all children are underweight, and one percent are severely underweight. Birth order and birth interval are among the most important factors affecting this index. Fifteen percent of children of birth order six and higher and 12 percent of children born after a birth interval of less than 2 years are classified as undernourished on the weight-for-age index.

Table 10.5 also shows the percentage of children under five years of age classified as undernourished (according to the three anthropometric indices) by selected background characteristics. There are striking differences in the percentage classified as stunted according to the mother's level of education. Undernutrition is not a problem among children of mothers with secondary education or higher; the percentage of children who are below the -2 SD cut-off point (4 percent) is close to that seen for the reference population (2.3 percent). In contrast, almost one-third of children whose mothers lack formal education are classified as stunted. There are also urban-rural and regional differences. Stunting is more common in rural (22 percent) than in urban areas (13 percent). The highest level of stunting is seen in the East region (30 percent) and the lowest levels are in the West and Central regions (10 and 12 percent, respectively). Similar results are observed for the weight-for-height and weight-for-age indices.

In summary, the anthropometric findings from the TDHS-98 suggest that, for many Turkish children, deterioration in nutritional status is observed after 6 months of age. In particular, stunting is a serious problem among children, especially in the East region. A comparison with the 1998 findings with the results of the TDHS-93 indicates that there was little change in the nutritional status of children in Turkey during the five-year period between the surveys. For example, the proportion stunted in 1998 is only slightly lower than the level observed in 1993 (18 percent). Improvements in the nutritional status of Turkish children will be obtained when children are no longer exposed to a number of risk factors, especially short birth intervals and high parity. An intersectoral approach is necessary to discourage mothers from introducing supplementary food too early, to train mothers on the timely introduction of appropriate supplementation, and to assist couples to keep the number of children within their desired limits and ensure optimal birth spacing through effective family planning.

10.3 Maternal Nutrition

In order to assess women's nutritional status, women who had given birth in the five-year period before the interview were weighed by an electronic scale and their heights were measured by a wooden measuring board (i.e., with the same equipment used to obtain children's measurements).

Table 10.6 presents the distribution as well as the means and standard deviations of the anthropometric indicators for eligible women: height, weight, and body mass index. Indicators based on a woman's weight-for-height exclude pregnant women and women with a birth within the 2 months preceding the interview. The table shows that anthropometric measures are available for most eligible women, with height or weight measurements are missing for 4 percent of respondents.

Balanced nutrition during childhood and the adolescent period has a positive impact on linear growth, whereas poor nutrition and experience of a severe illnesses, particularly in early childhood, can affect growth negatively. Maternal height is useful in predicting the risk of delivery complications since short stature is frequently associated with a small pelvis size. The height below which women are considered to be at risk of such complications is in the range of 140-150 centimetres, with 145 centimetres being a standardly accepted cutoff for identifying maternal malnutrition. According to the TDHS-98 results, the mean height for mothers was 156 centimetres, one centimetre higher than the mean reported in the TDHS-93. Two percent of mothers were shorter than 145 centimetres, and 13 percent were below 150 centimetres. The mean maternal weight was 63.3 kilograms. More than one-fourth (26 percent) of mothers weighed more than 70 kilograms.

The body mass index (BMI) assess the relation between height and weight and is calculated by dividing the weight in kilograms by the squared height in metres. A body mass index of less than 18.5 is used to identify cases of chronic undernutrition. A BMI higher than 25.0 is often used to identify women with problems with overweight and obesity, although there is no standard definition of obesity. In the TDHS-98, the mean BMI of nonpregnant mothers was 26. The mothers' BMI fell below 18.5 in 3 percent of cases. Fifty-two percent of the mothers had a BMI above 25.0, including 19 percent who had a BMI of at least 30.

Table 10.7 shows the nutritional status of mothers by selected background characteristics. Younger generations of women appear to be taller than women age 35 and over. The level of education is also directly associated with women's height. Mothers who have had no education or did not complete primary education are, on the average, 3.2 centimeters shorter than those with secondary education or more. Educational level also appears to be related with the body mass index. Mothers with no education had an average BMI of 27 while mothers with secondary or more education had an average BMI of 24.6.

Table 10.7 Nutritional status of mothers by background characteristics

Among women who had a birth in the five years preceding the survey, mean height and percentage under 145 centimeters, mean body mass index (BMI) of women, and percentage of women whose BMI is less than 18.5 (kg/m²), by selected background characteristics, Turkey 1998

Background characteristic	Height			BMI		
	Mean	Percentage <145 cm	Number of women	Mean	Percentage <18.5	Number of women
Age						
15-19	156.6	0.0	133	23.0	6.8	102
20-24	156.4	2.3	672	24.7	3.9	562
25-29	156.4	1.5	807	25.7	2.3	697
30-34	156.2	1.6	535	26.7	1.4	481
35-49	155.6	4.2	377	28.5	1.3	341
Residence						
Urban	156.5	1.9	1,643	26.0	2.8	1,442
Rural	155.7	2.4	881	25.8	2.0	741
Region						
West	156.6	2.0	821	25.9	3.1	733
South	156.3	1.5	372	26.2	2.8	321
Central	156.3	1.9	589	25.7	2.6	516
North	156.0	1.9	193	26.4	2.4	166
East	155.6	2.8	549	26.0	1.4	446
Education						
No educ./Pri. incomp.	154.9	4.9	596	27.0	1.8	498
Pri. comp./Sec. incomp.	156.3	1.3	1,586	25.9	2.8	1,373
Sec. comp./+	158.1	0.5	342	24.6	2.7	311
Total	156.2	2.1	2,524	26.0	2.5	2,183

Note: Table includes only women who had a birth in the five years preceding the survey. The BMI index excludes pregnant women and those who are less than three months postpartum.

CHAPTER 11

KNOWLEDGE OF AIDS AND SEXUALLY TRANSMITTED DISEASES

Attila Hancıoğlu and A. Sinan Türkyılmaz

The TDHS-98 included questions to assess the knowledge of sexually transmitted diseases (STDs) and several questions on AIDS to assess the knowledge and attitudes of respondents regarding transmission mechanisms and prevention of infection with the AIDS virus. All women age 15-49 years and eligible husbands were asked whether they had ever heard of AIDS, and if so, what were their sources of information concerning prevention and treatment of the disease, and what were their personal perceptions about the risk of getting the disease.

11.1 Source of Information about AIDS

Tables 11.1.1 and 11.1.2 show the percentage of all women and the husbands of currently married women who have heard of AIDS by source of information, according to background characteristics. In the TDHS-98, a respondent may report having heard about AIDS from more than one source. Overall, 84 percent of women and 93 percent of husbands have heard about AIDS. Regional differentials are significant in this respect. While 92 percent of women living in the West region have heard of AIDS, the figure declines to 58 percent in the East region. Although the West-East differential is not as great for husbands, there is still a significant differential (97 versus 78 percent). As expected, urban women and husbands are more knowledgeable about AIDS than their rural counterparts.

Eighty percent of all women received information about AIDS from television, 28 percent from newspapers or magazines, 11 percent from friends or relatives, and 10 percent from radio broadcasts. Sources of information for husbands appear to have a slightly different pattern; 85 percent of husbands received information about AIDS from television, 54 percent from newspapers or magazines, 22 percent from radio broadcasts, and 14 percent received from friends or relatives. For both women and husbands, the television is the leading source of information on AIDS. For husbands, newspapers are also a common source of information on AIDS. It is notable that neither schools, nor religious institutions are major sources of information.

The percentage of women and husbands who have heard of AIDS varies by age. Older husbands appear to be less knowledgeable about AIDS. For women, though, the age differential is not as great. For both women and husbands, those in their 20s and 30s are more likely to have heard of AIDS than older and younger groups.

As with many other indicators, level of education is closely related to AIDS knowledge. Almost all women with secondary education have heard of AIDS, while this figure declines to 55 percent for women with less than primary education. A similar differential exists for husbands.

More than one-third of currently married women and more than one-quarter of never-married women mentioned having heard of sexually transmitted diseases (STDs) other than AIDS (Table 11.2). STDs are more widely known by husbands: more than half of the husbands reported knowing STDs. Regional and urban-rural differentials in this regard are more pronounced for women than husbands. In the West region, 4 in every 10 of currently married women have knowledge of STDs; in the East region this proportion is as low as 2 in every 10 women. Both women and husbands are more likely to be informed about STDs if they have secondary or higher level education. It is noteworthy that irrespective

Table 11.1.1 Knowledge of AIDS and sources of AIDS information: women

Percentage of all women 15-49 who have ever heard of AIDS, percentage who received information about AIDS from specific sources, and mean number of sources of information about AIDS, by background characteristics, Turkey 1998

Background characteristic	Ever heard of AIDS	Source of AIDS information among those who have heard of AIDS											Number of women	Mean number of sources
		Radio	TV	Newspaper	Pamphlet	Health worker	Mosque/Church	School	Community meeting	Friend/Relative	Work place	Other source		
Age														
15-19	83.4	11.1	78.2	27.8	4.5	2.2	0.0	13.6	0.0	11.7	0.3	4.0	1,720	1.8
20-24	87.9	12.2	83.6	32.0	4.0	3.5	0.0	3.8	0.0	11.9	0.3	3.1	1,558	1.8
25-29	88.0	11.3	83.6	30.6	4.2	5.0	0.0	2.2	0.0	9.7	0.3	3.3	1,397	1.7
30-39	83.6	9.4	80.5	27.2	2.9	3.7	0.0	0.7	0.0	10.6	0.4	3.6	2,283	1.7
40-49	79.5	8.3	74.5	21.9	1.9	2.7	0.1	0.5	0.0	11.3	0.4	3.1	1,618	1.6
Marital status														
Currently married	83.2	9.5	79.5	24.8	2.6	3.5	0.0	1.0	0.0	10.9	0.3	3.1	5,921	1.6
Formerly married	87.4	11.4	79.9	27.3	3.0	5.5	0.0	0.6	0.0	14.5	0.4	3.0	276	1.7
Never married	86.6	12.5	81.2	35.2	5.5	3.0	0.0	12.0	0.0	11.1	0.4	4.2	2,380	1.9
Residence														
Urban	90.3	11.4	86.0	34.2	4.3	4.0	0.0	4.9	0.0	11.9	0.4	3.6	5,704	1.8
Rural	72.2	8.3	68.0	14.9	1.7	2.1	0.0	2.3	0.0	9.3	0.2	3.2	2,872	1.5
Region														
West	92.3	11.3	86.8	36.2	3.9	4.1	0.1	4.6	0.0	12.3	0.2	3.3	3,204	1.8
South	82.5	11.3	78.6	23.9	3.1	3.3	0.0	2.9	0.0	12.0	0.4	3.7	1,258	1.7
Central	90.6	11.3	87.3	26.3	3.9	3.2	0.0	5.2	0.0	9.8	0.4	3.8	1,985	1.7
North	86.4	9.1	83.0	25.8	3.2	4.4	0.0	4.2	0.0	12.7	0.6	3.0	692	1.7
East	58.1	6.8	54.6	15.2	2.1	1.8	0.0	2.1	0.0	8.3	0.3	3.3	1,437	1.6
Education														
No educ./Pri. incomp.	54.9	4.3	50.6	3.3	0.3	1.2	0.0	0.4	0.0	10.1	0.1	2.4	1,861	1.3
Pri. comp./Sec. incomp.	90.2	10.9	86.5	25.3	1.7	2.9	0.0	3.1	0.0	12.0	0.1	2.7	5,158	1.6
Sec.comp./+	99.9	15.8	93.5	65.1	12.9	7.6	0.0	11.6	0.0	8.8	1.3	7.2	1,556	2.2
Total	84.3	10.4	80.0	27.7	3.4	3.4	0.0	4.0	0.0	11.0	0.3	3.4	8,576	1.7

Note: Mean number of sources is based on respondents who have heard of AIDS.

Table 11.1.2 Knowledge of AIDS and sources of AIDS information: husbands

Percentage of husbands who have ever heard of AIDS, percentage who received information about AIDS from specific sources, and mean number of sources of information about AIDS, by background characteristics, Turkey 1998

Background characteristic	Ever heard of AIDS	Source of AIDS information among those who have heard of AIDS											Number of husbands	Mean number of sources
		Radio	TV	News-paper	Pamph-let	Health worker	Mosque/ Church	School	Com-munity meeting	Friend/ Rela-tive	Work place	Other source		
Age														
15-24	90.9	18.6	83.1	49.6	9.1	6.5	0.0	2.1	2.9	15.2	0.0	10.4	112	2.2
25-29	95.5	22.2	89.0	57.5	6.7	2.1	0.7	0.7	1.8	12.5	1.6	7.0	342	2.1
30-39	94.9	24.4	88.0	60.3	5.6	3.0	0.1	0.7	1.5	11.5	1.6	4.3	716	2.1
40-49	92.8	18.0	82.9	50.8	5.9	4.5	0.2	0.7	2.1	15.6	0.9	5.3	575	2.0
50+	84.1	22.3	74.7	39.8	1.0	2.7	1.6	0.0	3.6	16.3	0.2	4.6	226	2.0
Residence														
Urban	95.1	23.8	87.2	62.3	6.7	3.6	0.4	0.8	2.2	13.0	1.5	6.6	1,347	2.2
Rural	88.3	16.7	79.9	36.4	3.0	3.0	0.5	0.5	1.8	15.0	0.5	3.0	624	1.8
Region														
West	97.1	25.8	89.5	62.8	7.0	3.6	0.6	0.4	1.7	6.9	1.4	7.7	767	2.1
South	91.9	25.1	84.8	47.4	3.7	4.7	0.5	1.2	4.3	20.6	2.1	3.7	285	2.2
Central	95.9	18.0	86.2	55.7	5.7	2.6	0.1	1.3	2.0	17.8	0.8	4.0	481	2.0
North	93.3	13.8	86.0	50.1	4.8	3.9	0.0	0.0	1.2	19.3	0.8	3.4	150	2.0
East	77.8	17.0	70.0	37.0	3.6	2.9	0.5	0.3	1.2	14.7	0.4	4.7	287	2.0
Education														
No educ./Pri. incomp	67.8	15.2	56.3	16.1	0.7	2.5	0.9	0.0	2.0	16.4	0.5	2.5	204	1.7
Pri. comp./Sec. incomp.	94.1	20.9	86.9	49.4	3.3	3.3	0.5	0.0	2.3	14.9	1.2	3.5	1,253	2.0
Sec. comp./+	100.0	25.9	91.3	80.8	13.0	4.1	0.0	2.7	1.5	9.3	1.3	11.3	513	2.4
Total	92.9	21.6	84.9	54.1	5.5	3.4	0.4	0.7	2.1	13.6	1.2	5.4	1,971	2.1

Note: Mean number of sources is based on respondents who have heard of AIDS.

Table 11.2 Knowledge of AIDS and sexually transmitted diseases (STDs)

Percentage of currently married women, never-married women, and husbands who have heard about AIDS and about other sexually transmitted diseases (STDs), by selected background characteristics, Turkey 1998

Background characteristic	Currently married women			Never-married women			Husbands		
	Knows about AIDS	Knows about STDs, excluding AIDS	Number of women	Knows about AIDS	Knows about STDs, excluding AIDS	Number of women	Knows about AIDS	Knows about STDs, excluding AIDS	Number of women
Age									
15-19	79.5	19.1	262	84.0	20.8	1,454			
20-24	86.2	28.1	924	90.0	32.1	612	90.9	44.0	112 [†]
25-29	87.6	38.4	1,196	90.0	43.1	181	95.5	50.5	342
30-39	83.2	37.4	2,104	93.8	40.3	104	94.9	55.1	716
40-49	78.2	37.3	1,434	96.5	47.5	28	92.8	55.6	575
50+	-	-	0	-	-	0	84.1	57.5	226
Residence									
Urban	88.9	40.2	3,978	93.7	31.7	1,522	95.1	59.0	1,347
Rural	71.5	25.5	1,943	74.0	17.5	858	88.3	43.4	624
Region									
West	91.0	41.5	2,261	95.9	34.5	819	97.1	57.1	767
South	81.3	33.3	851	84.0	23.0	366	91.9	48.9	285
Central	89.7	35.9	1,426	93.0	24.9	502	95.9	56.6	481
North	84.4	30.7	474	92.1	27.2	198	93.3	54.2	150
East	54.6	23.5	909	64.3	17.5	495	77.8	47.0	287
Education									
No educ./Pri. incomp.	56.0	21.4	1,546	40.5	11.8	226	67.8	37.5	204
Pri. comp./Sec. incomp.	91.2	32.9	3,570	87.4	16.3	1,460	94.1	46.0	1,253
Sec. comp./+	100.0	72.8	804	99.8	52.9	693	100.0	80.3	513
Total	83.2	35.3	5,921	86.6	26.6	2,380	92.9	54.1	1,971

[†] Total includes 3 respondents age 15-19.

of region and place of residence, of educational level and age, AIDS is more widely known by women and husbands. This might be due to the wider coverage of the AIDS disease in the media; it might also be due to the fact that the TDHS-98 questionnaires did not specify STDs but probed respondents to provide names of such diseases. In the case of AIDS, on the other hand, the disease was specifically mentioned by the interviewer.

11.2 Knowledge of Ways to Prevent AIDS

Three percent of women and 2 percent of husbands who have heard of AIDS believe that there are no ways of preventing the contraction of the disease (Tables 11.3.1 and 11.3.2). A further 27 percent of women and 23 percent of husbands were misinformed about the ways of avoiding contraction. Additionally, 30 percent of women and 10 percent of husbands who said that there was a way to avoid getting AIDS then failed to describe ways to avoid AIDS. In other words, although AIDS is generally known by women and husbands, knowledge of ways to avoid it appears to be poor among a substantial minority of both groups.

Table 11.3.1 Knowledge of ways to avoid HIV/AIDS: women

Percentage of all women knowing about AIDS by knowledge of ways to avoid HIV/AIDS, according to selected background characteristics, Turkey 1998

Background characteristic	Ways to avoid AIDS											Don't know specific ways	Percentage with misinformation ¹	Number of women
	No way to avoid AIDS	Abstain from sex	Use condoms	Have only one sex partner	Avoid sex with prostitutes	Avoid sex with homosexuals	Avoid transfusions	Avoid injections	Avoid kissing	Avoid mosquito bite	Other			
Age														
15-19	4.4	7.2	10.4	10.6	8.8	0.2	7.4	2.2	0.6	0.0	26.5	37.8	27.1	1,434
20-24	3.1	5.3	19.1	15.3	14.0	0.8	8.8	2.4	1.4	0.0	26.2	32.5	27.2	1,368
25-29	4.1	4.2	18.6	21.0	14.0	0.3	12.2	3.4	1.1	0.1	24.4	29.2	25.2	1,229
30-39	2.9	5.0	15.7	21.1	20.2	0.3	11.9	4.1	2.0	0.0	25.3	24.9	26.8	1,909
40-49	1.9	7.9	12.7	15.7	19.8	0.8	12.5	3.6	2.1	0.1	27.7	24.7	29.5	1,286
Marital status														
Currently married	3.1	5.5	15.1	18.6	18.1	0.5	10.8	3.2	1.7	0.1	24.5	28.7	25.8	4,925
Formerly married	1.4	6.5	22.1	13.4	16.5	0.0	12.4	4.6	1.2	0.0	27.8	26.2	28.1	241
Never married	3.8	6.8	14.8	13.4	9.6	0.4	9.8	3.0	1.1	0.0	29.4	32.0	30.1	2,060
Residence														
Urban	2.8	5.7	18.9	19.1	15.8	0.5	13.1	3.8	1.6	0.0	26.4	25.4	27.7	5,152
Rural	4.4	6.3	6.2	11.7	15.2	0.5	4.2	1.8	1.2	0.1	24.9	40.0	25.8	2,074
Region														
West	3.0	3.6	21.5	21.6	16.3	0.5	12.4	3.5	1.9	0.0	24.6	26.1	26.2	2,957
South	3.3	10.6	10.3	10.8	13.1	0.1	8.2	3.3	1.3	0.1	24.4	35.6	25.2	1,037
Central	3.8	6.0	13.5	15.0	13.7	0.5	9.6	3.2	1.1	0.0	31.3	27.5	32.1	1,799
North	1.9	7.8	10.0	20.0	22.6	0.6	13.2	3.2	1.8	0.3	20.6	28.0	21.9	598
East	4.1	6.3	6.6	9.9	15.5	0.6	7.2	2.1	0.8	0.0	25.2	40.1	25.9	835
Education														
No educ./Pri. incomp.	3.8	6.4	6.2	10.7	20.2	0.3	4.3	1.7	1.7	0.1	19.2	40.4	20.8	1,021
Pri. comp./Sec. incomp.	3.7	6.1	11.4	14.7	15.7	0.3	8.1	2.2	1.3	0.0	23.9	34.1	24.9	4,650
Sec. comp./+	1.6	4.9	32.7	27.8	12.4	1.0	21.9	7.1	1.8	0.1	36.6	8.9	37.9	1,555
Total	3.3	5.9	15.2	16.9	15.6	0.5	10.6	3.2	1.5	0.0	26.0	29.6	27.1	7,226

¹ Includes avoiding kissing and avoiding mosquito bites.

Table 11.3.2 Knowledge of ways to avoid HIV/AIDS: husbands

Percentage of all husbands knowing about AIDS by knowledge of ways to avoid HIV/AIDS, according to selected background characteristics, Turkey 1998

Background characteristic	Ways to avoid AIDS											Don't know specific ways	Percentage with misinformation ¹	Number of husbands	
	No way to avoid AIDS	Abstain from sex	Use condoms	Have only one sex partner	Avoid sex with prostitutes	Avoid sex with homosexuals	Avoid transfusions	Avoid injections	Avoid kissing	Avoid mosquito bite	Other				
Age															
15-24	3.0	0.7	30.7	11.5	30.8	0.7	7.9	1.6	1.5	0.0	21.9	12.8	22.7	102	
25-29	3.3	3.3	27.2	27.5	26.0	3.1	13.3	3.9	1.7	0.2	18.8	10.0	20.4	327	
30-39	1.7	5.1	21.4	25.6	29.4	2.2	16.8	4.6	2.6	0.0	21.0	8.9	23.3	679	
40-49	1.4	6.8	17.6	22.0	32.8	2.0	17.1	3.4	2.9	0.2	21.6	9.3	23.1	534	
50+	2.4	2.7	9.5	9.9	37.6	3.6	12.6	5.2	1.9	0.0	25.0	11.6	25.4	190	
Residence															
Urban	1.6	4.8	23.3	22.7	32.5	2.7	19.2	5.1	3.0	0.1	22.5	7.0	24.6	1,281	
Rural	3.2	4.6	14.5	21.8	26.6	1.5	6.4	1.5	1.1	0.0	18.2	16.0	18.9	551	
Region															
West	1.5	4.4	26.1	19.3	33.7	2.6	23.6	7.4	2.2	0.0	20.0	5.5	21.7	745	
South	3.4	5.3	14.9	21.8	35.2	3.3	11.8	1.7	2.6	0.3	14.8	16.6	16.9	262	
Central	1.3	4.7	19.2	21.9	27.2	1.7	9.5	1.8	3.0	0.0	25.4	8.4	26.8	462	
North	2.3	2.4	18.1	41.7	22.5	2.0	8.9	2.8	3.2	0.3	22.8	13.3	25.2	140	
East	3.8	6.9	13.5	23.0	27.6	1.9	8.1	0.9	1.0	0.3	23.1	16.1	24.3	223	
Education															
No educ./Pri.incomp.	2.7	2.8	6.6	11.4	33.6	0.0	6.4	1.1	1.4	0.0	18.0	23.6	18.0	139	
Pri. comp./Sec. incomp.	2.7	5.6	17.9	19.0	32.3	1.5	10.8	2.2	2.2	0.1	18.8	11.6	20.7	1,180	
Sec. comp./+	0.4	3.5	30.7	33.5	26.3	4.9	28.1	9.0	3.0	0.2	27.6	1.6	29.3	513	
Total	2.1	4.8	20.6	22.5	30.7	2.4	15.3	4.0	2.4	0.1	21.2	9.7	22.9	1,832	

¹ Includes avoiding kissing and avoiding mosquito bites.

Women and husbands who stated that AIDS is preventable generally were able to state more than one way to avoid it. Among women, the most common responses were that AIDS can be prevented by having sex with one partner (17 percent), by avoiding sex with prostitutes (16 percent), and by using condoms (15 percent). Similarly, among husbands, the most common responses were that AIDS can be prevented by avoiding sex with prostitutes (31 percent), by having sex with one partner (23 percent), by using condoms (21 percent) or by avoiding blood transfusions (15 percent). Sizeable percentages of respondents also mentioned other ways (26 percent for women and 21 percent for husbands). A small percentage of women and husbands cited avoiding having sex with homosexuals and avoiding injections (less than 5 percent) as ways to avoid getting AIDS.

Knowledge of ways to prevent AIDS varies by region and educational level. The percentage of women who say that AIDS can be avoided by having only one sex partner varies from 10 percent in the East region to 22 percent in the West region; for husbands, percentages for this category vary from 19 percent in the West region to 42 percent in the North region. The percentage of husbands and women who stated using condoms as a way to avoid AIDS varies from 7 percent for husbands and 6 percent for women for the lowest educational category, to 31 percent for husbands and 33 percent for women for those with the highest educational qualifications.

11.3 Perception of Risk of AIDS

More than two-thirds of women and husbands who have heard of AIDS stated that a healthy-looking person could have AIDS (Tables 11.4.1 and 11.4.2). Significant proportions of women and husbands (17 and 13 percent, respectively) did not know whether this was possible. Differentials by subgroups are again significant and similar patterns are applicable for both women and husbands. Women and husbands living in the West region, those living in urban areas, and those with higher educational qualifications are more likely to know that it is possible for a person with AIDS to look healthy. Younger women and husbands are also more knowledgeable about this fact than their older counterparts.

It appears that the general perception of the disease in Turkey is that it is almost always fatal. Sixty-five percent of women and 74 percent of husbands stated that AIDS was almost always a fatal disease. Very few women and husbands think that AIDS almost never kills. It is interesting to note that subgroup differentials are not great in this respect, meaning that the perception of AIDS as a fatal disease is universal among all subgroups of Turkish society.

Table 11.4.1 Knowledge of AIDS-related issues: women

Percent distribution of women by responses to questions on various AIDS-related issues, according to selected background characteristics, Turkey 1998

Background characteristic	Can a healthy-looking person have the AIDS virus?			Is AIDS a fatal disease?				Number of women
	Yes	No	Don't know/ Missing	Almost never	Sometimes	Almost always	Don't know/ Missing	
Age								
15-19	73.0	15.9	11.1	4.1	22.2	59.0	14.7	1,434
20-24	72.2	13.4	14.4	3.5	17.3	64.6	14.6	1,368
25-29	67.8	14.6	17.5	3.0	16.9	64.6	15.5	1,229
30-39	63.8	16.4	19.8	1.8	14.4	68.4	15.4	1,909
40-49	60.9	16.2	22.9	1.8	13.6	67.6	17.0	1,286
Marital status								
Currently married	64.4	15.7	19.9	2.4	14.8	66.0	16.9	4,925
Formerly married	65.4	15.0	19.6	3.3	18.7	62.9	15.1	241
Never married	74.7	14.7	10.6	3.7	21.5	62.9	11.9	2,060
Residence								
Urban	71.5	13.9	14.7	2.6	15.8	68.8	12.8	5,152
Rural	57.2	19.2	23.6	3.2	19.3	55.7	21.9	2,074
Region								
West	73.8	11.2	15.0	2.8	14.7	69.6	12.9	2,957
South	61.5	19.9	18.5	3.6	16.6	63.6	16.2	1,037
Central	66.5	14.2	19.3	2.3	17.7	61.6	18.4	1,799
North	65.8	17.9	16.2	1.6	22.3	57.8	18.3	598
East	54.8	25.3	19.8	3.3	18.4	63.0	15.2	835
Education								
No educ./Pri. incomp.	48.7	22.6	28.6	2.1	15.2	59.4	23.3	1,021
Pri. comp./Sec. incomp.	64.4	16.9	18.6	3.4	16.0	63.3	17.3	4,650
Sec. comp./+	88.5	6.0	5.5	1.2	20.3	73.7	4.8	1,555
Total	67.4	15.4	17.2	2.8	16.8	65.0	15.4	7,226

Table 11.4.2 Knowledge of AIDS-related issues: husbands

Percent distribution of husbands by responses to questions on various AIDS-related issues, according to selected background characteristics, Turkey 1998

Background characteristic	Can a healthy-looking person have the AIDS virus?			Is AIDS a fatal disease?			Number of husbands	
	Yes	No	Don't know/ Missing	Almost never	Sometimes	Almost always		Don't know/ Missing
Age								
15-24	75.7	19.3	5.1	1.3	15.8	71.9	11.0	102
25-29	75.7	13.2	11.2	0.6	16.2	73.4	9.8	327
30-39	69.8	18.5	11.7	0.9	16.7	74.9	7.5	679
40-49	62.2	21.9	15.9	1.3	13.8	75.2	9.7	534
50+	61.4	20.2	18.4	2.2	11.3	72.5	14.0	190
Marital status								
Currently married	68.1	18.8	13.1	1.2	15.2	74.3	9.4	1,832
Residence								
Urban	72.3	17.5	10.2	1.0	16.3	76.0	6.6	1,281
Rural	58.3	21.7	20.0	1.5	12.4	70.2	15.9	551
Region								
West	71.9	18.1	10.0	0.9	15.9	76.8	6.4	745
South	64.7	18.0	17.2	1.3	11.9	73.1	13.7	262
Central	71.6	16.5	11.9	0.7	17.8	71.1	10.4	462
North	56.2	25.6	18.2	0.6	14.5	77.6	7.2	140
East	59.6	22.3	18.1	3.4	11.3	71.7	13.6	223
Education								
No educ./Pri. incomp.	47.2	25.0	27.8	3.1	8.8	74.8	13.3	139
Pri. comp./Sec. incomp.	63.7	21.0	15.3	1.0	12.7	74.6	11.7	1,180
Sec. comp./+	83.9	12.0	4.1	1.0	22.4	73.5	3.1	513
Number of non-married sexual partners								
0	68.1	18.8	13.1	1.2	15.2	74.3	9.4	1,832
Total	68.1	18.8	13.1	1.2	15.2	74.3	9.4	1,832

REFERENCES

- Hacettepe University, Institute of Population Studies (HIPS). 1980. *Turkish Fertility Survey, 1978. First Report, Volume I: Methodology and Findings*. Ankara: HIPS.
- Hacettepe University, Institute of Population Studies (HIPS). 1987. *1983 Turkish Population and Health Survey*. Ankara: HIPS.
- Hacettepe University, Institute of Population Studies (HIPS). 1989. *1988 Turkish Population and Health Survey*. Ankara: HIPS.
- Hancıoğlu, A. and B. Akadlı Ergöçmen. 1992. Some Social Aspects of Turkish Marital Unions and their Relationship with Early Age Mortality. *Turkish Journal of Population Studies*, 14:3-26.
- Krasovec, K. and M.A. Anderson. 1991. *Maternal Nutrition and Pregnancy Outcomes: Anthropometric Assessment*, Scientific Publication No. 529. Washington D.C.: Pan American Health Organization.
- Martorell, R. and J.P. Habicht. 1986. Growth in Early Childhood in Developing Countries, in *Human Growth: A Comprehensive Treatise*. ed. F. Falkner and J.M. Tanner, 3:241-262. New York: Plenum Press.
- Ministry of Health, Hacettepe University Institute of Population Studies, and Macro International Inc 1994. *Turkish Demographic and Health Survey 1993*. Ankara, Turkey.
- Population Reference Bureau, 1999. *World Population Data Sheet. Book Edition*. Washington D.C.
- Rutstein, S.O. 1983. *Infant and Child Mortality: Levels, Trends, and Demographic Differentials*. WFS Comparative Studies No. 38. Voorburg, Netherlands: International Statistical Institute.
- Rutstein, S.O. 1984. *Infant and Child Mortality: Levels, Trends, and Demographic Differentials.. Revised Edition*. WFS Comparative Studies No. 43. Voorburg, Netherlands: International Statistical Institute.
- State Institute of Statistics (SIS). 1991. *1989 Turkish Demographic Survey*. Ankara: SIS.
- State Institute of Statistics (SIS). 1993. *1990 Census of Population: Social and Economic Characteristics of Population*. Ankara: SIS.
- State Institute of Statistics (SIS). 1997. *Statistical Yearbook of Turkey, 1997*. Ankara: SIS.
- State Institute of Statistics (SIS). 1999. Web site: [http:// www.die.gov.tr](http://www.die.gov.tr).
- Sullivan, J., G.T. Bicego and S.O. Rutstein. 1990. Assessment of the Quality of Data Used for the Direct Estimation of Infant and Child Mortality in the Demographic Surveys. In *An Assessment of DHS-I Data Quality*. DHS Methodological Reports, No.1. Columbia, Maryland: Institute for Resource Development/Macro Systems, Inc.
- United Nations Population Fund, 1995, *Programme Review and Strategy Development Report*. No.48, New York.

APPENDIX A

**PERSONNEL INVOLVED
IN THE TURKISH DEMOGRAPHIC
AND HEALTH SURVEY**

APPENDIX A

PERSONNEL INVOLVED IN THE TURKISH DEMOGRAPHIC AND HEALTH SURVEY

Project Director

Prof. Dr. Aykut Toros
Hacettepe University, Institute of Population Studies

Survey Director

Assoc. Prof. Dr. Banu Akadlı Ergöçmen
Hacettepe University, Institute of Population Studies

Field Director

Assist. Prof. Dr. İsmet Koç
Hacettepe University, Institute of Population Studies

Research Assistants

Tuba Dündar (regional coordinator)
Hacettepe University, Institute of Population Studies

Elif Kurtuluş (regional coordinator)
Hacettepe University, Institute of Population Studies

Burcu Savaş
Hacettepe University, Institute of Population Studies

Kutay Törüner
Hacettepe University, Institute of Population Studies

Steering Committee

Uğur Aytaç
General Directorate of MCH/FP, Ministry of Health

Dr. Mehmet Ali Biliker
General Directorate of MCH/FP, Ministry of Health

Prof. Dr. Çiğdem Bumin
Association for Voluntary Surgical Contraception (AVSC)

Hasibe Dedeş
State Institute of Statistics

Assoc. Prof. Dr. Banu Akadlı Ergöçmen
Hacettepe University, Institute of Population Studies

Nuray Fincancıoğlu
Human Resources Development Foundation

Prof. Dr. Ömer Gebizlioğlu
State Institute of Statistics

Assoc. Prof. Dr. Attila Hancıoğlu
Hacettepe University, Institute of Population Studies

Mehmet Konaş
State Planning Organization

Head of Data Processing

Assoc. Prof. Dr. Turgay Ünalın
Hacettepe University, Institute of Population Studies

Head of Listing

A. Sinan Türkyılmaz
Hacettepe University, Institute of Population Studies

Head of Sampling

Assoc. Prof. Dr. Attila Hancıoğlu
Hacettepe University, Institute of Population Studies

A. Sinan Türkyılmaz
Hacettepe University, Institute of Population Studies

Measure/DHS+ Staff

Dr. Edilberto Loaiza

Dr. Alfredo Aliaga

Dr. Ann Way

Han Ragers

Jeanne Cushing

Darrin Helsel

Dr. Sidney Moore

Kaye Mitchell

Dr. M. Rifat Köse

General Directorate of MCH/FP, Ministry of Health

Ahmet Özırmak
United Nations Population Fund (UNFPA)

Prof. Dr. Zafer Öztekin

Hacettepe University, Department of Public Health

Dr. Pınar Senlet

U.S. Agency for International Development (USAID)

Prof. Dr. Aykut Toros

Hacettepe University, Institute of Population Studies

Prof. Dr. Ergül Tunçbilek

Hacettepe University, Faculty of Medicine,

Department of Child Health

Prof. Dr. Sunday Üner

Hacettepe University, Institute of Population Studies

Dr. Samira Yener

State Planning Organization

APPENDIX B

SURVEY DESIGN

APPENDIX B

SURVEY DESIGN

A. Sinan Türkyılmaz and Alfredo Aliaga

The major features of **sample design** and **implementation** for the Turkish Demographic and Health Survey (TDHS) are described in this section. **Sample design** features that are discussed include: target sample size, choice of domains, sampling stages, stratification, degree of clustering, and the relationship of design decisions to the nature of the sample frame.¹ Aspects of the **sample implementation** include the cartographic and listing work that was needed to update, improve, or generate the ultimate sample lists of households or individuals, as well as the procedures for the final household selection.

This section also presents information on fieldwork, including descriptions of the recruitment and training of interviewers, the composition of interviewing teams, quality control procedures, and various practical problems encountered. Response rates² for urban and rural areas and regions are presented. An account is also given of the data processing and analysis, including a description of the calculation of the final weighting factors (design and non-response weights).

B.1 Sample Design and Implementation

A weighted, multistage, stratified cluster sampling approach was used in the selection of the TDHS-98 sample. The sample was designed in this fashion because of the need to provide estimates for a variety of characteristics for various domains. These domains, which are frequently employed in the tabulation of major indicators from the survey, are:

- Turkey as a whole;
- Urban and rural areas (each as a separate domain);
- Each of the major five regions of the country, namely the West, South, Central, North, and East regions.

The major objective of the TDHS sample design was to ensure that the survey would provide estimates with acceptable precision for these domains for most of the important demographic characteristics, such as fertility, infant and child mortality, and contraceptive prevalence, as well as for the health indicators. The different populations covered by the TDHS survey were defined as the total population for the Household Questionnaire, as well as all ever-married women younger than age 50 for the Women Questionnaire, all never-married women between ages 15 and 49 for the Single Women Questionnaire, and all husbands (of eligible currently married women) for the Husband Questionnaire. The aim was to survey these populations by designing a sample of households and interviewing an adult member of the household in order to collect information on household members. All women in the household who were identified as eligible in the

¹ For an additional description of these aspects of sample designs for DHS surveys, see the *DHS Sampling Manual*, Basic Documentation Series, No. 8, pp. 59-66.

² For a more complete discussion of the calculation of response rates, see the *DHS Sampling Manual*, Basic Documentation Series, No. 8, pp. 55-57.

household schedule were interviewed.³ In addition, husbands of eligible currently married women were interviewed in a subsample of one-half of all households.

B.2 Sample Frame

Different criteria have been used to describe "urban" and "rural" settlements in Turkey. In the demographic surveys of the 1970s a population size of 2,000 was used to differentiate between urban and rural settlements. In the 1980s, this was increased to 10,000 and, in some surveys in the 1990s, to 20,000. A number of surveys used the administrative status of settlements in combination with population size for the purpose of differentiation.

The urban frame of the 1998 TDHS consisted of a list of provincial centres, district centres, and other settlements with populations larger than 10,000, regardless of administrative status. In turn, the rural frame consists of all district centres, subdistricts and villages not included in the urban frame. Initial information on these settlements was obtained from the preliminary results of 1997 Population Count. The preliminary results of 1997 Population Count provided a computerized list of all settlements (provincial and district centres, subdistricts and villages) and their population. The population counts were taken from the cumulative enumeration forms for settlements, which were filled by supervisors during the Population Count.

B.3 Stratification

Currently Turkey is divided administratively into 80 provinces. This figure was 67 for a long time, with new provinces formed since the late 1980s. For purposes of selection in prior surveys in Turkey, these provinces have been grouped into five regions, as described in Chapter 1. This regional breakdown has been popularised as a powerful variable for understanding the demographic, social, cultural, and economic differences between different parts of the country. The five regions, West, South, Central, North, and East regions, include varying numbers of provinces.

One of the priorities of the TDHS was to produce a sample design that was methodologically and conceptually consistent with the designs of previous demographic surveys carried out by the Hacettepe Institute of Population Studies. In surveys prior to the 1993, the five-region division of the country was used for stratification. In the 1993 TDHS, a more detailed stratification taking into account subregions was employed to obtain a better dispersion of the sample. The criteria for subdividing the five major regions into subregions were the infant mortality rates of each province, estimated from the 1990 Population Census using indirect techniques.⁴ Using the infant mortality estimates as well as geographic proximity, the provinces in each region were grouped into 14 subregions at the time of the 1993 TDHS. The sub-regional division developed during the 1993 TDHS was used in the 1998 survey.

³ Although all women and husbands who were permanent residents of or were visitors to the sampled households were interviewed during the fieldwork, the tabulations were restricted to those who had slept in the household the night before the interview, i.e., the analysis was based on the de facto population.

⁴ See Hancioğlu, A. 1991. *Indirect estimation of mortality from information on the survival status of a close relative: Turkey 1970-1985*, Unpublished Doctoral Dissertation, Hacettepe Institute of Population Studies, Ankara.

The 80 provinces of Turkey were classified into 5 regions and 14 subregions as follows:

Region	Sub region	Province	Number of provinces
West	1	Edirne, İstanbul, Kırklareli, Tekirdağ	4
West	2	Balıkesir, Kocaeli, Sakarya, Çanakkale, Bursa, Yalova	6
West	3	İzmir, Denizli, Manisa, Aydın	4
South	4	Muğla, Burdur, Isparta, Antalya	4
South	5	Hatay, Adana, İçel, Gaziantep, Kilis, Osmaniye	6
Central	6	Çankırı, Çorum, Yozgat, Tokat, Amasya	5
Central	7	Bilecik, Eskişehir, Uşak, Kütahya, Afyon	5
Central	8	Ankara, Kırşehir, Nevşehir, Bolu, Konya, Kayseri, Niğde, Aksaray, Karaman, Kırıkkale	10
North	9	Trabzon, Rize, Giresun, Ordu, Artvin	5
North	10	Samsun, Kastamonu, Zonguldak, Sinop, Bartın, Karabük	6
East	11	Mardin, Diyarbakır, Siirt, Hakkari, Bitlis, Van, Batman, Şırnak	8
East	12	Kars, Bingöl, Ağrı, Muş, Erzurum, Ardahan, Iğdır	7
East	13	Şanlıurfa, Malatya, Adıyaman, K.Maraş, Sivas	5
East	14	Tunceli, Elazığ, Erzincan, Gümüşhane, Bayburt	5

B.4 Sample Allocation

The target sample size of 10,000 households was allocated among the five major divisions using the sampling error estimates from the TDHS-93 in combination with the power allocation technique⁵ with the expectation that the target sample size would provide about 8,000 completed individual interviews. During the power allocation calculations, the aim was to keep the allocation as similar as possible to the 1993 TDHS. The optimal distribution (with power 0.4) among the five major regions is shown in Table B.1. For purposes of comparison, Table B.1 also shows the allocation of the TDHS-93 sample and the allocation if the TDHS-98 sample had been distributed proportional to the size of the population in each region.

Region	Allocation in TDHS-93	TDHS-98	
		Proportional allocation	Power allocation
West	2,700	4,000	2,800
South	1,700	1,400	1,800
Central	2,100	2,300	2,100
North	1,500	800	1,500
East	2,000	1,500	2,000

To have an adequate representation of clusters within each of the five major regions, it was decided to select 25 households per standard urban segments (each consisting of 100 households) and 15 households per standard rural segment. It was also determined that 70 percent of the 10,000 households would be located in urban settlements and 30 percent in rural settlements. On this basis, the total number of selected standard segments by regions is shown in Table B.2.

⁵ See Bankier, M.D., 1988. "Power allocations: Determining sample sizes for subnational areas," *The American Statistician*, 42:(3):174-177)

Region	Urban segments (Population > 10000) (Cluster size = 25 HHs)	Rural segments (Population < 10000) (Cluster size = 15 HHs)	Number of segments
West	79	56	135
South	51	36	87
Central	59	42	101
North	41	30	71
East	50	36	86
Total	280	200	480

B.5 Sample Selection

Selection Procedures

The lists of settlements of urban settlements (settlements with 10,000 or more population) and rural settlements (settlements with less than 10,000 population) constituted the frame for the first stage of the sample selection. For the selection of the first-stage sample, settlements were grouped within each of the 14 subregions, and a systematic random sample of settlements with probability proportional to size (PPS) based on the preliminary 1997 Population Count was selected from the settlement lists. The output from this first stage of the selection was a list of all of the settlements included in the 1998 TDHS sample along with the number of clusters to be drawn from each settlement.

In Turkey, settlements are not divided into small areal units with well-defined boundaries (e.g., census enumeration areas) that can be used for conducting surveys. For some settlements, however, household lists were available from the Structure Schedules that were prepared in 1995 by many municipalities in collaboration with the State Institute of Statistics (SIS). Household lists from the Structure Schedules were available for settlements from which 340 clusters in the TDHS-98 sample were to be drawn. For those settlements, the household lists were subdivided into segments of approximately 100 households. The list of these segments constituted the frame for the selection of the 340 clusters. For each of the selected clusters, SIS provided a list of the dwellings units with their full addresses (quarter, area, avenue/street, building and door number).

SIS was not able to provide a frame from the 1995 Structure Schedules for settlements from which 140 clusters were to be drawn for the TDHS-98. For these settlements, the list of households had to be prepared in the field. In the case of small settlements (less than 250 households), the entire settlement was listed. In the case of the small number of settlements in which there were more than 250 households, 200 households were listed and an estimate of the remaining number of households in the settlement was obtained through a quick count.

Listing and Mapping Activities

Although the SIS had dwelling lists for many clusters, they did not have the corresponding maps. For this reason, the selected clusters were formed with streets that were not always adjacent to each other. Moreover, the lists provided by the SIS did not reflect changes that may have occurred during the period from the 1995 to the survey date. Two types of changes were possible: those that could be updated during listing, such as the construction of a new building on the street, a change in the use of a building (e.g., a flat can be used as an office instead of a dwelling), or changes in the names of streets, and those that were more problematic, e.g., the appearance of new quarters in urban centres.

In an effort to develop strategies for dealing with these as well as other possible problems that might arise, a pilot listing activity was undertaken in the capital, Ankara, before the actual listing activity began. The final listing forms, sketch map formats, and listing and mapping manuals were developed based on this experience.

Forty (40) university students were trained for the main listing activity. Listing teams were formed following a four-day training program in the beginning of June 1998. Each team was provided with maps describing the location of the settlements they were expected to visit as well as other materials needed for the listing. Sixteen (16) listing teams were constituted with one mapper and one lister. The listing operation started on 8th of June. It was carried under the supervision of the research assistants and regional coordinators from the Hacettepe Institute of Population Studies.

The cluster (standard segment) size was around 100 households for most of the clusters in urban areas. Only two urban clusters had extremely low numbers of households; in order to obtain 100 households in these clusters, adjacent streets were added to the original cluster. In some of the selected villages, the total populations also were small, and, therefore, the original cluster did not include 100 households. In these cases, the village that was nearest to the selected village was included in the sample, and the names of these villages were provided to the listing teams; the lists of 100 households were completed from the two villages.

Most of the listing activity was completed before the training for the main fieldwork began in July. Overall, the quality of the listing work produced by the listers was good although it varied somewhat largely in response to problems the listing teams experienced in working in some geographic areas. In particular, there were some problems with the listing of clusters of Adana province where there had been an earthquake. There also were problems with the lists for İçel province. Finally, three clusters were not listed due to problems of accessibility; information on these clusters is presented later in this Appendix.

B.6 Questionnaire Development and Pre-test

Questionnaires

Four main types of questionnaires were used to collect the TDHS data: the Household Questionnaire and the Individual Questionnaires for ever-married women of reproductive ages, for never-married women and for husbands. The contents of these questionnaires were based on the DHS Model "A" Questionnaire, which was designed for the DHS program for use in countries with high contraceptive prevalence. Additions, deletions and modifications were made to the DHS model questionnaire in order to collect information particularly relevant to Turkey. Attention also was paid to ensuring the comparability of the TDHS findings with previous demographic surveys carried out by the Hacettepe Institute of Population Studies. In the process of designing the TDHS questionnaires, national and international population and health agencies were consulted for their comments.

All TDHS questionnaires were developed in English and then translated into Turkish. English versions of the Household and Individual questionnaires are reproduced in Appendix E.

The Household Questionnaire was used to enumerate all usual members of and visitors to the selected households and to collect information relating to the socioeconomic position of the households. In the first part of the Household Questionnaire, basic information was collected on the age, sex, educational attainment, marital status, and relationship to the head of household of each person listed as a household member or visitor. The objective of the first part of the Household Questionnaire was to obtain the information needed to identify women and husbands who were eligible for the individual interview as well as to provide basic demographic data for Turkish households. The second part of the Household Questionnaire included questions on the welfare of the elderly people. In the third part of the Household Questionnaire, questions were included on the

dwelling unit, such as the number of rooms, the flooring material, the source of water, and the type of toilet facilities, and on the household's ownership of a variety of consumer goods.

The Individual Questionnaire for women obtained information on the following subjects:

- Background characteristics
- Reproduction
- Marriage
- Knowledge and use of family planning
- Maternal care and breastfeeding
- Immunisation and health
- Fertility preferences
- Husband's background
- Women's work and status
- Sexually transmitted diseases and AIDS
- Maternal and child anthropometry.

The Individual Questionnaire for ever-married women included a monthly calendar, which was used to record fertility, contraception, marriage and migration histories for a period of approximately six years beginning in January 1993 up to the survey month. In addition, fieldwork teams measured the heights and weights of children under age five and of all women at ages 15-49.

The Individual Questionnaire for never-married women covered the following subjects:

- Background characteristics
- Reproduction
- Marriage
- Knowledge and use of family planning
- Fertility preferences
- Migration
- Women's work and status
- Sexually transmitted diseases and AIDS
- Anthropometry.

The Individual Questionnaire for husbands covered the following topics:

- Background characteristics
- Reproduction
- Knowledge and use of family planning
- Marriage
- Fertility preferences
- Sexually transmitted diseases and AIDS
- Attitudes

As mentioned earlier, the DHS Model "A" Questionnaire for women and men served as the base for the Turkish questionnaires. The DHS model questionnaires were modified to include subjects of particular interest in Turkey. The following is a list of some of the main differences between the standard DHS questionnaire and the TDHS questionnaire.

Information on the mother tongues and second languages known by the respondent, her husband, and their parents was collected in the TDHS.

- Additional questions were asked to respondents regarding their cumulative numbers of abortions, miscarriages and stillbirths; specific questions regarding the last abortion were also included.
- A separate section on nuptiality was included in the Individual Questionnaire of the TDHS; this included a number of questions already in the standard DHS questionnaire, as well as questions on the type of marriage, arrangement of marriage, and consanguinity, etc.
- In the husbands questionnaire withdrawal users were asked two additional questions to determine whether they were using this method in combination with other methods.
- Respondents were asked a series of additional questions concerning their attitudes and beliefs regarding the pill, the IUD, the condom and withdrawal. The questions probed whether women thought these methods were reliable, easy to use, or harmful to their health and whether their husbands opposed their use.
- A series of reproductive health questions were included.
- Questions on the foods children were given for the last seven days on cough/acute respiratory tract infection and fever were not included.
- Questions to ascertain the general level of knowledge on diarrhoea were not included
- A number of questions regarding recent sexual activity as well as initiation of sexual activity were not included.
- A separate section dealing with the attitudes, beliefs and behaviour of women regarding intramarital relationships, child-rearing, and status of women was included in the TDHS Individual Questionnaire.

Pre-test

In May 1998, a pre-test was conducted to ensure that the questions in the TDHS questionnaires were in a logical sequence; that the wording of the questions was comprehensible, appropriate and meaningful; and that the precoded answers were adequate.

Nineteen interviewers were trained at the Hacettepe Institute of Population Studies for a period of two weeks. The training period included both classroom training and interviews in the field. The interviewers were mostly university students and graduates. In addition to the interviewers, research assistants, who would later become regional coordinators and supervisors, also received training.

Fieldwork for the pretest was carried out in one district in central Ankara, two districts in squatter housing areas of Ankara, and three villages in Ankara province. A total of 185 household, 172 ever-married women, 72 husband and 45 never-married women interviews were completed during the pretest. Frequency distributions and cross tabulations were obtained shortly after the completion of the interviews. Based on the evaluation of these results and on the feedback obtained from the interviewers, several minor changes were made to the TDHS questionnaires.

B.7 Data Collection Activities

Staff Recruitment and Training

Candidates for the positions of interviewers, field editors, supervisors and measurers were solicited in announcements sent to the all universities in Ankara and from the Institute of Population Studies files of field staff who had worked on previous surveys. All candidates for the field staff positions were interviewed in four groups by the staff of the Institute of Population Studies using interview guidelines prepared for this purpose. Individuals who met a number of the requirements and had the necessary qualifications were accepted into the training program.

All candidates for the field staff positions were at least high school graduates and the majority were university students. Previous survey experience was not among the qualifications for the candidates for the position of interviewers in order to ensure that the trainees had no biases that might result from their previous experience. Approximately 120 applicants were accepted for the training program.

Training of the candidates for the fieldwork positions was conducted in July 1998 for three weeks at the Hacettepe Institute of Population Studies. The training program included general lectures related to the demographic situation in Turkey, family planning and mother and child health, questionnaire training, role playing and mock interviews, field practice in areas not covered in the survey and quizzes to test the progress and capabilities of the candidates. A variety of materials were used during the training sessions, including manuals for supervisors and editors, and for interviewers.

All trainees received the same classroom training during the first two weeks of the training period; at the end of the third week, supervisors, field editors, and measurers were selected from among the candidates, and a number of unsuccessful candidates were eliminated at this stage. Separate classroom training sessions were organised for supervisors, field editors, and measurers.

Fieldwork

Fieldwork for the TDHS, including initial interviews, callbacks and reinterviews began in the first week of August 1998 and was completed at the end of November 1998.

Fieldwork activities were completed in two stages. In the first stage, data collection was carried out by 12 teams, each consisting of a supervisor, a field editor, and 4 or 5 female and one male interviewers, depending on the workload of that specific team. The male interviewer and field editor worked as measurers as well. The first stage of the fieldwork was completed by the end of September, at which point a number of fieldwork staff, as agreed initially, discontinued working in the field. Four new teams were set up from among the staff of the 12 teams that had worked in the first stage of fieldwork. The teams at this second stage had the same composition as those in the first stage. These teams continued with data collection activities until the end of November.

Four regional coordinators were responsible for visiting the fieldwork teams in turn, checking the quality of data collected, and reporting periodically to the field director in Ankara.

Fieldwork teams visited 76 of the 80 provinces in Turkey. The TDHS fieldwork was a relatively rapid operation because of the specific conditions prevailing in the country, i.e., a large proportion of the fieldwork staff consisted of students who had to begin school in October and climatic conditions in many parts of the country limited access after October.

A total of 480 clusters were selected for the TDHS sample. Of these, interviews were successfully completed in 476 clusters. Due to problems of access and lack of security, three clusters were not listed and, consequently, were not visited by the fieldwork teams; in addition, a cluster that had been listed could not be visited by the fieldwork teams.

B.8 Data Processing and Analysis

Office Editing

The questionnaires were returned to the Institute of Population Studies by the fieldwork teams for data processing as soon as interviews were completed in a province. The office editing staff checked that the questionnaires for all the selected households and eligible respondents were returned from the field. The comparatively few questions that had not been precoded (e.g., occupation) were coded at this time.

Machine Entry and Editing

The data were entered and edited on microcomputers using the Integrated System for Survey Analysis (ISSA), a package program specifically developed to process DHS data. ISSA allows range, skip, and consistency errors to be detected and corrected at the data entry stage. The machine entry and editing activities were initiated within one week after the beginning of the fieldwork and were completed a few days after the completion of the fieldwork.

Advantage was taken of the fact that data processing activities ran concurrently with fieldwork. Field check tables from edited data were periodically produced for each interviewing team. These focused on such potential problems as high proportions of incomplete households and displacement of eligible respondents and were used to check the progress and quality of data from the field.

B.9 Calculation of Sample Weights

As mentioned earlier, the TDHS sampling plan is not a self-weighted one. In particular, a disproportionate number of sample units were chosen from the North, East and South regions, since there would have been inadequate numbers of observations for these areas if the target number of households had been proportionally allocated across regions. The following describes the procedure for calculating the weights to be used in the analysis of the TDHS-98 results. Since the final selection was not implemented proportionally in urban and rural areas within each subregion, and since there was some variation in urban and rural nonresponse rates, separate weights are calculated for rural and urban areas within each of the 14 subregions.

The major component of the weight is the reciprocal of the sampling fraction employed in calculating the number of units in that particular stratum (subregion):

$$W_h = 1 / f_h .$$

The term $f(h)$, the sampling fraction at the h^{th} stratum, is the product of the probabilities of selection at every stage in a stratum:

$$f_h = P_{1h} * P_{2h} * P_{3h}$$

where P_{ih} is the probability of the sample unit in the i -th sample stage for the h -th strata.

A second component taken into account in the calculation of the weights is the level of nonresponse for the household and the individual interviews. The adjustment for household nonresponse is equal to the inverse value of:

$$R_{hh} = \text{Completed households/Eligible households.}$$

Eligible households include households where interviews were completed, households where there were no competent respondents, households where interviews were postponed and eventually not completed, refusals, and those dwellings that were not found by the fieldwork teams.

Similarly, the adjustment for nonresponse in the women's survey is equal to the inverse value of:

$$R_{ww} = \text{Completed women questionnaires/Eligible women.}$$

Approximately half of the households were selected for the husband questionnaires. The rule for the selection of a household for husband interview was very simple. If the cluster was even-numbered, then the households whose number was even were selected for husband interview or vice versa. A separate set of sampling weights were calculated for the husband sample following procedures similar to those described above. For the husband survey, the adjustment for nonresponse is defined as:

$$R_{mm} = \text{Completed husband questionnaires/Eligible husbands.}$$

The weights for the subregions regions and the factors compensating for nonresponse are shown in Table B.3.1 for women and Table B.3.2 for husbands.

The weights for the TDHS-98 also include an adjustment for the missing clusters (one cluster in subregion 7 and three clusters in subregion 11).

The unadjusted weights for the households were calculated by multiplying the above factors for each subregion; they were then standardised by multiplying these weights by the ratio of the number of completed interviewed households to the total unadjusted weighted number of households. A similar standardisation procedure was followed in obtaining the weights for the individual women's and the husbands' data. The final weights for households and individual women and husband are shown in Table B.4.

Table B.3.1 Design weights and nonresponse factors: women sample

Design weights and nonresponse factors by subregion and urban-rural residence for the women and husbands samples, Turkey 1998

Region	Subregion	Residence	Inverse of overall sampling fraction	Household level	Women level
West	Subregion 01	Urban	2452082 / 950	875 / 792	893 / 761
		Rural	194415 / 405	374 / 366	389 / 358
West	Subregion 02	Urban	951430 / 475	400 / 384	370 / 345
		Rural	422277 / 210	189 / 186	181 / 166
West	Subregion 03	Urban	1110291 / 550	478 / 445	437 / 395
		Rural	537244 / 225	206 / 198	164 / 145
South	Subregion 04	Urban	363925 / 400	321 / 292	329 / 296
		Rural	332057 / 165	145 / 143	156 / 140
South	Subregion 05	Urban	909842 / 875	782 / 748	909 / 851
		Rural	361621 / 375	343 / 329	417 / 391
Central	Subregion 06	Urban	245110 / 275	251 / 235	247 / 233
		Rural	270512 / 105	98 / 96	131 / 119
Central	Subregion 07	Urban	360313 / 275	215 / 205	203 / 175
		Rural	249081 / 120	108 / 105	123 / 104
Central	Subregion 08	Urban	1509029 / 900	799 / 693	769 / 680
		Rural	574728 / 405	372 / 356	428 / 395
North	Subregion 09	Urban	254793 / 500	373 / 363	425 / 385
		Rural	268104 / 225	190 / 184	238 / 217
North	Subregion 10	Urban	288549 / 525	430 / 376	453 / 422
		Rural	279613 / 225	193 / 183	252 / 234
East	Subregion 11	Urban	346485 / 425	359 / 341	499 / 434
		Rural	210022 / 135	121 / 119	217 / 202
East	Subregion 12	Urban	181206 / 250	209 / 191	277 / 258
		Rural	212196 / 120	113 / 112	174 / 154
East	Subregion 13	Urban	488554 / 450	354 / 330	406 / 375
		Rural	304478 / 195	170 / 166	224 / 200
East	Subregion 14	Urban	115357 / 125	92 / 85	105 / 92
		Rural	95092 / 45	36 / 36	52 / 49

Table B.3.2 Design weights and nonresponse factors: husbands sample

Design weights and nonresponse factors by subregion and urban-rural residence for the women and husbands samples, Turkey 1998

Region	Subregion	Residence	Inverse of overall sampling fraction	Household level	Husband level
West	Subregion 01	Urban	2 * 2452082 / 950	440 / 394	288 / 158
		Rural	2 * 194415 / 405	191 / 187	133 / 107
West	Subregion 02	Urban	2 * 951430 / 475	214 / 206	137 / 94
		Rural	2 * 422277 / 210	97 / 95	59 / 47
West	Subregion 03	Urban	2 * 1110291 / 550	240 / 228	161 / 105
		Rural	2 * 537244 / 225	100 / 99	53 / 35
South	Subregion 04	Urban	2 * 363925 / 400	159 / 144	110 / 71
		Rural	2 * 332057 / 165	74 / 72	53 / 37
South	Subregion 05	Urban	2 * 909842 / 875	397 / 379	299 / 210
		Rural	2 * 361621 / 375	169 / 165	113 / 82
Central	Subregion 06	Urban	2 * 245110 / 275	123 / 116	89 / 66
		Rural	2 * 270512 / 105	50 / 50	53 / 36
Central	Subregion 07	Urban	2 * 360313 / 275	103 / 99	72 / 39
		Rural	2 * 249081 / 120	58 / 58	50 / 26
Central	Subregion 08	Urban	2 * 1509029 / 900	397 / 348	263 / 163
		Rural	2 * 574728 / 405	190 / 182	125 / 83
North	Subregion 09	Urban	2 * 254793 / 500	195 / 190	133 / 95
		Rural	2 * 268104 / 225	95 / 91	70 / 44
North	Subregion 10	Urban	2 * 288549 / 525	215 / 188	147 / 102
		Rural	2 * 279613 / 225	94 / 87	73 / 58
East	Subregion 11	Urban	2 * 346485 / 425	172 / 163	147 / 62
		Rural	2 * 210022 / 135	60 / 59	53 / 33
East	Subregion 12	Urban	2 * 181206 / 250	105 / 96	88 / 49
		Rural	2 * 212196 / 120	57 / 57	52 / 29
East	Subregion 13	Urban	2 * 488554 / 450	181 / 170	123 / 77
		Rural	2 * 304478 / 195	80 / 78	56 / 34
East	Subregion 14	Urban	2 * 115357 / 125	47 / 44	30 / 21
		Rural	2 * 95092 / 45	18 / 18	14 / 8

Table B.4 Final sample weights

Final weights by subregion and urban-rural residence, Turkey 1998

Region	Subregion	Residence	Household weight in the entire sample	Women's weight in the entire sample	Household's weight in the subsample	Husband's weights in the subsample
West	Subregion 01	Urban	1.885393	2.048465	1.909400	2.276100
		Rural	0.324320	0.326288	0.324786	0.264012
West	Subregion 02	Urban	1.379494	1.369817	1.378349	1.313744
		Rural	1.350938	1.363847	1.360054	1.116527
West	Subregion 03	Urban	1.433671	1.468568	1.407601	1.411481
		Rural	1.642477	1.720027	1.597657	1.582160
South	Subregion 04	Urban	0.661275	0.680529	0.665451	0.674232
		Rural	1.349177	1.391956	1.370117	1.283485
South	Subregion 05	Urban	0.718738	0.710829	0.721503	0.671814
		Rural	0.664704	0.656369	0.654266	0.589628
Central	Subregion 06	Urban	0.629423	0.617794	0.626045	0.552091
		Rural	1.738844	1.772332	1.706582	1.643083
Central	Subregion 07	Urban	0.908531	0.975793	0.902981	1.090198
		Rural	1.411572	1.545734	1.374960	1.729201
Central	Subregion 08	Urban	1.278134	1.338301	1.267055	1.336973
		Rural	0.980411	0.983592	0.981338	0.966518
North	Subregion 09	Urban	0.346201	0.353848	0.346440	0.317187
		Rural	0.813514	0.826118	0.824011	0.857309
North	Subregion 10	Urban	0.415574	0.413042	0.416361	0.392416
		Rural	0.866543	0.864041	0.889434	0.732095
East	Subregion 11	Urban	0.567472	0.604109	0.569858	0.883591
		Rural	1.045872	1.040272	1.047999	1.100733
East	Subregion 12	Urban	0.524389	0.521283	0.525146	0.616774
		Rural	1.179574	1.233996	1.171349	1.373569
East	Subregion 13	Urban	0.770011	0.771883	0.765700	0.799893
		Rural	1.057232	1.096348	1.060831	1.142653
East	Subregion 14	Urban	0.660409	0.697870	0.652996	0.610059
		Rural	1.397141	1.372800	1.399786	1.601986

B.10 Coverage of the Sample

The results of sample implementation for the household and the individual interviews for the country as a whole, for urban and rural areas, and for the five regions of Turkey are shown in Tables B.5 and Table B.6. The results indicate that, of the 9,970 households selected, the TDHS fieldwork teams successfully completed interviews with 8,059 (81 percent). The main reasons that eligible households were not interviewed were that some of the listed dwelling units were found to be vacant at the time of the interview or the household was away for an extended period. A total of 8,956 households were located and visited, of which 8,059 households were successfully interviewed. Overall, the household response rate was calculated as 93.8 percent. The household response rate was higher in rural areas than in urban areas and highest in the South and East regions.

In the interviewed households, 9,468 eligible women were identified, of whom 91 percent were interviewed. Eligibility for the individual interview required that the woman was in the reproductive ages (15-49) and was present in the household on the night before the interview. Among the small number of eligible women not interviewed in the survey, the principal reason for nonresponse was the failure to find the woman

at home after repeated visits to the household. The eligible woman response rate was higher in rural areas than in urban areas and was higher in the South and North regions than in the other three regions.

In half of the households (4,983 households), which were selected as a subsample, husband questionnaires were administered. In these households, 3,043 husbands were identified as eligible, i.e., they were currently married to women in reproductive ages listed in the household schedule. Interviews were completed with 1,971 husbands (65 percent). The highest response rate is in North region for husband questionnaires.

The overall response rate for women in the TDHS was calculated as 85 percent. It ranged from 82 percent in the Central region to 88 percent in the South region, when it was calculated for women. The overall response rate for husbands in the TDHS was 61 percent.

Table B.5 Sample implementation: women

Percent distribution of households and eligible women in the 1998 TDHS sample by results of the individual and household interviews, and response rates, according to region and urban-rural residence, Turkey 1998

Result	Region					Residence		Total
	West	South	Central	North	East	Urban	Rural	
Selected households								
Completed (C)	83.9	83.3	80.3	74.8	79.1	78.4	86.5	80.8
Household present but no competent respondent at home (HP)	3.1	2.5	4.3	2.7	0.9	3.3	1.7	2.8
Postponed	0.5	0.2	0.3	0.1	0.1	0.3	0.2	0.3
Refused (HR)	1.4	1.5	2.5	1.2	1.1	2.0	0.6	1.6
Dwelling not found (DNF)	0.3	0.2	0.2	1.4	2.2	1.0	0.2	0.8
Household absent (HA)	6.0	5.6	7.2	9.6	8.5	7.8	5.6	7.2
Dwelling vacant/ address not a dwelling (DV)	4.8	6.7	5.0	9.9	7.0	6.9	5.1	6.3
Dwelling destroyed (DD)	0.0	0.1	0.1	0.1	0.4	0.1	0.1	0.1
Other	0.0	0.0	0.0	0.1	0.7	0.2	0.0	0.2
Total percent	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Number	2,827	1,815	2,104	1,479	1,745	6,989	2,981	9,970
Household response rate (HRR)¹	94.0	95.0	91.7	93.3	94.9	92.3	97.0	93.8
Eligible women								
Completed (EWC)	89.2	92.7	89.7	92.0	90.3	90.2	91.4	90.6
Not at home (EWNH)	6.9	4.5	5.7	5.0	5.6	5.8	5.4	5.7
Postponed (EWP)	0.5	0.1	0.6	0.0	0.1	0.4	0.1	0.3
Refused (EWR)	1.8	1.1	1.4	0.5	1.5	1.6	0.8	1.4
Partly completed (EWPC)	0.7	0.3	1.3	1.0	1.6	0.9	1.1	1.0
Other (EWO)	0.9	1.3	1.2	1.5	0.9	1.1	1.2	1.1
Total percent	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Number	2,434	1,811	1,901	1,368	1,954	6,322	3,146	9,468
Eligible woman response rate (EWRR)²	89.2	92.7	89.7	92.0	90.3	90.2	91.4	90.6
Overall response rate (ORR)³	83.8	88.1	82.3	85.8	85.7	83.2	88.6	84.9

Note: The household response rate is calculated for completed households as a proportion of completed, no competent respondent, postponed, refused, and dwelling not found. The eligible woman response rate is calculated for completed interviews as a proportion of completed, not at home, postponed, refused, partially completed and "other." The overall response rate is the product of the household and woman response rates.

¹ Using the number of households falling into specific response categories, the household response rate (HRR) is calculated as:

$$\frac{C}{C + HP + P + R + DNF}$$

² Using the number of eligible women falling into specific response categories, the eligible woman response rate (EWRR) is calculated as:

$$\frac{EWC}{EWC + EWNH + EWP + EWR + EWPC + EWO}$$

³ The overall response rate (ORR) is calculated as:

$$ORR = HRR * EWRR$$

Table B.6 Sample implementation: husbands

Percent distribution of households and eligible husbands in the 1998 TDHS sample by results of the individual and household interviews, and response rates, according to region and urban-rural residence, Turkey 1998

Result	Region					Residence		Total
	West	South	Central	North	East	Urban	Rural	
Selected households								
Completed (C)	85.5	84.0	81.0	75.2	78.6	79.1	87.3	81.5
Household present but no competent respondent at home (HP)	3.4	2.2	3.5	3.4	1.0	3.3	1.6	2.8
Postponed	0.4	0.3	0.4	0.1	0.1	0.4	0.1	0.3
Refused (HR)	1.1	1.5	2.5	1.4	0.7	1.8	0.5	1.4
Dwelling not found (DNF)	0.3	0.2	0.1	0.9	2.2	0.9	0.1	0.7
Absent (HA)	5.9	5.6	8.1	8.9	9.5	8.0	6.0	7.4
Dwelling vacant/not a dwelling (DV)	3.5	6.1	4.3	9.7	6.7	6.2	4.1	5.6
Dwelling destroyed (DD)	0.0	0.0	0.2	0.1	0.3	0.1	0.2	0.1
Other (O)	0.0	0.0	0.0	0.1	0.9	0.2	0.1	0.2
Total percent	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Number	1,414	905	1,053	739	872	3,496	1,487	4,983
Household response rate (HRR)¹	94.3	95.1	92.6	92.8	95.1	92.5	97.4	94.0
Eligible husbands								
Completed (EHC)	65.7	69.6	63.3	70.7	55.7	62.9	68.9	64.8
Not at home (EHNH)	27.7	27.5	27.3	26.2	41.5	30.9	27.7	29.9
Postponed (EHP)	0.8	0.0	0.6	0.0	0.4	0.6	0.1	0.4
Refused (EHR)	3.7	2.1	6.0	1.4	1.4	3.8	1.7	3.2
Partly completed (EHPC)	0.4	0.2	0.9	0.5	0.2	0.5	0.2	0.4
Other (EHO)	1.7	0.7	1.8	1.2	0.9	1.3	1.4	1.3
Total percent	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Number	831	575	652	423	562	2,087	956	3,043
Eligible husband response rate (EHRR)²	65.7	69.6	63.3	70.7	55.7	62.9	68.9	64.8
Overall response rate (ORR)³	62.0	66.2	58.7	65.6	53.0	58.2	67.1	60.9

Note: The household response rate is calculated for completed households as a proportion of completed, no competent respondent, postponed, refused, and dwelling not found. The eligible husband response rate is calculated for completed interviews as a proportion of completed, not at home, postponed, refused, partially completed and "other." The overall response rate is the product of the household and woman response rates.

¹ Using the number of households falling into specific response categories, the household response rate (HRR) is calculated as:

$$\frac{C}{C + HP + P + R + DNF}$$

² Using the number of eligible husbands falling into specific response categories, the eligible husband response rate (EHRR) is calculated as:

$$\frac{EHC}{EHC + EHNH + EHP + EHR + EHPC + EHO}$$

³ The overall response rate (ORR) is calculated as:

$$ORR = HRR * EHRR$$

APPENDIX C

ESTIMATES OF SAMPLING ERRORS

APPENDIX C

ESTIMATES OF SAMPLING ERRORS

Alfredo Aliaga and A. SinanTurkyılmaz

The estimates from a sample survey are affected by two types of errors: (1) nonsampling errors, and (2) sampling errors. Nonsampling errors are the results of mistakes made in implementing data collection and data processing, such as failure to locate and interview the correct household, misunderstanding of the questions on the part of either the interviewer or the respondent, and data entry errors. Although numerous efforts were made during the implementation of the TDHS to minimize this type of error, nonsampling errors are impossible to avoid and difficult to evaluate statistically.

Sampling errors, on the other hand, can be evaluated statistically. The sample of respondents selected in the TDHS is only one of many samples that could have been selected from the same population, using the same design and expected size. Each of these samples would yield results that differ somewhat from the results of the actual sample selected. Sampling errors are a measure of the variability between all possible samples. Although the degree of variability is not known exactly, it can be estimated from the survey results.

A sampling error is usually measured in terms of the *standard error* for a particular statistic (mean, percentage, etc.), which is the square root of the variance. The standard error can be used to calculate confidence intervals within which the true value for the population can reasonably be assumed to fall. For example, for any given statistic calculated from a sample survey, the value of that statistic will fall within a range of plus or minus two times the standard error of that statistic in 95 percent of all possible samples of identical size and design.

If the sample of respondents had been selected as a simple random sample, it would have been possible to use straightforward formulas for calculating sampling errors. However, the TDHS sample is the result of a three-stage stratified design, and, consequently, it was necessary to use more complex formulae. The computer software used to calculate sampling errors for the TDHS is the ISSA Sampling Error Module (SAMPERR). This module used the Taylor linearization method of variance estimation for survey estimates that are means or proportions. The Jackknife repeated replication method is used for variance estimation of more complex statistics such as fertility and mortality rates.

The Taylor linearization method treats any percentage or average as a ratio estimate, $r = y/x$, where y represents the total sample value for variable y , and x represents the total number of cases in the group or subgroup under consideration. The variance of r is computed using the formula given below, with the standard

$$\text{var}(r) = \frac{1-f}{x^2} \sum_{h=1}^H \left[\frac{m_h}{m_h-1} \left(\sum_{i=1}^{m_h} z_{hi}^2 - \frac{z_h^2}{m_h} \right) \right]$$

error being the square root of the variance in which:

$$z_{hi} = y_{hi} - r \cdot x_{hi}, \text{ and } z_h = y_h - r \cdot x_h$$

where h represents the stratum which varies from 1 to H,
 m_h is the total number of enumeration areas selected in the h^{th} stratum,
 y_{hi} is the sum of the values of variable y in EA i in the h^{th} stratum,
 x_{hi} is the sum of the number of cases in EA i in the h^{th} stratum, and
 f is the overall sampling fraction, which is so small that it is ignored.

The Jackknife repeated replication method derives estimates of complex rates from each of several replications of the parent sample, and calculates standard errors for these estimates using simple formulae. Each replication considers *all but one* clusters in the calculation of the estimates. Pseudo-independent replications are thus created. In the TDHS, there were 476 non-empty clusters. Hence, 476 replications were created. The variance of a rate r is calculated as follows:

$$\text{var}(r) = \frac{1}{k(k-1)} \sum_{i=1}^k (r_i - r)^2$$

in which

$$r_i = kr - (k-1)r_{(i)}$$

where r is the estimate computed from the full sample of 476 clusters,
 $r_{(i)}$ is the estimate computed from the reduced sample of 476 clusters (i^{th} cluster excluded), and
 k is the total number of clusters.

In addition to the standard error, SAMPERR computes the design effect (DEFT) for each estimate, which is defined as the ratio between the standard error using the given sample design and the standard error that would result if a simple random sample had been used. A DEFT value of 1.0 indicates that the sample design is as efficient as a simple random sample, while a value greater than 1.0 indicates the increase in the sampling error due to the use of a more complex and less statistically efficient design. SAMPERR also computes the relative error and confidence limits for the estimates.

Sampling errors for the TDHS are calculated for a number of variables considered to be of primary interest. Results for women and for husbands are presented in this appendix for the country as a whole, for urban and rural areas, and for each of the five regions: West, South, Central, North and East. For each variable, the type of statistic (mean, proportion, or rate) and the base population are given in Table C.1. Tables C.2-C.9 present the value of the statistic (R), its standard error (SE), the number of unweighted (N) and weighted (WN) cases, the design effect (DEFT), the relative standard error (SE/R), and the 95 percent confidence limits ($R \pm 2SE$), for each variable. The DEFT is considered undefined when the standard error considering simple random sample is zero (when the estimate is close to 0 or 1).

In general, the relative standard errors for most estimates for the country as a whole are small, except for estimates of very small proportions. There are some differentials in the relative standard errors for the estimates for sub-populations. For example, for the contraceptive prevalence rate (CPR), i.e. the proportion of currently married women aged 15-49 who were using any method of contraception at the time of the interview, the relative standard error for the country as a whole, for urban areas, and for rural areas are 1.2 percent, 1.4 percent, and 2.7 percent, respectively.

To obtain the 95 percent confidence limits for the CPR, one adds and subtracts twice the standard error to the sample estimate, i.e. $0.639 \pm 2 \times 0.007$. The results indicate that there is a high probability (95 percent) that the true value of the CPR for the country as a whole lies between 62.3 percent and 65.4 percent.

Table C.1 List of selected variables for sampling errors, Turkey 1998

Variable	Estimate	Base population
WOMEN		
Urban	Proportion	All women 15-49
No education	Proportion	All women 15-49
With secondary education or higher	Proportion	All women 15-49
Never married	Proportion	All women 15-49
Currently married	Proportion	All women 15-49
Married before age 20	Proportion	All women 15-49
Sex before age 18	Proportion	All women 15-49
Children ever born	Mean	All women 15-49
Children ever born to women over 40	Mean	All women 40-49
Children surviving	Mean	All women 15-49
Know any contraceptive method	Proportion	Currently married women 15-49
Know any modern contraceptive method	Proportion	Currently married women 15-49
Ever used any contraceptive method	Proportion	Currently married women 15-49
Currently using any method	Proportion	Currently married women 15-49
Currently using a modern method	Proportion	Currently married women 15-49
Currently using pill	Proportion	Currently married women 15-49
Currently using IUD	Proportion	Currently married women 15-49
Currently using injections	Proportion	Currently married women 15-49
Currently using Norplant	Proportion	Currently married women 15-49
Currently using condom	Proportion	Currently married women 15-49
Currently using female sterilisation	Proportion	Currently married women 15-49
Currently using calendar/rhythm method	Proportion	Currently married women 15-49
Currently using withdrawal	Proportion	Currently married women 15-49
Using public sector source	Proportion	Current users of modern method
Want no more children	Proportion	Currently married women 15-49
Want to delay at least 2 years	Proportion	Currently married women 15-49
Ideal number of children	Mean	All women 15-49
Mothers received tetanus injection	Proportion	Births in last 5 years
Mothers received medical care at birth	Proportion	Births in last 5 years
Had diarrhoea in the last 2 weeks	Proportion	Children under 5
Treated with ORS packets	Proportion	Children under 5 with diarrhoea in last 2 weeks
Sought medical treatment	Proportion	Children under 5 with diarrhoea in last 2 weeks
Having health card, seen	Proportion	Children 12-23 months
Received BCG vaccination	Proportion	Children 12-23 months
Received DPT vaccination (3 doses)	Proportion	Children 12-23 months
Received polio vaccination (3 doses)	Proportion	Children 12-23 months
Received measles vaccination	Proportion	Children 12-23 months
Fully immunised	Proportion	Children 12-23 months
Weight-for-height (below -2 SD)	Proportion	Children 0-47 months
Height-for-age (below -2 SD)	Proportion	Children 0-47 months
Weight-for-age (below -2 SD)	Proportion	Children 0-47 months
Total fertility rate (3 years)	Rate	Women-years of exposure to child-bearing
Neonatal mortality rate (0-4 years)	Rate	Number of births exposed to death
Infant mortality rate (0-4 years)	Rate	Number of births exposed to death
Child mortality rate (0-4 years)	Rate	Number of births exposed to death
Under-five mortality rate (0-4 years)	Rate	Number of births exposed to death
Postneonatal mortality rate (0-4 years)	Rate	Number of births exposed to death
HUSBANDS		
Urban	Proportion	All husbands
No education	Proportion	All husbands
With secondary education or higher	Proportion	All husbands
Know any contraceptive method	Proportion	All husbands
Know any modern contraceptive method	Proportion	All husbands
Ever used any contraceptive method	Proportion	All husbands
Currently using any method	Proportion	All husbands
Currently using a modern method	Proportion	All husbands
Currently using pill	Proportion	All husbands
Currently using IUD	Proportion	All husbands
Currently using injections	Proportion	All husbands
Currently using condom	Proportion	All husbands
Currently using female sterilisation	Proportion	All husbands
Currently using calendar/rhythm method	Proportion	All husbands
Currently using withdrawal	Proportion	All husbands
Want no more children	Proportion	All husbands
Want to delay at least 2 years	Proportion	All husbands
Ideal number of children	Mean	All husbands

Table C.2 Sampling errors - National sample: Turkey 1998

Variable	Value (R)	Standard error (SE)	Number of cases		Design effect (DEFT)	Relative error (SE/R)	Confidence limits	
			Unweighted (N)	Weighted (WN)			R-2SE	R+2SE
WOMEN								
Urban	0.665	0.017	8576	8576	3.299	0.025	0.631	0.699
No education	0.167	0.006	8576	8576	1.499	0.036	0.155	0.179
With secondary education or higher	0.303	0.010	8576	8576	2.107	0.035	0.282	0.324
Never married	0.277	0.006	8576	8576	1.188	0.021	0.266	0.289
Currently married	0.690	0.006	8576	8576	1.197	0.009	0.678	0.702
Married before age 20	0.525	0.009	6813	6856	1.558	0.018	0.507	0.544
Sex before age 18	0.000	0.000	6813	6856	NA	NA	0.000	0.000
Children ever born	2.007	0.027	8576	8576	1.137	0.014	1.952	2.062
Children ever born to women over 40	4.220	0.080	1602	1618	1.237	0.019	4.059	4.381
Children surviving	1.802	0.022	8576	8576	1.066	0.012	1.758	1.846
Know any contraceptive method	0.989	0.002	5893	5921	1.309	0.002	0.986	0.993
Know any modern contraceptive method	0.987	0.002	5893	5921	1.297	0.002	0.983	0.991
Ever used any contraceptive method	0.842	0.006	5893	5921	1.306	0.007	0.829	0.854
Currently using any method	0.639	0.008	5893	5921	1.255	0.012	0.623	0.654
Currently using a modern method	0.377	0.009	5893	5921	1.350	0.023	0.360	0.394
Currently using pill	0.044	0.003	5893	5921	1.295	0.079	0.037	0.050
Currently using IUD	0.198	0.007	5893	5921	1.340	0.035	0.184	0.212
Currently using injections	0.005	0.001	5893	5921	1.066	0.202	0.003	0.007
Currently using Norplant	0.000	0.000	5893	5921	NA	NA	0.000	0.000
Currently using condom	0.082	0.005	5893	5921	1.326	0.058	0.073	0.092
Currently using female sterilisation	0.042	0.003	5893	5921	1.231	0.076	0.036	0.049
Currently using calendar/rhythm method	0.011	0.002	5893	5921	1.579	0.197	0.007	0.015
Currently using withdrawal	0.244	0.008	5893	5921	1.355	0.031	0.229	0.259
Using public sector source	0.558	0.016	2206	2240	1.483	0.028	0.527	0.589
Want no more children	0.621	0.007	5893	5921	1.093	0.011	0.607	0.634
Want to delay at least 2 years	0.136	0.005	5893	5921	1.202	0.039	0.126	0.147
Ideal number of children	2.357	0.017	8191	8215	1.422	0.007	2.323	2.392
Mothers received tetanus injection	0.438	0.015	3565	3459	1.502	0.033	0.408	0.467
Mothers received medical care at birth	0.806	0.014	3565	3459	1.664	0.018	0.778	0.834
Had diarrhoea in the last 2 weeks	0.297	0.010	3403	3299	1.180	0.034	0.277	0.317
Treated with ORS packets	0.143	0.013	1044	979	1.081	0.089	0.118	0.168
Sought medical treatment	0.365	0.018	1044	979	1.098	0.049	0.330	0.401
Having health card, seen	0.385	0.021	707	689	1.107	0.054	0.344	0.427
Received BCG vaccination	0.885	0.016	707	689	1.327	0.018	0.853	0.918
Received DPT vaccination (3 doses)	0.587	0.019	707	689	1.007	0.032	0.549	0.625
Received polio vaccination (3 doses)	0.644	0.021	707	689	1.126	0.032	0.602	0.685
Received measles vaccination	0.785	0.021	707	689	1.347	0.027	0.743	0.827
Fully immunised	0.457	0.023	707	689	1.183	0.049	0.412	0.502
Weight-for-height (below -2 SD)	0.019	0.003	2782	2677	1.004	0.139	0.014	0.024
Height-for-age (below -2 SD)	0.160	0.009	2782	2677	1.254	0.059	0.141	0.179
Weight-for-age (below -2 SD)	0.083	0.006	2782	2677	1.113	0.075	0.070	0.095
Total fertility rate (3 years)	2.609	0.079	NA	24175	1.513	0.030	2.451	2.768
Neonatal mortality rate(0-4 years)	25.814	2.980	3670	3561	1.085	0.115	19.854	31.774
Infant mortality rate (0-4 years)	42.702	4.659	3675	3566	1.312	0.109	33.384	52.020
Child mortality rate (0-4 years)	9.773	1.917	3683	3573	1.073	0.196	5.938	13.607
Under-five mortality rate (0-4 years)	52.058	5.187	3688	3578	1.254	0.100	41.683	62.432
Postneonatal mortality rate (0-4 years)	16.888	2.887	3675	3566	1.252	0.171	11.114	22.662
HUSBANDS								
Urban	0.683	0.019	1971	1971	1.793	0.028	0.646	0.721
No education	0.066	0.007	1971	1971	1.173	0.099	0.053	0.080
With secondary education or higher	0.468	0.016	1971	1971	1.440	0.035	0.436	0.500
Know any contraceptive method	0.979	0.004	1971	1971	1.239	0.004	0.971	0.987
Know any modern contraceptive method	0.971	0.005	1971	1971	1.350	0.005	0.961	0.981
Ever used any contraceptive method	0.821	0.011	1971	1971	1.282	0.013	0.799	0.843
Currently using any method	0.626	0.013	1971	1971	1.165	0.020	0.601	0.652
Currently using a modern method	0.422	0.014	1971	1971	1.300	0.034	0.393	0.451
Currently using pill	0.068	0.007	1971	1971	1.210	0.101	0.054	0.081
Currently using IUD	0.188	0.010	1971	1971	1.168	0.055	0.168	0.209
Currently using injections	0.005	0.002	1971	1971	1.099	0.353	0.001	0.008
Currently using condom	0.114	0.009	1971	1971	1.212	0.076	0.097	0.131
Currently using female sterilisation	0.039	0.004	1971	1971	1.020	0.115	0.030	0.047
Currently using calendar/rhythm method	0.016	0.004	1971	1971	1.300	0.228	0.009	0.024
Currently using withdrawal	0.184	0.010	1971	1971	1.183	0.056	0.163	0.205
Want no more children	0.623	0.012	1971	1971	1.088	0.019	0.600	0.647
Want to delay at least 2 years	0.158	0.009	1971	1971	1.091	0.057	0.140	0.175
Ideal number of children	2.710	0.051	1870	1874	1.006	0.019	2.607	2.812

Table C.3 Sampling errors - Urban areas: Turkey 1998

Variable	Value (R)	Standard error (SE)	Number of cases		Design effect (DEFT)	Relative error (SE/R)	Confidence limits	
			Unweighted (N)	Weighted (WN)			R-2SE	R+2SE
WOMEN								
Urban	1.000	0.000	5702	5704	NA	0.000	1.000	1.000
No education	0.141	0.007	5702	5704	1.574	0.051	0.127	0.156
With secondary education or higher	0.386	0.014	5702	5704	2.230	0.037	0.357	0.415
Never married	0.267	0.006	5702	5704	1.092	0.024	0.254	0.280
Currently married	0.697	0.007	5702	5704	1.147	0.010	0.683	0.711
Married before age 20	0.499	0.012	4624	4670	1.596	0.024	0.475	0.522
Sex before age 18	0.000	0.000	4624	4670	NA	NA	0.000	0.000
Children ever born	1.866	0.035	5702	5704	1.276	0.019	1.797	1.935
Children ever born to women over 40	3.821	0.104	1059	1079	1.405	0.027	3.612	4.030
Children surviving	1.694	0.029	5702	5704	1.219	0.017	1.637	1.752
Know any contraceptive method	0.993	0.001	3966	3978	0.925	0.001	0.991	0.995
Know any modern contraceptive method	0.992	0.001	3966	3978	0.927	0.001	0.989	0.995
Ever used any contraceptive method	0.864	0.007	3966	3978	1.212	0.008	0.851	0.877
Currently using any method	0.667	0.009	3966	3978	1.219	0.014	0.649	0.685
Currently using a modern method	0.408	0.010	3966	3978	1.284	0.025	0.388	0.428
Currently using pill	0.046	0.004	3966	3978	1.234	0.089	0.038	0.054
Currently using IUD	0.210	0.009	3966	3978	1.321	0.041	0.193	0.228
Currently using injections	0.005	0.001	3966	3978	1.076	0.230	0.003	0.008
Currently using Norplant	0.000	0.000	3966	3978	NA	NA	0.000	0.000
Currently using condom	0.093	0.006	3966	3978	1.313	0.065	0.081	0.105
Currently using female sterilisation	0.047	0.004	3966	3978	1.238	0.089	0.039	0.055
Currently using calendar/rhythm method	0.014	0.003	3966	3978	1.674	0.226	0.007	0.020
Currently using withdrawal	0.238	0.010	3966	3978	1.429	0.041	0.219	0.258
Using public sector source	0.544	0.019	1591	1629	1.555	0.036	0.505	0.582
Want no more children/sterilised	0.610	0.009	3966	3978	1.100	0.014	0.593	0.627
Want to delay at least 2 years	0.138	0.007	3966	3978	1.195	0.048	0.125	0.151
Ideal number of children	2.304	0.019	5449	5481	1.306	0.008	2.266	2.342
Mothers received tetanus injection	0.435	0.018	2294	2162	1.460	0.040	0.399	0.470
Mothers received medical care at birth	0.877	0.011	2294	2162	1.284	0.013	0.855	0.899
Had diarrhoea in the last 2 weeks	0.261	0.012	2206	2081	1.160	0.045	0.238	0.285
Treated with ORS packets	0.164	0.018	623	544	1.098	0.110	0.128	0.201
Sought medical treatment	0.427	0.023	623	544	1.050	0.055	0.380	0.474
Having health card, seen	0.455	0.028	448	424	1.168	0.062	0.398	0.512
Received BCG vaccination	0.916	0.016	448	424	1.216	0.018	0.883	0.949
Received DPT vaccination (3 doses)	0.639	0.024	448	424	1.013	0.037	0.591	0.686
Received polio vaccination (3 doses)	0.690	0.026	448	424	1.140	0.037	0.638	0.741
Received measles vaccination	0.822	0.028	448	424	1.502	0.034	0.766	0.878
Fully immunised	0.512	0.029	448	424	1.191	0.057	0.454	0.571
Weight-for-height (below -2 SD)	0.017	0.003	1814	1696	1.067	0.197	0.010	0.024
Height-for-age (below -2 SD)	0.126	0.008	1814	1696	0.973	0.066	0.109	0.142
Weight-for-age (below -2 SD)	0.062	0.006	1814	1696	0.907	0.092	0.051	0.073
Total fertility rate (3 years)	2.386	0.086	NA	16165	1.489	0.036	2.213	2.559
Neonatal mortality rate (0-4 years)	23.513	3.591	2359	2221	1.085	0.153	16.330	30.696
Infant mortality rate (0-4 years)	35.217	4.655	2363	2225	1.165	0.132	25.907	44.527
Child mortality rate (0-4 years)	7.455	2.053	2366	2226	1.077	0.275	3.350	11.561
Under-five mortality rate (0-4 years)	42.410	5.306	2370	2231	1.159	0.125	31.798	53.021
Postneonatal mortality rate (0-4 years)	11.704	2.398	2363	2225	1.043	0.205	6.907	16.501
HUSBANDS								
Urban	1.000	0.000	1312	1347	NA	0.000	1.000	1.000
No education	0.047	0.006	1312	1347	1.036	0.129	0.035	0.059
With secondary education or higher	0.563	0.019	1312	1347	1.403	0.034	0.524	0.601
Know any contraceptive method	0.986	0.004	1312	1347	1.147	0.004	0.978	0.993
Know any modern contraceptive method	0.981	0.005	1312	1347	1.326	0.005	0.971	0.991
Ever used any contraceptive method	0.841	0.012	1312	1347	1.214	0.015	0.817	0.866
Currently using any method	0.660	0.015	1312	1347	1.173	0.023	0.629	0.691
Currently using a modern method	0.459	0.018	1312	1347	1.300	0.039	0.423	0.494
Currently using pill	0.074	0.008	1312	1347	1.151	0.112	0.058	0.091
Currently using IUD	0.200	0.013	1312	1347	1.176	0.065	0.174	0.226
Currently using injections	0.005	0.002	1312	1347	1.033	0.397	0.001	0.009
Currently using condom	0.132	0.012	1312	1347	1.236	0.087	0.109	0.155
Currently using female sterilisation	0.038	0.005	1312	1347	1.042	0.145	0.027	0.049
Currently using calendar/rhythm method	0.020	0.005	1312	1347	1.308	0.256	0.010	0.030
Currently using withdrawal	0.178	0.012	1312	1347	1.150	0.068	0.154	0.202
Want no more children/sterilised	0.617	0.014	1312	1347	1.009	0.022	0.590	0.644
Want to delay at least 2 years	0.169	0.011	1312	1347	1.037	0.063	0.148	0.191
Ideal number of children	2.676	0.065	1251	1292	1.010	0.024	2.546	2.805

Table C.4 Sampling errors - Rural areas: Turkey 1998

Variable	Value (R)	Standard error (SE)	Number of cases		Design effect (DEFT)	Relative error (SE/R)	Confidence limits	
			Unweighted (N)	Weighted (WN)			R-2SE	R+2SE
WOMEN								
Urban	0.000	0.000	2874	2872	NA	NA	0.000	0.000
No education	0.219	0.016	2874	2872	2.030	0.072	0.187	0.250
With secondary education or higher	0.138	0.012	2874	2872	1.876	0.088	0.114	0.162
Never married	0.299	0.011	2874	2872	1.341	0.038	0.276	0.322
Currently married	0.676	0.011	2874	2872	1.297	0.017	0.654	0.699
Married before age 20	0.583	0.016	2189	2187	1.506	0.027	0.551	0.615
Sex before age 18	0.000	0.000	2189	2187	NA	NA	0.000	0.000
Children ever born	2.285	0.051	2874	2872	1.071	0.022	2.184	2.387
Children ever born to women over 40	5.018	0.140	543	539	1.178	0.028	4.738	5.298
Children surviving	2.016	0.040	2874	2872	0.983	0.020	1.937	2.096
Know any contraceptive method	0.982	0.005	1927	1943	1.580	0.005	0.972	0.991
Know any modern contraceptive method	0.976	0.005	1927	1943	1.539	0.005	0.966	0.987
Ever used any contraceptive method	0.796	0.016	1927	1943	1.714	0.020	0.765	0.828
Currently using any method	0.581	0.016	1927	1943	1.407	0.027	0.549	0.612
Currently using a modern method	0.314	0.014	1927	1943	1.343	0.045	0.285	0.342
Currently using pill	0.038	0.006	1927	1943	1.447	0.165	0.026	0.051
Currently using IUD	0.173	0.011	1927	1943	1.283	0.064	0.150	0.195
Currently using injections	0.003	0.001	1927	1943	1.003	0.416	0.001	0.006
Currently using Norplant	0.000	0.000	1927	1943	NA	NA	0.000	0.000
Currently using condom	0.060	0.007	1927	1943	1.303	0.117	0.046	0.074
Currently using female sterilisation	0.033	0.005	1927	1943	1.182	0.145	0.024	0.043
Currently using calendar/rhythm method	0.005	0.002	1927	1943	1.115	0.357	0.001	0.009
Currently using withdrawal	0.255	0.012	1927	1943	1.218	0.047	0.231	0.280
Using public sector source	0.597	0.027	615	610	1.377	0.046	0.542	0.651
Want no more children/sterilised	0.642	0.012	1927	1943	1.129	0.019	0.617	0.666
Want to delay at least 2 years	0.134	0.010	1927	1943	1.239	0.072	0.114	0.153
Ideal number of children	2.464	0.039	2742	2734	1.725	0.016	2.387	2.541
Mothers received tetanus injection	0.443	0.027	1271	1297	1.648	0.061	0.388	0.497
Mothers received medical care at birth	0.688	0.030	1271	1297	1.849	0.044	0.627	0.748
Had diarrhoea in the last 2 weeks	0.357	0.019	1197	1217	1.273	0.053	0.319	0.395
Treated with ORS packets	0.116	0.018	421	435	1.065	0.151	0.081	0.151
Sought medical treatment	0.289	0.025	421	435	1.091	0.086	0.239	0.338
Having health card, seen	0.275	0.029	259	265	1.067	0.107	0.216	0.333
Received BCG vaccination	0.837	0.032	259	265	1.424	0.039	0.772	0.901
Received DPT vaccination (3 doses)	0.503	0.032	259	265	1.046	0.064	0.439	0.568
Received polio vaccination (3 doses)	0.570	0.035	259	265	1.131	0.061	0.500	0.639
Received measles vaccination	0.727	0.032	259	265	1.173	0.044	0.663	0.791
Fully immunised	0.368	0.035	259	265	1.187	0.096	0.298	0.439
Weight-for-height (below -2 SD)	0.023	0.005	968	981	0.992	0.207	0.013	0.032
Height-for-age (below -2 SD)	0.220	0.021	968	981	1.522	0.096	0.177	0.262
Weight-for-age (below -2 SD)	0.119	0.014	968	981	1.310	0.117	0.091	0.147
Total fertility rate (3 years)	3.085	0.151	NA	8009	1.479	0.049	2.783	3.387
Neonatal mortality rate (0-4 years)	29.634	5.236	1311	1340	1.088	0.177	19.163	40.105
Infant mortality rate (0-4 years)	55.014	9.449	1312	1341	1.426	0.172	36.116	73.912
Child mortality rate (0-4 years)	13.749	3.805	1317	1346	1.072	0.277	6.138	21.360
Under-five mortality rate (0-4 years)	68.007	10.251	1318	1347	1.318	0.151	47.504	88.509
Postneonatal mortality rate (0-4 years)	25.380	6.367	1312	1341	1.359	0.251	12.646	38.114
HUSBANDS								
Urban	0.000	0.000	659	624	NA	NA	0.000	0.000
No education	0.108	0.016	659	624	1.294	0.145	0.077	0.140
With secondary education or higher	0.264	0.026	659	624	1.532	0.100	0.212	0.317
Know any contraceptive method	0.964	0.010	659	624	1.332	0.010	0.945	0.983
Know any modern contraceptive method	0.950	0.012	659	624	1.356	0.012	0.927	0.973
Ever used any contraceptive method	0.777	0.022	659	624	1.377	0.029	0.733	0.822
Currently using any method	0.554	0.023	659	624	1.173	0.041	0.508	0.599
Currently using a modern method	0.344	0.024	659	624	1.274	0.069	0.297	0.392
Currently using pill	0.054	0.012	659	624	1.348	0.220	0.030	0.077
Currently using IUD	0.162	0.016	659	624	1.136	0.101	0.129	0.195
Currently using injections	0.004	0.003	659	624	1.253	0.735	0.000	0.011
Currently using condom	0.075	0.010	659	624	1.008	0.138	0.054	0.096
Currently using female sterilisation	0.041	0.008	659	624	0.989	0.187	0.026	0.056
Currently using calendar/rhythm method	0.009	0.004	659	624	1.211	0.494	0.000	0.018
Currently using withdrawal	0.197	0.020	659	624	1.277	0.100	0.158	0.237
Want no more children/sterilised	0.637	0.023	659	624	1.218	0.036	0.591	0.682
Want to delay at least 2 years	0.132	0.016	659	624	1.221	0.122	0.100	0.164
Ideal number of children	2.785	0.098	619	582	1.201	0.035	2.588	2.982

Table C.5 Sampling errors - West region: Turkey 1998

Variable	Value (R)	Standard error (SE)	Number of cases		Design effect (DEFT)	Relative error (SE/R)	Confidence limits	
			Unweighted (N)	Weighted (WN)			R-2SE	R+2SE
WOMEN								
Urban	0.815	0.019	2170	3204	2.240	0.023	0.778	0.852
No education	0.099	0.008	2170	3204	1.289	0.084	0.082	0.115
With secondary education or higher	0.371	0.020	2170	3204	1.923	0.054	0.331	0.411
Never married	0.256	0.010	2170	3204	1.052	0.039	0.236	0.275
Currently married	0.706	0.010	2170	3204	1.054	0.015	0.685	0.726
Married before age 20	0.486	0.019	1806	2666	1.574	0.038	0.449	0.523
Sex before age 18	0.000	0.000	1806	2666	NA	NA	0.000	0.000
Children ever born	1.690	0.047	2170	3204	1.230	0.028	1.596	1.784
Children ever born to women over 40	3.434	0.133	443	645	1.360	0.039	3.169	3.700
Children surviving	1.532	0.038	2170	3204	1.172	0.025	1.456	1.608
Know any contraceptive method	0.993	0.002	1549	2261	0.935	0.002	0.989	0.997
Know any modern contraceptive method	0.992	0.002	1549	2261	0.954	0.002	0.988	0.996
Ever used any contraceptive method	0.902	0.008	1549	2261	1.025	0.009	0.887	0.918
Currently using any method	0.705	0.013	1549	2261	1.130	0.019	0.679	0.732
Currently using a modern method	0.405	0.016	1549	2261	1.301	0.040	0.373	0.437
Currently using pill	0.054	0.007	1549	2261	1.172	0.125	0.040	0.067
Currently using IUD	0.205	0.014	1549	2261	1.360	0.068	0.177	0.233
Currently using injections	0.003	0.001	1549	2261	1.062	0.508	0.000	0.006
Currently using Norplant	0.000	0.000	1549	2261	NA	NA	0.000	0.000
Currently using condom	0.094	0.010	1549	2261	1.294	0.102	0.075	0.113
Currently using female sterilisation	0.042	0.006	1549	2261	1.239	0.151	0.029	0.054
Currently using calendar/rhythm method	0.016	0.005	1549	2261	1.550	0.312	0.006	0.026
Currently using withdrawal	0.276	0.016	1549	2261	1.380	0.057	0.245	0.307
Using public sector source	0.489	0.029	636	919	1.483	0.060	0.430	0.548
Want no more children/sterilised	0.609	0.013	1549	2261	1.014	0.021	0.584	0.634
Want to delay at least 2 years	0.134	0.010	1549	2261	1.103	0.071	0.115	0.153
Ideal number of children	2.164	0.021	2105	3113	1.241	0.010	2.121	2.207
Mothers received tetanus injection	0.365	0.025	694	1031	1.241	0.068	0.316	0.415
Mothers received medical care at birth	0.923	0.012	694	1031	1.087	0.013	0.898	0.947
Had diarrhoea in the last 2 weeks	0.218	0.019	671	995	1.156	0.086	0.180	0.255
Treated with ORS packets	0.128	0.034	144	217	1.221	0.262	0.061	0.196
Sought medical treatment	0.370	0.046	144	217	1.116	0.124	0.278	0.461
Having health card, seen	0.478	0.046	148	221	1.128	0.096	0.386	0.570
Received BCG vaccination	0.958	0.018	148	221	1.105	0.019	0.921	0.994
Received DPT vaccination (3 doses)	0.614	0.034	148	221	0.852	0.055	0.546	0.682
Received polio vaccination (3 doses)	0.723	0.042	148	221	1.152	0.058	0.639	0.807
Received measles vaccination	0.806	0.049	148	221	1.527	0.061	0.707	0.905
Fully immunised	0.502	0.044	148	221	1.077	0.088	0.414	0.590
Weight-for-height (below -2 SD)	0.015	0.005	515	763	1.012	0.362	0.004	0.026
Height-for-age (below -2 SD)	0.099	0.013	515	763	0.968	0.130	0.073	0.125
Weight-for-age (below -2 SD)	0.038	0.009	515	763	1.053	0.245	0.019	0.056
Total fertility rate (3 years)	2.032	0.122	NA	9140	1.406	0.060	1.788	2.277
Neonatal mortality rate (0-4 years)	25.288	6.286	711	1056	1.074	0.249	12.716	37.859
Infant mortality rate (0-4 years)	32.789	7.584	712	1058	1.107	0.231	17.621	47.957
Child mortality rate (0-4 years)	5.647	3.145	711	1056	1.115	0.557	0.000	11.937
Under-five mortality rate (0-4 years)	38.251	8.207	712	1058	1.125	0.215	21.837	54.665
Postneonatal mortality rate (0-4 years)	7.501	3.155	712	1058	1.034	0.421	1.192	13.810
HUSBANDS								
Urban	0.823	0.018	546	767	1.127	0.022	0.786	0.860
No education	0.036	0.009	546	767	1.077	0.240	0.019	0.053
With secondary education or higher	0.522	0.026	546	767	1.238	0.051	0.469	0.575
Know any contraceptive method	0.987	0.006	546	767	1.222	0.006	0.975	0.999
Know any modern contraceptive method	0.982	0.008	546	767	1.429	0.008	0.966	0.998
Ever used any contraceptive method	0.881	0.018	546	767	1.278	0.020	0.846	0.916
Currently using any method	0.700	0.024	546	767	1.197	0.034	0.653	0.747
Currently using a modern method	0.438	0.027	546	767	1.276	0.062	0.384	0.493
Currently using pill	0.078	0.013	546	767	1.108	0.164	0.052	0.103
Currently using IUD	0.194	0.021	546	767	1.215	0.106	0.153	0.235
Currently using injections	0.002	0.002	546	767	1.001	0.999	0.000	0.006
Currently using condom	0.126	0.017	546	767	1.206	0.136	0.092	0.161
Currently using female sterilisation	0.026	0.007	546	767	1.047	0.272	0.012	0.041
Currently using calendar/rhythm method	0.025	0.008	546	767	1.218	0.324	0.009	0.042
Currently using withdrawal	0.233	0.020	546	767	1.116	0.087	0.193	0.273
Want no more children/sterilised	0.614	0.019	546	767	0.887	0.030	0.577	0.651
Want to delay at least 2 years	0.178	0.017	546	767	1.022	0.094	0.145	0.212
Ideal number of children	2.328	0.055	530	744	1.007	0.024	2.218	2.43

Table C.6 Sampling errors - South region: Turkey 1998

Variable	Value (R)	Standard error (SE)	Number of cases		Design effect (DEFT)	Relative error (SE/R)	Confidence limits	
			Unweighted (N)	Weighted (WN)			R-2SE	R+2SE
WOMEN								
Urban	0.641	0.034	1678	1258	2.887	0.053	0.573	0.709
No education	0.182	0.014	1678	1258	1.518	0.078	0.154	0.211
With secondary education or higher	0.260	0.021	1678	1258	1.934	0.080	0.219	0.302
Never married	0.291	0.012	1678	1258	1.090	0.042	0.266	0.315
Currently married	0.676	0.012	1678	1258	1.069	0.018	0.652	0.701
Married before age 20	0.511	0.018	1329	997	1.336	0.036	0.474	0.547
Sex before age 18	0.000	0.000	1329	997	NA	NA	0.000	0.000
Children ever born	2.050	0.051	1678	1258	0.935	0.025	1.947	2.153
Children ever born to women over 40	4.461	0.154	317	244	1.071	0.034	4.153	4.768
Children surviving	1.893	0.046	1678	1258	0.922	0.024	1.801	1.984
Know any contraceptive method	0.993	0.003	1126	851	1.190	0.003	0.987	0.999
Know any modern contraceptive method	0.993	0.003	1126	851	1.190	0.003	0.987	0.999
Ever used any contraceptive method	0.820	0.015	1126	851	1.337	0.019	0.790	0.851
Currently using any method	0.603	0.014	1126	851	0.930	0.022	0.576	0.630
Currently using a modern method	0.351	0.016	1126	851	1.118	0.045	0.319	0.383
Currently using pill	0.023	0.004	1126	851	1.003	0.195	0.014	0.032
Currently using IUD	0.209	0.015	1126	851	1.220	0.071	0.180	0.239
Currently using injections	0.005	0.002	1126	851	0.953	0.405	0.001	0.009
Currently using Norplant	0.000	0.000	1126	851	NA	NA	0.000	0.000
Currently using condom	0.062	0.007	1126	851	1.022	0.119	0.047	0.076
Currently using female sterilisation	0.045	0.007	1126	851	1.156	0.159	0.031	0.059
Currently using calendar/rhythm method	0.004	0.002	1126	851	0.957	0.446	0.000	0.008
Currently using withdrawal	0.242	0.013	1126	851	1.008	0.053	0.216	0.268
Using public sector source	0.672	0.027	402	299	1.148	0.040	0.618	0.726
Want no more children/sterilised	0.594	0.016	1126	851	1.100	0.027	0.562	0.626
Want to delay at least 2 years	0.129	0.010	1126	851	0.978	0.076	0.109	0.148
Ideal number of children	2.521	0.039	1605	1201	1.354	0.015	2.443	2.599
Mothers received tetanus injection	0.642	0.033	665	490	1.536	0.051	0.576	0.707
Mothers received medical care at birth	0.862	0.025	665	490	1.494	0.029	0.813	0.911
Had diarrhoea in the last 2 weeks	0.287	0.019	640	472	0.990	0.065	0.250	0.325
Treated with ORS packets	0.101	0.025	187	136	1.036	0.251	0.050	0.151
Sought medical treatment	0.479	0.039	187	136	0.992	0.082	0.401	0.558
Having health card, seen	0.356	0.051	127	95	1.188	0.143	0.254	0.458
Received BCG vaccination	0.927	0.026	127	95	1.133	0.028	0.875	0.980
Received DPT vaccination (3 doses)	0.674	0.047	127	95	1.111	0.069	0.581	0.768
Received polio vaccination (3 doses)	0.689	0.050	127	95	1.209	0.073	0.589	0.789
Received measles vaccination	0.871	0.035	127	95	1.175	0.040	0.801	0.940
Fully immunised	0.573	0.054	127	95	1.221	0.094	0.466	0.681
Weight-for-height (below -2 SD)	0.022	0.008	555	407	1.207	0.344	0.007	0.037
Height-for-age (below -2 SD)	0.135	0.016	555	407	1.070	0.117	0.104	0.167
Weight-for-age (below -2 SD)	0.089	0.012	555	407	0.965	0.139	0.064	0.114
Total fertility rate (3 years)	2.555	0.150	NA	3554	1.282	0.059	2.254	2.856
Neonatal mortality rate (0-4 years)	16.756	5.567	684	503	1.038	0.332	5.621	27.890
Infant mortality rate (0-4 years)	32.702	8.217	687	505	1.101	0.251	16.267	49.136
Child mortality rate (0-4 years)	10.692	4.234	686	504	0.928	0.396	2.224	19.161
Under-five mortality rate (0-4 years)	43.044	9.775	689	506	1.055	0.227	23.494	62.595
Postneonatal mortality rate (0-4 years)	15.946	5.310	687	505	1.000	0.333	5.325	26.566
HUSBANDS								
Urban	0.663	0.036	400	285	1.505	0.054	0.592	0.735
No education	0.064	0.012	400	285	0.992	0.190	0.040	0.088
With secondary education or higher	0.384	0.026	400	285	1.057	0.067	0.333	0.435
Know any contraceptive method	0.973	0.011	400	285	1.325	0.011	0.952	0.995
Know any modern contraceptive method	0.960	0.014	400	285	1.376	0.014	0.933	0.987
Ever used any contraceptive method	0.818	0.023	400	285	1.185	0.028	0.773	0.864
Currently using any method	0.606	0.025	400	285	1.018	0.041	0.557	0.656
Currently using a modern method	0.417	0.028	400	285	1.145	0.068	0.360	0.473
Currently using pill	0.030	0.008	400	285	0.887	0.252	0.015	0.045
Currently using IUD	0.214	0.018	400	285	0.897	0.086	0.177	0.250
Currently using injections	0.007	0.004	400	285	0.942	0.559	0.000	0.015
Currently using condom	0.097	0.015	400	285	0.980	0.149	0.068	0.126
Currently using female sterilisation	0.053	0.010	400	285	0.926	0.196	0.032	0.074
Currently using calendar/rhythm method	0.009	0.005	400	285	0.971	0.497	0.000	0.019
Currently using withdrawal	0.180	0.019	400	285	1.003	0.107	0.142	0.219
Want no more children/sterilised	0.604	0.025	400	285	1.019	0.041	0.554	0.654
Want to delay at least 2 years	0.154	0.016	400	285	0.860	0.101	0.123	0.185
Ideal number of children	3.007	0.115	382	271	1.046	0.038	2.777	3.238

Table C.7 Sampling errors - Central region: Turkey 1998

Variable	Value (R)	Standard error (SE)	Number of cases		Design effect (DEFT)	Relative error (SE/R)	Confidence limits	
			Unweighted (N)	Weighted (WN)			R-2SE	R+2SE
WOMEN								
Urban	0.617	0.038	1706	1985	3.194	0.061	0.542	0.692
No education	0.098	0.009	1706	1985	1.254	0.092	0.080	0.117
With secondary education or higher	0.324	0.025	1706	1985	2.195	0.077	0.274	0.374
Never married	0.253	0.011	1706	1985	1.089	0.045	0.230	0.276
Currently married	0.719	0.012	1706	1985	1.121	0.017	0.694	0.743
Married before age 20	0.549	0.019	1381	1605	1.448	0.035	0.510	0.588
Sex before age 18	0.000	0.000	1381	1605	NA	NA	0.000	0.000
Children ever born	1.962	0.056	1706	1985	1.183	0.029	1.850	2.074
Children ever born to women over 40	3.845	0.158	321	375	1.264	0.041	3.529	4.161
Children surviving	1.746	0.046	1706	1985	1.145	0.026	1.655	1.838
Know any contraceptive method	1.000	0.000	1226	1426	NA	0.000	1.000	1.000
Know any modern contraceptive method	0.999	0.001	1226	1426	1.067	0.001	0.997	1.001
Ever used any contraceptive method	0.886	0.009	1226	1426	0.988	0.010	0.868	0.904
Currently using any method	0.683	0.015	1226	1426	1.103	0.021	0.654	0.712
Currently using a modern method	0.428	0.016	1226	1426	1.123	0.037	0.396	0.460
Currently using pill	0.049	0.008	1226	1426	1.279	0.161	0.033	0.065
Currently using IUD	0.242	0.014	1226	1426	1.153	0.058	0.214	0.270
Currently using injections	0.005	0.002	1226	1426	1.006	0.394	0.001	0.009
Currently using Norplant	0.000	0.000	1226	1426	NA	NA	0.000	0.000
Currently using condom	0.092	0.010	1226	1426	1.195	0.107	0.073	0.112
Currently using female sterilisation	0.034	0.005	1226	1426	1.041	0.159	0.023	0.045
Currently using calendar/rhythm method	0.011	0.004	1226	1426	1.229	0.338	0.003	0.018
Currently using withdrawal	0.237	0.014	1226	1426	1.128	0.058	0.209	0.264
Using public sector source	0.598	0.032	529	612	1.490	0.053	0.535	0.662
Want no more children/sterilised	0.663	0.013	1226	1426	0.942	0.019	0.637	0.688
Want to delay at least 2 years	0.143	0.012	1226	1426	1.214	0.085	0.119	0.167
Ideal number of children	2.291	0.032	1638	1906	1.254	0.014	2.227	2.355
Mothers received tetanus injection	0.550	0.032	694	795	1.514	0.059	0.485	0.614
Mothers received medical care at birth	0.899	0.017	694	795	1.238	0.019	0.866	0.933
Had diarrhoea in the last 2 weeks	0.310	0.021	662	762	1.068	0.067	0.269	0.352
Treated with ORS packets	0.160	0.030	210	236	1.132	0.189	0.099	0.220
Sought medical treatment	0.299	0.035	210	236	1.059	0.116	0.230	0.369
Having health card, seen	0.413	0.042	137	157	0.980	0.101	0.330	0.497
Received BCG vaccination	0.960	0.017	137	157	0.988	0.017	0.927	0.993
Received DPT vaccination (3 doses)	0.642	0.052	137	157	1.248	0.082	0.537	0.747
Received polio vaccination (3 doses)	0.655	0.050	137	157	1.224	0.077	0.554	0.756
Received measles vaccination	0.864	0.023	137	157	0.778	0.027	0.818	0.910
Fully immunised	0.518	0.055	137	157	1.255	0.105	0.409	0.628
Weight-for-height (below -2 SD)	0.013	0.004	556	643	0.813	0.301	0.005	0.021
Height-for-age (below -2 SD)	0.116	0.016	556	643	1.130	0.135	0.085	0.148
Weight-for-age (below -2 SD)	0.054	0.009	556	643	0.933	0.163	0.037	0.072
Total fertility rate (3 years)	2.564	0.132	NA	5616	1.304	0.052	2.299	2.828
Neonatal mortality rate (0-4 years)	26.779	6.130	722	827	0.997	0.229	14.520	39.039
Infant mortality rate (0-4 years)	41.261	8.886	723	828	1.168	0.215	23.489	59.033
Child mortality rate (0-4 years)	8.717	3.521	723	831	1.041	0.404	1.676	15.758
Under-five mortality rate (0-4 years)	49.618	10.068	726	832	1.193	0.203	29.483	69.753
Postneonatal mortality rate (0-4 years)	14.482	4.608	723	828	1.002	0.318	5.265	23.698
HUSBANDS								
Urban	0.617	0.046	413	481	1.912	0.074	0.525	0.709
No education	0.040	0.010	413	481	1.075	0.261	0.019	0.060
With secondary education or higher	0.512	0.038	413	481	1.552	0.075	0.436	0.589
Know any contraceptive method	0.983	0.006	413	481	1.006	0.006	0.971	0.996
Know any modern contraceptive method	0.977	0.007	413	481	0.991	0.008	0.962	0.992
Ever used any contraceptive method	0.871	0.018	413	481	1.090	0.021	0.835	0.907
Currently using any method	0.679	0.023	413	481	0.999	0.034	0.634	0.725
Currently using a modern method	0.499	0.027	413	481	1.092	0.054	0.445	0.553
Currently using pill	0.085	0.016	413	481	1.148	0.185	0.054	0.117
Currently using IUD	0.224	0.018	413	481	0.900	0.083	0.187	0.261
Currently using injections	0.009	0.005	413	481	1.154	0.598	0.000	0.020
Currently using condom	0.127	0.018	413	481	1.106	0.143	0.091	0.164
Currently using female sterilisation	0.048	0.010	413	481	0.927	0.204	0.028	0.067
Currently using calendar/rhythm method	0.016	0.007	413	481	1.109	0.426	0.002	0.030
Currently using withdrawal	0.158	0.022	413	481	1.200	0.137	0.115	0.201
Want no more children/sterilised	0.641	0.029	413	481	1.242	0.046	0.582	0.700
Want to delay at least 2 years	0.146	0.019	413	481	1.065	0.127	0.109	0.183
Ideal number of children	2.462	0.060	392	459	1.143	0.024	2.343	2.582

Table C.8 Sampling errors - North region: Turkey 1998

Variable	Value (R)	Standard error (SE)	Number of cases		Design effect (DEFT)	Relative error (SE/R)	Confidence limits	
			Unweighted (N)	Weighted (WN)			R-2SE	R+2SE
WOMEN								
Urban	0.449	0.039	1258	692	2.767	0.087	0.371	0.526
No education	0.183	0.015	1258	692	1.413	0.084	0.152	0.214
With secondary education or higher	0.278	0.021	1258	692	1.641	0.075	0.236	0.319
Never married	0.286	0.018	1258	692	1.396	0.062	0.250	0.321
Currently married	0.686	0.019	1258	692	1.419	0.027	0.649	0.723
Married before age 20	0.510	0.021	1019	556	1.349	0.041	0.468	0.552
Sex before age 18	0.000	0.000	1019	556	NA	NA	0.000	0.000
Children ever born	2.167	0.077	1258	692	1.208	0.036	2.013	2.321
Children ever born to women over 40	4.357	0.179	269	149	1.251	0.041	3.999	4.715
Children surviving	1.937	0.062	1258	692	1.148	0.032	1.814	2.061
Know any contraceptive method	0.998	0.002	862	474	1.244	0.002	0.995	1.000
Know any modern contraceptive method	0.990	0.005	862	474	1.563	0.005	0.980	1.000
Ever used any contraceptive method	0.880	0.014	862	474	1.258	0.016	0.852	0.908
Currently using any method	0.670	0.019	862	474	1.187	0.028	0.632	0.708
Currently using a modern method	0.352	0.027	862	474	1.643	0.076	0.298	0.405
Currently using pill	0.045	0.007	862	474	1.048	0.165	0.030	0.060
Currently using IUD	0.124	0.013	862	474	1.157	0.105	0.098	0.150
Currently using injections	0.000	0.000	862	474	NA	NA	0.000	0.000
Currently using Norplant	0.000	0.000	862	474	NA	NA	0.000	0.000
Currently using condom	0.092	0.011	862	474	1.139	0.122	0.069	0.114
Currently using female sterilisation	0.084	0.013	862	474	1.354	0.152	0.058	0.110
Currently using calendar/rhythm method	0.007	0.002	862	474	0.851	0.356	0.002	0.011
Currently using withdrawal	0.309	0.019	862	474	1.223	0.062	0.270	0.347
Using public sector source	0.451	0.031	318	168	1.099	0.068	0.389	0.512
Want no more children/sterilised	0.614	0.020	862	474	1.191	0.032	0.574	0.653
Want to delay at least 2 years	0.099	0.010	862	474	0.949	0.098	0.079	0.118
Ideal number of children	2.234	0.038	1237	680	1.468	0.017	2.158	2.311
Mothers received tetanus injection	0.600	0.035	460	271	1.364	0.059	0.529	0.670
Mothers received medical care at birth	0.897	0.019	460	271	1.128	0.021	0.860	0.935
Had diarrhoea in the last 2 weeks	0.264	0.036	444	260	1.707	0.135	0.193	0.336
Treated with ORS packets	0.168	0.044	115	69	1.225	0.265	0.079	0.256
Sought medical treatment	0.316	0.051	115	69	1.215	0.160	0.215	0.417
Having health card, seen	0.523	0.058	94	52	1.121	0.110	0.408	0.638
Received BCG vaccination	0.975	0.017	94	52	1.095	0.018	0.941	1.000
Received DPT vaccination (3 doses)	0.714	0.049	94	52	1.048	0.068	0.617	0.812
Received polio vaccination (3 doses)	0.738	0.046	94	52	1.015	0.062	0.646	0.830
Received measles vaccination	0.840	0.049	94	52	1.292	0.058	0.743	0.938
Fully immunised	0.589	0.053	94	52	1.052	0.090	0.483	0.696
Weight-for-height (below -2 SD)	0.016	0.009	361	211	1.445	0.580	0.003	0.034
Height-for-age (below -2 SD)	0.128	0.026	361	211	1.415	0.203	0.076	0.180
Weight-for-age (below -2 SD)	0.048	0.018	361	211	1.550	0.376	0.012	0.084
Total fertility rate (3 years)	2.679	0.173	NA	1935	1.181	0.065	2.333	3.025
Neonatal mortality rate (0-4 years)	19.657	6.821	475	282	1.130	0.347	6.016	33.298
Infant mortality rate (0-4 years)	42.036	9.878	475	282	1.076	0.235	22.281	61.792
Child mortality rate (0-4 years)	8.875	5.660	477	284	1.221	0.638	0.000	20.194
Under-five mortality rate (0-4 years)	50.538	11.613	477	284	1.043	0.230	27.312	73.765
Postneonatal mortality rate (0-4 years)	22.379	8.003	475	282	1.118	0.358	6.374	38.384
HUSBANDS								
Urban	0.467	0.058	299	150	1.999	0.124	0.351	0.582
No education	0.051	0.014	299	150	1.102	0.275	0.023	0.079
With secondary education or higher	0.407	0.043	299	150	1.507	0.105	0.322	0.493
Know any contraceptive method	0.969	0.017	299	150	1.656	0.017	0.936	1.000
Know any modern contraceptive method	0.961	0.017	299	150	1.544	0.018	0.926	0.995
Ever used any contraceptive method	0.792	0.038	299	150	1.598	0.047	0.717	0.867
Currently using any method	0.585	0.035	299	150	1.232	0.060	0.515	0.655
Currently using a modern method	0.369	0.038	299	150	1.349	0.102	0.294	0.445
Currently using pill	0.079	0.021	299	150	1.346	0.267	0.037	0.121
Currently using IUD	0.112	0.015	299	150	0.810	0.132	0.083	0.142
Currently using injections	0.000	0.000	299	150	NA	NA	0.000	0.000
Currently using condom	0.115	0.017	299	150	0.941	0.151	0.081	0.150
Currently using female sterilisation	0.057	0.015	299	150	1.140	0.268	0.027	0.088
Currently using calendar/rhythm method	0.004	0.003	299	150	0.793	0.705	0.000	0.010
Currently using withdrawal	0.207	0.021	299	150	0.904	0.103	0.164	0.249
Want no more children/sterilised	0.634	0.025	299	150	0.907	0.040	0.583	0.684
Want to delay at least 2 years	0.113	0.014	299	150	0.750	0.122	0.086	0.141
Ideal number of children	2.696	0.122	286	144	0.791	0.045	2.451	2.940

Table C.9 Sampling errors - East region: Turkey 1998

Variable	Value (R)	Standard error (SE)	Number of cases		Design effect (DEFT)	Relative error (SE/R)	Confidence limits	
			Unweighted (N)	Weighted (WN)			R-2SE	R+2SE
WOMEN								
Urban	0.522	0.056	1764	1437	4.698	0.107	0.410	0.634
No education	0.395	0.021	1764	1437	1.841	0.054	0.352	0.437
With secondary education or higher	0.171	0.018	1764	1437	2.061	0.108	0.134	0.208
Never married	0.344	0.015	1764	1437	1.306	0.043	0.315	0.374
Currently married	0.632	0.015	1764	1437	1.315	0.024	0.602	0.663
Married before age 20	0.612	0.017	1278	1033	1.278	0.028	0.578	0.647
Sex before age 18	0.000	0.000	1278	1033	NA	NA	0.000	0.000
Children ever born	2.659	0.074	1764	1437	0.986	0.028	2.510	2.808
Children ever born to women over 40	7.003	0.232	252	204	1.207	0.033	6.538	7.467
Children surviving	2.338	0.059	1764	1437	0.917	0.025	2.219	2.457
Know any contraceptive method	0.956	0.009	1130	909	1.546	0.010	0.937	0.975
Know any modern contraceptive method	0.947	0.010	1130	909	1.521	0.011	0.927	0.968
Ever used any contraceptive method	0.623	0.025	1130	909	1.711	0.040	0.574	0.673
Currently using any method	0.420	0.024	1130	909	1.617	0.057	0.372	0.467
Currently using a modern method	0.267	0.020	1130	909	1.501	0.074	0.228	0.307
Currently using pill	0.029	0.006	1130	909	1.174	0.204	0.017	0.040
Currently using IUD	0.140	0.011	1130	909	1.110	0.082	0.117	0.163
Currently using injections	0.011	0.003	1130	909	1.099	0.316	0.004	0.017
Currently using Norplant	0.000	0.000	1130	909	NA	NA	0.000	0.000
Currently using condom	0.052	0.008	1130	909	1.177	0.149	0.037	0.068
Currently using female sterilisation	0.033	0.005	1130	909	1.026	0.164	0.022	0.044
Currently using calendar/rhythm method	0.007	0.003	1130	909	1.091	0.377	0.002	0.013
Currently using withdrawal	0.144	0.014	1130	909	1.303	0.094	0.117	0.172
Using public sector source	0.652	0.031	321	243	1.174	0.048	0.589	0.714
Want no more children/sterilised	0.611	0.017	1130	909	1.184	0.028	0.577	0.646
Want to delay at least 2 years	0.158	0.013	1130	909	1.228	0.084	0.131	0.184
Ideal number of children	2.825	0.067	1606	1315	1.677	0.024	2.691	2.959
Mothers received tetanus injection	0.256	0.027	1052	871	1.713	0.107	0.201	0.310
Mothers received medical care at birth	0.523	0.036	1052	871	1.877	0.069	0.451	0.595
Had diarrhoea in the last 2 weeks	0.397	0.021	986	810	1.237	0.053	0.355	0.439
Treated with ORS packets	0.153	0.016	388	322	0.849	0.108	0.120	0.186
Sought medical treatment	0.374	0.029	388	322	1.121	0.079	0.315	0.433
Having health card, seen	0.208	0.028	201	165	0.965	0.135	0.152	0.264
Received BCG vaccination	0.664	0.051	201	165	1.536	0.077	0.562	0.767
Received DPT vaccination (3 doses)	0.406	0.031	201	165	0.897	0.077	0.343	0.468
Received polio vaccination (3 doses)	0.470	0.031	201	165	0.874	0.066	0.408	0.532
Received measles vaccination	0.617	0.043	201	165	1.240	0.069	0.531	0.702
Fully immunised	0.229	0.033	201	165	1.111	0.146	0.162	0.296
Weight-for-height (below -2 SD)	0.029	0.006	795	652	0.970	0.197	0.018	0.041
Height-for-age (below -2 SD)	0.300	0.023	795	652	1.281	0.075	0.255	0.346
Weight-for-age (below -2 SD)	0.171	0.017	795	652	1.177	0.097	0.138	0.204
Total fertility rate (3 years)	4.191	0.225	NA	3931	1.362	0.054	3.741	4.641
Neonatal mortality rate (0-4 years)	32.579	6.269	1078	892	1.113	0.192	20.042	45.116
Infant mortality rate (0-4 years)	61.525	12.419	1078	892	1.648	0.202	36.686	86.364
Child mortality rate (0-4 years)	15.354	5.008	1084	897	1.221	0.326	5.338	25.370
Under-five mortality rate (0-4 years)	75.934	13.537	1084	897	1.510	0.178	48.859	103.009
Postneonatal mortality rate (0-4 years)	28.946	8.887	1078	892	1.620	0.307	11.171	46.721
HUSBANDS								
Urban	0.555	0.062	313	287	2.217	0.112	0.430	0.680
No education	0.205	0.031	313	287	1.364	0.152	0.142	0.267
With secondary education or higher	0.364	0.043	313	287	1.591	0.119	0.278	0.451
Know any contraceptive method	0.959	0.014	313	287	1.264	0.015	0.931	0.987
Know any modern contraceptive method	0.948	0.018	313	287	1.440	0.019	0.912	0.984
Ever used any contraceptive method	0.594	0.038	313	287	1.367	0.064	0.518	0.670
Currently using any method	0.381	0.032	313	287	1.175	0.085	0.316	0.446
Currently using a modern method	0.285	0.032	313	287	1.249	0.112	0.221	0.348
Currently using pill	0.043	0.012	313	287	1.053	0.281	0.019	0.067
Currently using IUD	0.129	0.023	313	287	1.197	0.176	0.083	0.174
Currently using injections	0.007	0.005	313	287	1.026	0.712	0.000	0.016
Currently using condom	0.074	0.015	313	287	0.997	0.199	0.045	0.104
Currently using female sterilisation	0.032	0.011	313	287	1.090	0.340	0.010	0.054
Currently using calendar/rhythm method	0.005	0.004	313	287	0.917	0.717	0.000	0.013
Currently using withdrawal	0.089	0.016	313	287	1.009	0.183	0.056	0.121
Want no more children/sterilised	0.633	0.031	313	287	1.140	0.049	0.571	0.695
Want to delay at least 2 years	0.148	0.023	313	287	1.165	0.158	0.101	0.195
Ideal number of children	3.955	0.293	280	256	1.154	0.074	3.369	4.541

APPENDIX D

DATA QUALITY TABLES

APPENDIX D

DATA QUALITY TABLES

Table D.1 Household age distribution

Single-year age distribution of the de facto household population by sex (weighted), Turkey 1998

Age	Males		Females		Age	Males		Females	
	Number	Percent	Number	Percent		Number	Percent	Number	Percent
0	387	2.4	367	2.1	38	249	1.5	262	1.5
1	377	2.3	335	1.9	39	160	1.0	189	1.1
2	344	2.1	304	1.8	40	255	1.5	239	1.4
3	333	2.0	310	1.8	41	156	0.9	162	0.9
4	349	2.1	339	2.0	42	218	1.3	191	1.1
5	336	2.0	316	1.8	43	182	1.1	209	1.2
6	409	2.5	388	2.2	44	168	1.0	174	1.0
7	346	2.1	350	2.0	45	193	1.2	206	1.2
8	403	2.4	369	2.1	46	160	1.0	160	0.9
9	307	1.9	335	1.9	47	124	0.8	122	0.7
10	389	2.4	323	1.9	48	197	1.2	197	1.1
11	361	2.2	340	2.0	49	128	0.8	102	0.6
12	361	2.2	378	2.2	50	172	1.0	175	1.0
13	386	2.3	379	2.2	51	106	0.6	149	0.9
14	322	2.0	403	2.3	52	139	0.8	119	0.7
15	329	2.0	330	1.9	53	106	0.6	127	0.7
16	347	2.1	421	2.4	54	97	0.6	115	0.7
17	370	2.2	378	2.2	55	148	0.9	195	1.1
18	404	2.5	460	2.7	56	93	0.6	118	0.7
19	307	1.9	318	1.8	57	103	0.6	92	0.5
20	240	1.5	380	2.2	58	114	0.7	119	0.7
21	203	1.2	320	1.8	59	78	0.5	95	0.5
22	290	1.8	330	1.9	60	197	1.2	193	1.1
23	291	1.8	324	1.9	61	64	0.4	48	0.3
24	302	1.8	347	2.0	62	96	0.6	74	0.4
25	325	2.0	328	1.9	63	72	0.4	86	0.5
26	304	1.8	338	1.9	64	75	0.5	70	0.4
27	266	1.6	276	1.6	65	165	1.0	163	0.9
28	254	1.5	285	1.6	66	79	0.5	94	0.5
29	206	1.2	256	1.5	67	65	0.4	79	0.5
30	266	1.6	291	1.7	68	63	0.4	81	0.5
31	193	1.2	180	1.0	69	48	0.3	42	0.2
32	225	1.4	251	1.4	70+	501	3.0	602	3.5
33	255	1.6	277	1.6	Don't know/ Missing	14	0.1	8	0.0
34	236	1.4	256	1.5					
35	240	1.5	237	1.4					
36	199	1.2	229	1.3	Total	16,464	100.0	17,340	100.0
37	215	1.3	236	1.4					

Note: The de facto population includes all residents and nonresidents who slept in the household the night before the interview.

Table D.2 Age distribution of eligible and interviewed women

Percent distribution of the de facto household population of women age 10-54 and of interviewed women age 15-49, and the percentage of eligible women who were interviewed (weighted) by five-year age groups, Turkey 1998

Age	Household population		Women interviewed		Percent interviewed (weighted)
	Number	Percent	Number	Percent	
10-14	1,823	-	-	-	-
15-19	1,907	20.6	1,673	20.1	87.7
20-24	1,701	18.4	1,519	18.3	89.3
25-29	1,483	16.0	1,350	16.2	91.1
30-34	1,256	13.6	1,162	14.0	92.6
35-39	1,154	12.5	1,049	12.6	90.9
40-44	975	10.5	857	10.3	87.9
45-49	787	8.5	708	8.5	90.0
50-54	684	-	-	-	-
15-49	9,262	-	8,319	-	89.8

Note: The de facto population includes all residents and nonresidents who slept in the household the night before interview.

Table D.3 Completeness of reporting

Percentage of observations missing information for selected demographic and health questions (weighted), Turkey 1998

Subject	Reference group	Percentage missing information	Number of cases
Birth date	Births in last 15 years		
Month only		9.4	10,368
Month and year		0.8	10,368
Age at death	Deaths to births in last 15 years	0.8	683
Age/date at first union ¹	Ever-married women	2.0	6,196
Respondent's education	All women	0.0	8,576
Child's size at birth	Births in last 59 months	6.1	2,362
Anthropometry²	Living children age 0-59 months		
Height missing		16.9	3,299
Weight missing		12.8	3,299
Height or weight missing		17.6	3,299
Diarrhoea in last 2 weeks	Living children age 0-59 months	1.3	3,299

¹ Both year and age missing

² Child not measured

Table D.4 Births by calendar years

Distribution of births by Western calendar years for living (L), dead (D), and all (T) children, according to reporting completeness, sex ratio at birth, and ratio of births by calendar year, Turkey 1998

Year	Number of births			Percentage with complete birth date ¹			Sex ratio at birth ²			Calendar ratio ³			Male			Female		
	L	D	T	L	D	T	L	D	T	L	D	T	L	D	T	L	D	T
94	532	17	549	100.0	100.0	100.0	105.0	161.9	106.4	NA	NA	NA	273	10	283	260	6	266
93	712	28	740	100.0	100.0	100.0	107.2	103.7	107.0	124.2	90.5	122.4	368	14	383	344	14	358
92	615	46	661	99.8	95.5	99.5	109.4	70.7	106.2	92.6	155.9	95.3	321	19	340	294	27	320
91	616	30	646	100.0	100.0	100.0	113.8	120.1	114.1	95.3	79.4	94.4	328	17	344	288	14	302
90	677	31	708	99.6	100.0	99.6	104.2	279.1	108.4	109.7	89.0	108.6	345	23	368	332	8	340
89	619	39	658	99.7	100.0	99.8	115.2	124.7	115.7	87.6	94.5	88.0	331	22	353	287	17	305
88	736	52	787	88.4	58.1	86.4	96.6	93.5	96.4	116.5	117.2	116.5	361	25	386	374	27	401
87	645	49	694	88.6	71.1	87.4	111.4	80.7	108.9	89.6	102.7	90.4	340	22	362	305	27	332
86	704	44	748	86.8	67.9	85.7	102.7	154.7	105.1	114.6	92.6	113.0	357	27	383	347	17	365
85	584	46	630	89.1	54.8	86.6	105.7	121.2	106.8	NA	NA	NA	300	25	325	284	21	305
90-94	3,152	152	3,304	99.9	98.7	99.8	107.8	120.3	108.4	NA	NA	NA	1,635	83	1,719	1,516	69	1,586
85-89	3,287	230	3,517	90.4	69.2	89.0	105.7	110.2	106.0	NA	NA	NA	1,689	121	1,810	1,598	109	1,707
80-84	3,118	284	3,401	84.8	63.3	83.0	104.8	121.4	106.1	NA	NA	NA	1,595	156	1,751	1,522	128	1,651
75-79	2,833	392	3,225	84.3	57.6	81.0	101.4	121.9	103.7	NA	NA	NA	1,426	216	1,642	1,406	177	1,583
< 75	3,067	694	3,761	80.7	54.3	75.8	99.1	128.3	103.9	NA	NA	NA	1,526	390	1,916	1,541	304	1,845
All	15,456	1,753	17,209	88.1	62.3	85.5	103.8	122.5	105.6	-	-	-	7,872	965	8,837	7,584	788	8,371

NA = Not applicable

¹ Both year and month of birth given

² $(B_m/B_f)*100$, where B_m and B_f are the numbers of male and female births, respectively

³ $[2B_x/(B_{x-1}+B_{x+1})]*100$, where B_x is the number of births in calendar year x

Table D.5 Reporting of age at death in days

Distribution of reported deaths under one month of age by age at death in days and the percentage of neonatal deaths reported to occur at ages 0-6 days, for five-year periods preceding the survey (unweighted), Turkey 1998

Age at death (in days)	Number of years preceding the survey				Total 0-19
	0-4	5-9	10-14	15-19	
0	19	27	22	26	94
1	20	20	27	14	80
2	14	15	4	10	42
3	12	7	13	15	47
4	7	2	5	1	15
5	2	3	0	3	8
6	2	0	1	3	6
7	2	11	6	16	35
8	1	0	1	0	1
9	0	0	2	1	3
10	0	2	6	5	14
11	0	0	0	2	2
12	0	3	1	1	5
13	1	1	1	1	3
15	4	5	6	9	25
16	1	1	0	0	2
17	0	0	3	3	5
18	0	0	1	5	6
19	0	1	0	0	1
20	1	3	3	2	9
22	1	0	1	0	3
23	0	1	1	1	4
25	0	1	3	4	8
27	0	0	1	3	4
28	0	1	1	0	1
30	1	3	2	4	10
Missing	0	2	0	0	2
Total 0-30 ¹	88	106	110	127	432
Percent early neonatal ²	85.6	69.4	64.5	56.1	67.6

¹ Includes cases for which age at death (in exact days) is not known
² (0-6 days/0-30 days) * 100

Table D.6 Reporting of age at death in months

Distribution of reported deaths under two years of age by age at death in months and the percentage of infant deaths reported to occur at ages under one month, for five-year periods preceding the survey (unweighted), Turkey 1998

Age at death (in months)	Number of years preceding the survey				Total 0-19
	0-4	5-9	10-14	15-19	
<1 ^a	88	108	110	127	434
1	4	13	18	27	62
2	7	17	18	22	64
3	3	14	19	28	65
4	7	4	14	23	47
5	7	6	11	11	35
6	4	7	13	21	46
7	4	3	4	13	24
8	7	4	4	11	26
9	5	4	7	11	27
10	1	1	4	1	7
11	2	3	1	9	15
12	4	12	9	16	41
13	1	0	0	2	2
14	1	1	0	0	3
15	0	1	0	0	1
16	0	0	0	1	1
18	0	6	5	5	16
19	0	0	0	1	1
20	1	0	0	0	1
22	0	0	0	1	1
24+	0	1	1	0	1
1 Year	4	3	5	3	16
Total 0-11 ^b	139	185	223	305	852
Percent neonatal ^c	63.5	58.6	49.6	41.6	51.0

^a Includes deaths under 1 month reported in days

^b Includes cases for which age at death (in exact months) is not known

^c (under 1 month/under 1 year) * 100

APPENDIX E

QUESTIONNAIRES

HACETTEPE UNIVERSITY INSTITUTE OF POPULATION STUDIES
1998 TURKISH DEMOGRAPHIC AND HEALTH SURVEY
HOUSEHOLD QUESTIONNAIRE

IDENTIFICATION	
CLUSTER NO.....	PROVINCE _____
HOUSEHOLD NO.....	DISTRICT _____
REGION.....	SUB-DISTRICT _____
URBAN (1) / RURAL (2).....	VILLAGE _____
	QUARTER _____
	STREET _____ NO _____

INTERVIEWER VISITS				
	1	2	3	FINAL VISIT
DATE: DAY-MONTH	_ _	_ _	_ _	_ _
NAME AND SURNAME OF INTERVIEWER	_____	_____	_____	_ _
RESULT (*)	_	_	_	_
NEXT VISIT DATE TIME	_ _	_ _		TOTAL NUMBER OF VISITS

<p>(*) RESULT CODES:</p> <p>1 COMPLETED</p> <p>2 HOUSEHOLD PRESENT BUT NO COMPETENT RESPONDENT AT HOME</p> <p>3 HOUSEHOLD ABSENT</p> <p>4 POSTPONED</p> <p>5 REFUSED</p> <p>6 DWELLING VACANT/ ADDRESS NOT A DWELLING</p> <p>7 DWELLING DESTROYED</p> <p>8 DWELLING NOT FOUND</p> <p>9 OTHER _____ (SPECIFY)</p>	<p>TOTAL IN HOUSEHOLD.....</p> <p>TOTAL ELIGIBLE EVER MARRIED WOMEN.....</p> <p>TOTAL ELIGIBLE SINGLE WOMEN.....</p> <p>SELECTED FOR HUSBANDS</p> <p>YES..... 1 → TOTAL ELIGIBLE HUSBANDS....</p> <p>NO..... 2</p>
--	--

<p>SUPERVISOR</p> <p style="text-align: center;">_ _</p> <p>DAY-MONTH</p> <p style="text-align: center;">_ _</p>	<p>FIELD EDITOR</p> <p style="text-align: center;">_ _</p> <p>DAY-MONTH</p> <p style="text-align: center;">_ _</p>	<p>KEYER</p> <p style="text-align: center;">_ _</p> <p>DAY-MONTH</p> <p style="text-align: center;">_ _</p>
--	--	---

HOUR MINUTE

SECTION I. Now I would like some information about people in this household, such as age and education.

HH LINE NO	HOUSEHOLD LIST ADD BY ASKING A-B-C-D	RELATIONSHIP TO HEAD OF HH	RESIDENCE		SEX	AGE	DATE OF BIRTH
			Does..... usually live in this house? YES..... 1 NO 2	Did sleep here last night? YES..... 1 NO 2			
(01)	(02)	(03)	(04)	(05)	(06)	(07)	(08)
01		01	1 2	1 2	1 2	<input type="text"/>	<input type="text"/>
02		<input type="text"/>	1 2	1 2	1 2	<input type="text"/>	<input type="text"/>
03		<input type="text"/>	1 2	1 2	1 2	<input type="text"/>	<input type="text"/>
04		<input type="text"/>	1 2	1 2	1 2	<input type="text"/>	<input type="text"/>
05		<input type="text"/>	1 2	1 2	1 2	<input type="text"/>	<input type="text"/>
06		<input type="text"/>	1 2	1 2	1 2	<input type="text"/>	<input type="text"/>
07		<input type="text"/>	1 2	1 2	1 2	<input type="text"/>	<input type="text"/>
08		<input type="text"/>	1 2	1 2	1 2	<input type="text"/>	<input type="text"/>
09		<input type="text"/>	1 2	1 2	1 2	<input type="text"/>	<input type="text"/>
10		<input type="text"/>	1 2	1 2	1 2	<input type="text"/>	<input type="text"/>

I want to be sure that I have completed the full list of everyone in this household.

1. Are there any other persons such as small children or infants? YES → ADD TO LIST
NO
2. Are there any others who are not members of your family but live here, such as lodgers, friends, servants? YES → ADD TO LIST
NO

IF THE HOUSEHOLD LIST COMPRISES MORE THAN 10 PERSONS, TICK HERE AND CONTINUE LISTING THE HOUSEHOLD ON A SEPARATE FORM. PROCEED WITH THE REST OF THE INTERVIEW ON THE ADDITIONAL FORM.

HH LINE NO	PLACE OF BIRTH	MATERNAL SURVIVAL		PATERNAL SURVIVAL		EDUCATION ASK IF AGED 6 AND OVER		
		Is 's natural mother alive?	RECORD LINE NO. IF LISTED IN THE HOUSE. RECORD "96" IF LIVING ELSEWHERE.	Is 's natural father alive?	RECORD LINE NO. IF LISTED IN THE HOUSE. RECORD "96" IF LIVING ELSEWHERE	Has ever been to school?	What is the highest level of school attended? What is the highest grade completed at that level?	Did graduate from this school? (Did receive diploma?)
		ALIVE 1 DEAD 2 DK 8 } (12)		ALIVE 1 DEAD 2 DK 8 } (14)		YES 1 NO 2 DK 8 } (18)	LEVEL/GRADE	YES 1 NO 2 DK 8
	(09)	(10)	(11)	(12)	(13)	(14)	(15)	(16)
01	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8
02	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8
03	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8
04	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8
05	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8
06	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8
07	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8
08	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8
09	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8
10	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8	<input type="text"/>	1 2 8

(*) CODES FOR RELATIONSHIP TO THE HOUSEHOLD HEAD:

01 HEAD	07 PARENT-IN-LAW	13 FIRST DEGREE COUSIN
02 WIFE/HUSBAND	08 SIBLING	14 GRAND PARENT
03 SON/DAUGHTER	09 SIBLING -IN- LAW	15 SIBLING'S CHILD
04 SON/DAUGHTER IN LAW	10 FATHER'S SIBLING	16 OTHER RELATIVE
05 GRANDCHILD	11 MOTHER'S SIBLING	17 NOT RELATED
06 PARENT	12 STEP CHILD	98 DK

LEVEL CODES

1	PRIMARY
2	SECONDARY
3	HIGH SCHOOL
4	UNIVERSITY
8	DK

CODES FOR GRADE

0	LESS THAN ONE YEAR/PREPARATORY
8	DK

HH LINE NO	ASK IF AGED LESS THAN 25	MARITAL STATUS			ELIGIBLE WOMAN		ELIGIBLE HUSBAND
		ASK IF AGED 12 AND OVER					
	Is still attending school? YES..... 1 NO 2 DK..... 8	Has ... ever married? YES..... 1 NO 2 → (21) DK..... 8 ↓ NEXT PERSON	What is 's marital status? MARRIED..... 1 WIDOWED..... 2 DIVORCED..... 3 → (22) SEPERATED..... 4 DK..... 8	RECORD LINE NO OF SPOUSE AND SKIP TO QST. 22 IF SPOUSE NOT IN THE HOUSEHOLD LIST, WRITE "96".	CIRCLE LINE NUMBER IF SINGLE WOMAN AGE 15-49 AND SKIP TO NEXT PERSON	CIRCLE LINE NUMBER IF EVER MARRIED WOMAN AGE 15-49 AND SKIP TO NEXT PERSON	AFTER COMPLETING THE HH LIST CIRCLE LINE NUMBER IF HE IS CURRENT HUSBAND OF A CURRENTLY MARRIED WOMAN CIRCLED IN QST 22
	(17)	(18)	(19)	(20)	(21)	(22)	(23)
01	1 2 8	1 2 8	1 2 3 4 8	<input type="text"/>	01	01	01
02	1 2 8	1 2 8	1 2 3 4 8	<input type="text"/>	02	02	02
03	1 2 8	1 2 8	1 2 3 4 8	<input type="text"/>	03	03	03
04	1 2 8	1 2 8	1 2 3 4 8	<input type="text"/>	04	04	04
05	1 2 8	1 2 8	1 2 3 4 8	<input type="text"/>	05	05	05
06	1 2 8	1 2 8	1 2 3 4 8	<input type="text"/>	06	06	06
07	1 2 8	1 2 8	1 2 3 4 8	<input type="text"/>	07	07	07
08	1 2 8	1 2 8	1 2 3 4 8	<input type="text"/>	08	08	08
09	1 2 8	1 2 8	1 2 3 4 8	<input type="text"/>	09	09	09
10	1 2 8	1 2 8	1 2 3 4 8	<input type="text"/>	10	10	10

SECTION 2. WELFARE OF ELDERLY

30	CHECK QUESTION 07: IN HOUSEHOLD LIST THERE IS AT LEAST ONE PERSON AGE 65 AND OVER <input type="checkbox"/>	ALL PERSONS ARE LESS THAN 65 <input type="checkbox"/>	→ 40
-----------	---	---	------

ENTER THE NAME AND LINE NUMBER OF EACH ELDERLY LISTED IN THE HOUSEHOLD SCHEDULE. ASK QUESTIONS ABOUT EACH OF THE ELDERLY SEPARATELY. BEGIN WITH THE ELDERLY HIGHEST ON THE HOUSEHOLD LIST. (IF THERE ARE MORE THAN 2 ELDERLY, USE ADDITIONAL QUESTIONNAIRES).

31	FROM QUESTION 02 FROM QUESTION 01	NAME _____ LINE NUMBER..... <input type="text"/>	NAME _____ LINE NUMBER..... <input type="text"/>	
-----------	--	---	---	--

32	Does have any living children? IF YES: How many? IF NO WRITE "00".	NO OF CHILDREN..... <input type="text"/>	NO OF CHILDREN..... <input type="text"/>	
-----------	--	--	--	--

33	Does have any living step children? IF YES: How many? IF NO WRITE "00".	NO OF STEP CHILDREN..... <input type="text"/>	NO OF STEP CHILDREN..... <input type="text"/>	
-----------	---	---	---	--

34		SKIP TO QST. 36. IF HAS NO LIVING CHILDREN OR STEP CHILDREN.	SKIP TO QST. 36. IF HAS NO LIVING CHILDREN OR STEP CHILDREN.	
-----------	--	--	--	--

35	Of 's own or step children, where does the nearest living child live? In the same house, very close, or another place?	SAME HOUSE.....1 NEXT DOOR/VERY CLOSE.....2 SAME COMMUNITY.....3 CLOSE COMMUNITY.....4 DISTANT COMMUNITY.....5 OTHER COUNTRY.....6 DK.....8	SAME HOUSE.....1 NEXT DOOR/VERY CLOSE.....2 SAME COMMUNITY.....3 CLOSE COMMUNITY.....4 DISTANT COMMUNITY.....5 OTHER COUNTRY.....6 DK.....8	
-----------	--	---	---	--

36	Who takes the prime responsibility for 's needs, health and welfare?	HIMSELF/HERSELF.....1 SPOUSE.....2 CHILDREN/STEP CHILDREN.....3 CHILDREN IN LAW.....4 CLOSE RELATIVE.....5 DISTANT RELATIVE.....6 NEIGHBOURS.....7 OTHER.....8 (SPECIFY)	HIMSELF/HERSELF.....1 SPOUSE.....2 CHILDREN/STEP CHILDREN.....3 CHILDREN IN LAW.....4 CLOSE RELATIVE.....5 DISTANT RELATIVE.....6 NEIGHBOURS.....7 OTHER.....8 (SPECIFY)	
-----------	--	--	--	--

37	Does have any income?	YES.....1 NO.....2 → 39	YES.....1 NO.....2 → 39	
-----------	-----------------------------	----------------------------	----------------------------	--

38	What are the source(s) of this income? RECORD ALL MENTIONED.	PENSION (SELF).....A PENSION (DIRECT).....B OLD AGE PENSION.....C RENT/INTEREST.....D FROM RELATIVE HERE.....E FROM RELATIVE ABROAD.....F CURRENTLY WORKING.....G OTHER.....I (SPECIFY)	PENSION (SELF).....A PENSION (DIRECT).....B OLD AGE PENSION.....C RENT/INTEREST.....D FROM RELATIVE HERE.....E FROM RELATIVE ABROAD.....F CURRENTLY WORKING.....G OTHER.....I (SPECIFY)	
-----------	---	---	---	--

39		RETURN TO QST. 32 IF THERE IS ANOTHER ELDERLY IN THE HOUSE; SKIP TO QST. 40 OTHERWISE.	RETURN TO QST. 32 IF THERE IS ANOTHER ELDERLY IN THE HOUSE; SKIP TO QST. 40 OTHERWISE.	
-----------	--	--	--	--

SECTION 3. HOUSING CHARACTERISTICS

<p>40</p>	<p>Does this house belong to a household member, is it rented from someone else, is it a lodging, or do you just live here without having to pay anything?</p>	<p>OWNED BY A HOUSEHOLD MEMBER 1 RENTED 2 LODGING 3 NO RENT PAID 4 OTHER 7 (SPECIFY)</p>	
<p>41</p>	<p>What is the source of drinking water for members of your household?</p>	<p>PIPED WATER PIPED WATER IN HOUSE/GARDEN 11 → 43 PUBLIC PIPED WATER OUTSIDE HOUSE/GARDEN 12 WELL WATER WELL IN HOUSE/GARDEN 21 → 43 PUBLIC WELL 22 SURFACE WATER PIPED SURFACE WATER IN HOUSE/GARDEN 31 → 43 SPRING/PUBLIC FOUNTAIN 32 RIVER/STREAM/POND/LAKE/DAM 33 RAINWATER 41 → 43 TANKER TRUCK 51 BOTTLED WATER/DEMI JOHN/PET WATER 61 → 43 WATER STATION 71 OTHER 96 (SPECIFY)</p>	
<p>42</p>	<p>How long does it take you to go there, get water, and come back?</p>	<p>MINUTE <input type="text" value=""/><input type="text" value=""/><input type="text" value=""/> ON PREMISES 996</p>	
<p>43</p>	<p>Is the toilet inside the house or outside?</p>	<p>NO FACILITY/BUSH/FIELD 0 → 45 INSIDE 1 OUTSIDE 2 INSIDE AND OUTSIDE 3 OTHER 7 (SPECIFY)</p>	
<p>44</p>	<p>What type of toilet system do you have in your household? PROBE IF THE TOILET SYSTEM IS CONNECTED TO DRAINAGE SYSTEM. IF CONNECTED: Is the toilet used by only the members of your household or is it shared with other households? IF NOT CONNECTED: What is the facility for excreta disposal?</p>	<p>FLUSH TOILET OWN FLUSH TOILET 11 SHARED FLUSH TOILET 12 PIT TOILET OPEN PIT 21 CLOSED PIT 22 NO FACILITY/ BUSH/ FIELD 31 OTHER 96 (SPECIFY)</p>	
<p>45</p>	<p>What is the main source of heating in winter for your house?</p>	<p>RADIATOR (CENTRAL HEATING) 1 RADIATOR (PRIVATE) 2 NATURAL GAS STOVE 3 STOVE (COAL, WOOD) 4 OTHER 7 (SPECIFY)</p>	

50	LINE NO. OF RESPONDENT TO THE HOUSEHOLD SCHEDULE	LINE NO..... <input type="text"/>	
51	LANGUAGE USED FOR CONDUCTING THE HOUSEHOLD QUESTIONNAIRE	TURKISH 1 → 53 KURDISH 2 ARABIC 3 OTHER 7 (SPECIFY)	
52	WAS AN INTERPRETER USED?	YES 1 NO 2	
53	RECORD THE TIME	HOUR <input type="text"/> MINUTE..... <input type="text"/>	

GO BACK TO THE FRONT COVER AND
COMPLETE THE NECESSARY INFORMATION.

PROVINCE CODES

01 ADANA	21 DIYARBAKIR	41 KOCAELI	61 TRABZON
02 ADIYAMAN	22 EDİRNE	42 KONYA	62 TUNCELI
03 AFYON	23 ELAZIĞ	43 KÜTAHYA	63 ŞANLIURFA
04 AĞRI	24 ERZINCAN	44 MALATYA	64 UŞAK
05 AMASYA	25 ERZURUM	45 MANISA	65 VAN
06 ANKARA	26 ESKİŞEHİR	46 K.MARAŞ	66 YOZGAT
07 ANTALYA	27 GAZİANTEP	47 MARDİN	67 ZONGULDAK
08 ARTVIN	28 GİRESUN	48 MUĞLA	68 AKSARAY
09 AYDIN	29 GÜMÜŞHANE	49 MUŞ	69 BAYBURT
10 BALIKESİR	30 HAKKARI	50 NEVŞEHİR	70 KARAMAN
11 BİLECİK	31 HATAY	51 NIĞDE	71 KIRIKKALE
12 BİNGÖL	32 İSPARTA	52 ORDU	72 BATMAN
13 BİTLİS	33 İÇEL	53 RİZE	73 ŞİRNAK
14 BOLU	34 İSTANBUL	54 SAKARYA	74 BARTIN
15 BURDUR	35 İZMİR	55 SAMSUN	75 ARDAHAN
16 BURSA	36 KARS	56 SİİRT	76 İĞDIR
17 ÇANAKKALE	37 KASTAMONU	57 SİNOP	77 YALOVA
18 ÇANKIRI	38 KAYSERİ	58 SİVAS	78 KARABÜK
19 ÇORUM	39 KIRKLARELİ	59 TEKİRDAĞ	79 KILIS
20 DENİZLİ	40 KİRŞEHİR	60 TOKAT	80 OSMANİYE

90 OTHER COUNTRY

**CONVERSION OF YEARS OF BIRTH FROM
RUMI CALENDAR TO MILADI CALENDAR
YEARS :**

RUMI YEAR + 584 = MILADI YEAR

HACETTEPE UNIVERSITY INSTITUTE OF POPULATION STUDIES
 1998 TURKISH DEMOGRAPHIC AND HEALTH SURVEY
 EVER-MARRIED WOMAN'S QUESTIONNAIRE

IDENTIFICATION			
CLUSTER NO.....	<input type="text"/>	<input type="text"/>	PROVINCE _____
HOUSEHOLD NO.....	<input type="text"/>	<input type="text"/>	DISTRICT _____
REGION.....	<input type="text"/>	<input type="text"/>	SUBDISTRICT _____
URBAN (1) / RURAL (2).....	<input type="text"/>	<input type="text"/>	VILLAGE _____
			QUARTER _____
			STREET _____ NO _____

NAME-SURNAME OF WOMAN _____	LINE NUMBER OF WOMAN.....	<input type="text"/>
-----------------------------	---------------------------	----------------------

INTERVIEWER VISITS				
	1	2	3	FINAL VISIT
DATE (DAY-MONTH)	___ ___	___ ___	___ ___	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>
INTERVIEWER'S NAME-SURNAME	_____	_____	_____	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>
RESULT (*)	___	___	___	<input type="text"/>
NEXT VISIT DAY-MONTH HOUR	___ ___ _____	___ ___ _____	_____	TOTAL NO OF VISITS <input type="text"/>

(*) RESULT CODES :

1 COMPLETED	5 PARTLY COMPLETED
2 NOT AT HOME	
3 POSTPONED	7 OTHER _____
4 REFUSED	(SPECIFY)

SUPERVISOR _____ <input type="text"/>	FIELD EDITOR _____ <input type="text"/>	KEYED BY _____ <input type="text"/>
DAY-MONTH <input type="text"/>	DAY-MONTH <input type="text"/>	DAY-MONTH <input type="text"/>

Age	Year of birth	
	Has not had birthday in 1998	Has already had birthday in 1998
	Does not know	
0	1997	--
1	1996	1997
2	1995	1996
3	1994	1995
4	1993	1994
5	1992	1993
6	1991	1992
7	1990	1991
8	1989	1990
9	1988	1989
10	1987	1988
11	1986	1987
12	1985	1986
13	1984	1985
14	1983	1984
15	1982	1983
16	1981	1982
17	1980	1981
18	1979	1980
19	1978	1979
20	1977	1978
21	1976	1977
22	1975	1976
23	1974	1975
24	1973	1974
25	1972	1973
26	1971	1972
27	1970	1971
28	1969	1970
29	1968	1969
30	1967	1968
31	1966	1967
32	1965	1966
33	1964	1965
34	1963	1964

Age	Year of birth	
	Has not had birthday in 1998	Has already had birthday in 1998
	Does not know	
35	1962	1963
36	1961	1962
37	1960	1961
38	1959	1960
39	1958	1959
40	1957	1958
41	1956	1957
42	1955	1956
43	1954	1955
44	1953	1954
45	1952	1953
46	1951	1952
47	1950	1951
48	1949	1950
49	1948	1949
50	1947	1948
51	1946	1947
52	1945	1946
53	1944	1945
54	1943	1944
55	1942	1943
56	1941	1942
57	1940	1941
58	1939	1940
59	1938	1939
60	1937	1938
61	1936	1937
62	1935	1936
63	1934	1935
64	1933	1934
65	1932	1933
66	1931	1932
67	1930	1931
68	1929	1930
69	1928	1929

SECTION 1. RESPONDENT'S BACKGROUND

101	RECORD THE TIME	HOUR - MINUTES..... <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>
102A	First I would like to ask some questions about you and the place you lived. For most of the time until you were 12 years old, where did you live ? _____ (NAME OF THE PLACE) Was it a province centre, district centre, a subdistrict or village ? Or did you live abroad ?	PROVINCE CENTRE.....1 DISTRICT CENTRE.....2 SUBDISTRICT OR VILLAGE.....3 ABROAD.....4 → 103
102B	In which province is this place now ? RECORD THE NAME AND CODE OF THE PROVINCE.	NAME OF PROVINCE <input type="text"/> _____ PROVINCE CODE
103	How long have you been living continuously in (NAME OF CURRENT PLACE OF INTERVIEW) ?	YEARS..... <input type="text"/> SINCE I WAS BORN.....95 VISITOR.....96 → 105
104A	Where did you live before you moved here ? _____ (NAME OF THE PLACE) Was that a province centre, district centre, a subdistrict or village ? Or did you live abroad ?	PROVINCE CENTRE.....1 DISTRICT CENTRE.....2 SUBDISTRICT OR VILLAGE.....3 ABROAD.....4 → 104C
104B	In which province is this place now ? RECORD THE NAME AND CODE OF THE PROVINCE.	NAME OF PROVINCE <input type="text"/> _____ PROVINCE CODE
104C	What was the main reason for you to move here ?	PERSONAL REASONS MARRIAGE.....01 EDUCATION.....02 LOOKING FOR JOB.....03 CHANGE OF JOB/APPOINTMENT.....04 RETURN TO HOMELAND.....05 SPOUSAL REASONS TO ACCOMPANY HUSBAND.....06 CHANGE OF JOB/APPOINTMENT..07 LOOKING FOR JOB.....08 FAMILIAL REASONS TO JOIN PARENTS.....09 CHANGE OF JOB/APPOINTMENT..10 LOOKING FOR JOB.....11 OTHER _____ 96 (SPECIFY)

105	In what month and year were you born ?	MONTH..... <input type="text"/> <input type="text"/> DON'T KNOW MONTH.....98 YEAR..... <input type="text"/> 9 <input type="text"/> <input type="text"/> DON'T KNOW YEAR.....9998
106	How old are you exactly ? What age have you completed ? CHECK ANSWERS TO 105 AND 106 USING AGE-BIRTH YEAR TABLE. IF INCONSISTENT PROBE AND CORRECT. AGE INFORMATION MUST BE RECORDED !	AGE IN COMPLETED YEARS..... <input type="text"/> <input type="text"/>
107	Have you ever attended school ?	YES.....1 NO.....2 →114
108	What is the highest level you have attended ?	PRIMARY.....1 SECONDARY.....2 HIGH SCHOOL.....3 UNIVERSITY.....4
109A	What is the highest grade you have completed at that level ?	GRADE..... <input type="text"/>
109B	Did you graduate (receive diploma) from this school ?	YES.....1 NO.....2
110	CHECK 106 : AGE 24 OR BELOW <input type="checkbox"/> AGE 25 OR ABOVE <input type="checkbox"/> →113	
111	Are you currently attending school ?	YES.....1 →113 NO.....2

112	What was the main reason you stopped attending school ?	GOT PREGNANT.....01 GOT MARRIED.....02 TO CARE FOR YOUNGER CHILDREN...03 FAMILY NEEDED HELP ON FARM OR IN BUSINESS.....04 COULD NOT PAY SCHOOL FEES.....05 NEEDED TO EARN MONEY.....06 GRADUATED/HAD ENOUGH SCHOOLING.07 DID NOT PASS ENTRANCE EXAMS...08 DID NOT LIKE SCHOOL.....09 SCHOOL NOT ACCESSIBLE/TOO FAR..10 PARENTS DID NOT SEND.....11 OTHER _____ 96 (SPECIFY) DON'T KNOW.....98
-----	---	---

113	CHECK 108 : ATTENDED <input type="checkbox"/> PRIMARY	ATTENDED <input type="checkbox"/> SECONDARY OR HIGHER	→115
-----	---	---	------

114	Can you read and understand a letter or newspaper easily, with difficulty, or not at all ?	EASILY.....1 WITH DIFFICULTY.....2 NOT AT ALL.....3 →116A
-----	--	---

115	How frequent do you read a newspaper or magazine ?	EVERY DAY/ALMOST EVERY DAY.....1 ONCE-TWICE A WEEK.....2 RARELY/SELDOM.....3 NEVER/ALMOST NEVER.....4
-----	--	--

116A	What is your mother tongue ? RECORD ONLY ONE RESPONSE.	TURKISH.....01 KURDISH AND DIALECTS (KURMANCI, GORANI, ZAZA ETC)...02 ARABIC.....03 GREEK.....04 ARMANIAN.....05 HEBREW(LADINO).....06 CIRCASSIAN.....07 GEORGIAN.....08 LAZ LANGUAGE.....09 PERSIAN.....10 BULGARIAN.....11 RUMANIAN.....12 SERBIAN.....13 ENGLISH.....14 GERMAN.....15 OTHER _____ 96 (SPECIFY)
------	---	--

116B	<p>In addition to your mother tongue, which language(s) can you speak ?</p> <p style="text-align: center;">RECORD ALL MENTIONED.</p>	<p>TURKISH.....A KURDISH AND DIALECTS (KURMANCI, GORANI, ZAZA ETC)....B ARABIC.....C GREEK.....D ARMANIAN.....E HEBREW (LADINO).....F CIRCASSIAN.....G GEORGIAN.....H LAZ LANGUAGE.....I PERSIAN.....J BULGARIAN.....K RUMANIAN.....L SERBIAN.....M ENGLISH.....N GERMAN.....O OTHER _____ U (SPECIFY) KNOWS NO OTHER LANGUAGE.....Y</p>
------	--	--

116C	<p>What is (was) your mother's and father's mother tongue ?</p> <p style="text-align: center;">USE THE CODES IN 116A.</p>	<p>MOTHER.....</p> <p>FATHER.....</p> <div style="display: flex; align-items: center; justify-content: center;"> <table border="1" style="border-collapse: collapse; text-align: center;"> <tr><td style="width: 20px; height: 20px;"></td><td style="width: 20px; height: 20px;"></td></tr> <tr><td style="width: 20px; height: 20px;"></td><td style="width: 20px; height: 20px;"></td></tr> </table> </div>				

117	<p>What is your religion ?</p> <p style="text-align: center;">IF THE ANSWER IS "MUSLIM" PROBE FOR RELIGIOUS SECT AND CIRCLE APPROPRIATE CODE.</p>	<p>MUSLIM</p> <p>SUNNI.....01 ALAWI.....02 OTHER _____ 03 (SPECIFY) CHRISTIAN.....06 JEWISH.....07 NO RELIGION.....10 OTHER _____ 96 (SPECIFY)</p>
-----	---	--

118	CHECK Q.4 IN THE HOUSEHOLD QUESTIONNAIRE :	
<p>THE WOMAN INTERVIEWED IS NOT A USUAL RESIDENT <input type="checkbox"/></p>	<p>THE WOMAN INTERVIEWED IS A USUAL RESIDENT <input type="checkbox"/></p>	→201

119A	<p>Now I would like to ask about the place in which you usually live.</p> <p>What is the name of the place in which you usually live ?</p> <p style="text-align: center;">_____</p> <p style="text-align: center;">(NAME OF PLACE)</p> <p>Is that a province centre, a district centre, a subdistrict or village, or are you living abroad ?</p>	<p>PROVINCE CENTRE.....1 DISTRICT CENTRE.....2 SUBDISTRICT OR VILLAGE.....3 ABROAD.....4 →120A</p>
------	--	---

119B	In which province is that located ?	NAME OF PROVINCE <input type="text"/> PROVINCE CODE <input type="text"/>
------	-------------------------------------	---

120A	Now I would like to ask about the household you usually live. How many persons do usually live in your house ?	NUMBER..... <input type="text"/>
------	---	----------------------------------

120B	Does the house you usually live belong to a household member, is it rented from someone else, is it a lodging, or do you just live here without having to pay anything ?	OWNED BY A HOUSEHOLD MEMBER.....1 RENTED.....2 LODGING.....3 NO MONEY PAID.....4 OTHER _____ 7 (SPECIFY)
------	--	---

121A	What is the source of drinking water for your household ?	PIPED WATER PIPED WATER IN HOUSE/GARDEN..11 →122A PUBLIC PIPED WATER OUTSIDE HOUSE/GARDEN.....12 WELL WATER WELL IN RESIDENCE/GARDEN....21 →122A PUBLIC WELL.....22 SURFACE WATER SPRING WATER PIPED INTO HOUSE/GARDEN.....31 →122A SPRING/PUBLIC FOUNTAIN.....32 RIVER/STREAM/ POND/LAKE/DAM.....33 RAINWATER.....41 →122A TANKER TRUCK.....51 BOTTLED WATER/DEMIJOHN/PET W...61 →122A WATER STATION.....71 OTHER _____ 96 (SPECIFY)
------	---	--

121B	How long does it take you to go there, get water, and come back ?	MINUTES..... <input type="text"/> ON PREMISES.....996
------	---	--

122A	Is the toilet inside the house or outside ?	NO FACILITY/BUSH/FIELD.....0 →123 INSIDE.....1 OUTSIDE.....2 INSIDE AND OUTSIDE.....3 OTHER _____ 7 (SPECIFY)
------	---	--

122B	<p>What type of toilet system do you have in your household?</p> <p>PROBE IF THE TOILET SYSTEM IS CONNECTED TO DRAINAGE SYSTEM.</p> <p>IF CONNECTED: Is toilet used by only the members of your household or is it shared with other households?</p> <p>IF NOT CONNECTED: What is the facility for excreta disposal?</p>	<p>CONNECTED TO DRAINAGE SYSTEM USED ONLY BY THIS HOUSEHOLD...11 SHARED WITH ANOTHER HOUSEHOLD...12</p> <p>PIT OPEN PIT.....21 CLOSED PIT.....22</p> <p>NO FACILITY.....31</p> <p>OTHER _____ 96 (SPECIFY)</p>
123	<p>What is the source of heating in winter for your house ?</p>	<p>RADIATOR (CENTRAL HEATING).....1 RADIATOR (PRIVATE).....2 NATURAL GAS STOVE.....3 STOVE (COAL/WOOD).....4</p> <p>OTHER _____ 7 (SPECIFY)</p>
124	<p>How many rooms in your household are used for sleeping ?</p>	<p>ROOMS USED FOR SLEEPING.... <input type="text"/></p>
125	<p>What is the main material of the floor ?</p>	<p>NATURAL FLOOR EARTH.....11 RUDIMENTARY WOOD PLANKS.....21 FINISHED FLOOR PARQUET/POLISHED WOOD.....31 KARO.....32 CEMENT.....34 CARPET/WALL TO WALL CARPET...35 MARLEY.....36 MOSAIC.....37</p> <p>OTHER _____ 96 (SPECIFY)</p>
126	<p>I would like to get an estimate of the total income that enters your household each month. Is the total amount of money earned by the members of your household in a month:</p>	
126A	<p>More than 100 million ?</p>	<p>YES.....1 NO.....2 → 126D</p>
126B	<p>More than 300 million ?</p>	<p>YES.....1 NO.....2 → 127</p>
126C	<p>More than 500 million ?</p>	<p>YES.....1 NO.....2 → 127</p>
126D	<p>Less than 50 million ?</p>	<p>YES.....1 NO.....2</p>

127

Do you have the following in your household ?

YES NO

Refrigerator	REFRIGERATOR.....1	2
Gas or electric oven	GAS/ELECTRIC OVEN.....1	2
Dishwasher	DISHWASHER.....1	2
Washing machine	WASHING MACHINE.....1	2
Vacuum cleaner	VACUUM CLEANER.....1	2
Television	TELEVISION.....1	2
Video	VIDEO.....1	2
Camera	CAMERA.....1	2
Music set with CD player	MUSIC SET WITH CD PLAYER...1	2
Telephone	TELEPHONE.....1	2
Cellular Telephone	CELLULAR TELEPHONE.....1	2
Car (excluding tractors, taxis, etc.)	CAR.....1	2
Computer	COMPUTER.....1	2

SECTION 2A. REPRODUCTION

201	Now I would like to ask about all the births you have had during your life. Have you ever given birth ?	YES.....1 NO.....2 → 206				
202	Do you have any sons or daughters to whom you have given birth who are now living with you ?	YES.....1 NO.....2 → 204				
203	How many sons live with you ? And how many daughters live with you ? IF NONE, RECORD "00".	SONS AT HOME..... DAUGHTERS AT HOME..... <table border="1" style="display: inline-table; vertical-align: middle; margin-left: 20px;"> <tr><td style="width: 20px; height: 20px;"></td><td style="width: 20px; height: 20px;"></td></tr> <tr><td style="width: 20px; height: 20px;"></td><td style="width: 20px; height: 20px;"></td></tr> </table>				
204	Do you have any sons or daughters to whom you have given birth who are alive but do not live with you ?	YES.....1 NO.....2 → 206				
205	How many sons are alive but do not live with you ? And how many daughters are alive but do not live with you ? IF NONE, RECORD "00".	SONS ELSEWHERE..... DAUGHTERS ELSEWHERE..... <table border="1" style="display: inline-table; vertical-align: middle; margin-left: 20px;"> <tr><td style="width: 20px; height: 20px;"></td><td style="width: 20px; height: 20px;"></td></tr> <tr><td style="width: 20px; height: 20px;"></td><td style="width: 20px; height: 20px;"></td></tr> </table>				
206	Have you ever given birth to a boy or a girl who was born alive but later died ? IF NO, PROBE: Any baby who cried or showed signs of life but only survived a few hours or days ?	YES.....1 NO.....2 → 208				
207	In all, how many boys have died ? And how many girls have died ? IF NONE, RECORD "00".	BOYS DEAD..... GIRLS DEAD..... <table border="1" style="display: inline-table; vertical-align: middle; margin-left: 20px;"> <tr><td style="width: 20px; height: 20px;"></td><td style="width: 20px; height: 20px;"></td></tr> <tr><td style="width: 20px; height: 20px;"></td><td style="width: 20px; height: 20px;"></td></tr> </table>				
208	SUM ANSWERS TO 203, 205, AND 207, AND ENTER TOTAL. IF NONE, RECORD "00".	TOTAL..... <table border="1" style="display: inline-table; vertical-align: middle; margin-left: 20px;"> <tr><td style="width: 20px; height: 20px;"></td><td style="width: 20px; height: 20px;"></td></tr> </table>				
209	CHECK 208 : Just to make sure that I have this right: you have had in TOTAL ____ live births during your life. Is that correct ? YES <input type="checkbox"/> NO <input type="checkbox"/> PROBE AND CORRECT 201-208 AS NECESSARY.					
210	CHECK 208 : ONE OR MORE LIVE BIRTHS <input type="checkbox"/> NO LIVE BIRTHS <input type="checkbox"/> → 227					

CONTINUE WITH THE BIRTH HISTORY (Q. 211)

BIRTH HISTORY

211 Now I would like to talk to you about all of your births. It is very important to learn about all of your births, whether still alive or not. Please let's start with the first one you had.
 RECORD NAMES OF ALL THE BIRTHS IN 212. RECORD TWINS AND TRIPLETS ON SEPARATE LINES.
 MAKE SURE TO RECORD DECEASED CHILDREN FROM MULTIPLE BIRTHS BEFORE THOSE SURVIVING.

212 What name was given to your (first/next) baby ? WRITE "BABY" IF THE BABY DIED BEFORE A NAME GIVEN.	213 RECORD SINGLE OR MULTIPLE BIRTH STATUS	214 Is (NAME) a boy or a girl ?	215 In what month and year was (NAME) born ? What is his/her birthday ? PROBE : In what season was he/she born ? NOTE: THE YEAR OF BIRTH FOR ALL CHILDREN, THE MONTH AND YEAR OF CHILDREN BORN AFTER 1993 HAVE TO BE DETERMINED !	216 Is (NAME) still alive ?						
01 <hr/> (NAME)	SINGLE.....1 MULTIPLE....2	BOY.....1 GIRL.....2	MONTH..... <table border="1" style="display: inline-table; vertical-align: middle;"><tr><td> </td><td> </td></tr></table> YEAR..... <table border="1" style="display: inline-table; vertical-align: middle;"><tr><td>1</td><td>9</td><td> </td><td> </td></tr></table>			1	9			YES.....1 NO.....2 219 ←
1	9									
02 <hr/> (NAME)	SINGLE.....1 MULTIPLE....2	BOY.....1 GIRL.....2	MONTH..... <table border="1" style="display: inline-table; vertical-align: middle;"><tr><td> </td><td> </td></tr></table> YEAR..... <table border="1" style="display: inline-table; vertical-align: middle;"><tr><td>1</td><td>9</td><td> </td><td> </td></tr></table>			1	9			YES.....1 NO.....2 219 ←
1	9									
03 <hr/> (NAME)	SINGLE.....1 MULTIPLE....2	BOY.....1 GIRL.....2	MONTH..... <table border="1" style="display: inline-table; vertical-align: middle;"><tr><td> </td><td> </td></tr></table> YEAR..... <table border="1" style="display: inline-table; vertical-align: middle;"><tr><td>1</td><td>9</td><td> </td><td> </td></tr></table>			1	9			YES.....1 NO.....2 219 ←
1	9									
04 <hr/> (NAME)	SINGLE.....1 MULTIPLE....2	BOY.....1 GIRL.....2	MONTH..... <table border="1" style="display: inline-table; vertical-align: middle;"><tr><td> </td><td> </td></tr></table> YEAR..... <table border="1" style="display: inline-table; vertical-align: middle;"><tr><td>1</td><td>9</td><td> </td><td> </td></tr></table>			1	9			YES.....1 NO.....2 219 ←
1	9									
05 <hr/> (NAME)	SINGLE.....1 MULTIPLE....2	BOY.....1 GIRL.....2	MONTH..... <table border="1" style="display: inline-table; vertical-align: middle;"><tr><td> </td><td> </td></tr></table> YEAR..... <table border="1" style="display: inline-table; vertical-align: middle;"><tr><td>1</td><td>9</td><td> </td><td> </td></tr></table>			1	9			YES.....1 NO.....2 219 ←
1	9									

<p>217 IF ALIVE: How old was (NAME) at his/her last birthday ? RECORD AGE IN COMPLETED YEARS. MAKE CALCULATIONS FOR CONSISTENCY.</p>	<p>218 IF ALIVE: Is (NAME) living with you ?</p>	<p>219 IF DEAD: How old was (NAME) when he/she died ? IF "1 YR.", PROBE: How many months old was (NAME)? RECORD DAYS IF LESS THAN 1 MONTH; MONTHS IF LESS THAN TWO YEARS; OR YEARS OTHERWISE.</p>	<p>220 FROM YEAR OF BIRTH OF (NAME) SUBTRACT YEAR OF PREVIOUS BIRTH. IS THE DIFFERENCE 4 OR MORE YEARS ?</p>	<p>221 Were there any other live births between (NAME OF PREVIOUS BIRTH) and (NAME) ?</p>
--	--	---	--	---

<p>AGE IN YEARS</p> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 5px auto;"></div>	<p>YES.....1 NO.....2 NEXT BIRTH</p>	<p>DAYS.....1 MONTHS.....2 YEARS.....3</p>	<table border="1" style="width: 100%; height: 100%; text-align: center;"> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> </table>								

<p>AGE IN YEARS</p> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 5px auto;"></div>	<p>YES.....1 NO.....2 220</p>	<p>DAYS.....1 MONTHS.....2 YEARS.....3</p>	<table border="1" style="width: 100%; height: 100%; text-align: center;"> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> </table>							<p>YES.....1 NO.....2 NEXT BIRTH</p>	<p>YES.....1 NO.....2</p>

<p>AGE IN YEARS</p> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 5px auto;"></div>	<p>YES.....1 NO.....2 220</p>	<p>DAYS.....1 MONTHS.....2 YEARS.....3</p>	<table border="1" style="width: 100%; height: 100%; text-align: center;"> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> </table>							<p>YES.....1 NO.....2 NEXT BIRTH</p>	<p>YES.....1 NO.....2</p>

<p>AGE IN YEARS</p> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 5px auto;"></div>	<p>YES.....1 NO.....2 220</p>	<p>DAYS.....1 MONTHS.....2 YEARS.....3</p>	<table border="1" style="width: 100%; height: 100%; text-align: center;"> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> </table>							<p>YES.....1 NO.....2 NEXT BIRTH</p>	<p>YES.....1 NO.....2</p>

<p>AGE IN YEARS</p> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 5px auto;"></div>	<p>YES.....1 NO.....2 220</p>	<p>DAYS.....1 MONTHS.....2 YEARS.....3</p>	<table border="1" style="width: 100%; height: 100%; text-align: center;"> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> </table>							<p>YES.....1 NO.....2 NEXT BIRTH</p>	<p>YES.....1 NO.....2</p>

IF THE ANSWER IS YES GO BACK AND MAKE THE NECESSARY CHANGES.

<p>212 What name was given to your (first/next) baby ?</p> <p>WRITE "BABY" IF THE BABY DIED BEFORE A NAME GIVEN.</p>	<p>213 RECORD SINGLE OR MULTIPLE BIRTH STATUS</p>	<p>214 Is (NAME) a boy or a girl ?</p>	<p>215 In what month and year was (NAME) born ? What is his/her birthday ?</p> <p>PROBE : In what season was he/she born ?</p> <p>NOTE: THE YEAR OF BIRTH FOR ALL CHILDREN, THE MONTH AND YEAR OF CHILDREN BORN AFTER 1993 HAVE TO BE DETERMINED !</p>	<p>216 Is (NAME) still alive ?</p>
--	---	--	--	------------------------------------

<p>06</p> <p>_____ (NAME)</p>	<p>SINGLE.....1</p> <p>MULTIPLE....2</p>	<p>BOY.....1</p> <p>GIRL.....2</p>	<p>MONTH.....</p> <p>YEAR..... 1 9</p>	<p>YES.....1</p> <p>NO.....2</p> <p>219 ←</p>
-------------------------------	--	------------------------------------	--	---

<p>07</p> <p>_____ (NAME)</p>	<p>SINGLE.....1</p> <p>MULTIPLE....2</p>	<p>BOY.....1</p> <p>GIRL.....2</p>	<p>MONTH.....</p> <p>YEAR..... 1 9</p>	<p>YES.....1</p> <p>NO.....2</p> <p>219 ←</p>
-------------------------------	--	------------------------------------	--	---

<p>08</p> <p>_____ (NAME)</p>	<p>SINGLE.....1</p> <p>MULTIPLE....2</p>	<p>BOY.....1</p> <p>GIRL.....2</p>	<p>MONTH.....</p> <p>YEAR..... 1 9</p>	<p>YES.....1</p> <p>NO.....2</p> <p>219 ←</p>
-------------------------------	--	------------------------------------	--	---

<p>09</p> <p>_____ (NAME)</p>	<p>SINGLE.....1</p> <p>MULTIPLE....2</p>	<p>BOY.....1</p> <p>GIRL.....2</p>	<p>MONTH.....</p> <p>YEAR..... 1 9</p>	<p>YES.....1</p> <p>NO.....2</p> <p>219 ←</p>
-------------------------------	--	------------------------------------	--	---

<p>10</p> <p>_____ (NAME)</p>	<p>SINGLE.....1</p> <p>MULTIPLE....2</p>	<p>BOY.....1</p> <p>GIRL.....2</p>	<p>MONTH.....</p> <p>YEAR..... 1 9</p>	<p>YES.....1</p> <p>NO.....2</p> <p>219 ←</p>
-------------------------------	--	------------------------------------	--	---

TICK HERE IF NUMBER OF LIVE BIRTHS IS MORE THAN 10 AND CONTINUE IN ANOTHER QUESTIONNAIRE FORM.

<p>217 IF ALIVE: How old was (NAME) at his/her last birthday? RECORD AGE IN COMPLETED YEARS. MAKE CALCULATIONS FOR CONSISTENCY.</p>	<p>218 IF ALIVE: Is (NAME) living with you?</p>	<p>219 IF DEAD: How old was (NAME) when he/she died? IF "1 YR.", PROBE: How many months old was (NAME)? RECORD DAYS IF LESS THAN 1 MONTH; MONTHS IF LESS THAN TWO YEARS; OR YEARS OTHERWISE.</p>	<p>220 FROM YEAR OF BIRTH OF (NAME) SUBTRACT YEAR OF PREVIOUS BIRTH. IS THE DIFFERENCE 4 OR MORE YEARS?</p>	<p>221 Were there any other live births between (NAME OF PREVIOUS BIRTH) and (NAME)?</p>
---	---	--	---	--

<p>AGE IN YEARS</p> <div style="border: 1px solid black; width: 30px; height: 20px; margin: 0 auto;"></div>	<p>YES.....1 NO.....2 ↓ 220</p>	<p>DAYS.....1 MONTHS.....2 YEARS.....3</p>	<table border="1" style="width: 100%; height: 100%; text-align: center;"> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> </table>									<p>YES.....1 NO.....2 NEXT BIRTH ←</p>	<p>YES.....1 NO.....2</p>

<p>AGE IN YEARS</p> <div style="border: 1px solid black; width: 30px; height: 20px; margin: 0 auto;"></div>	<p>YES.....1 NO.....2 ↓ 220</p>	<p>DAYS.....1 MONTHS.....2 YEARS.....3</p>	<table border="1" style="width: 100%; height: 100%; text-align: center;"> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> </table>									<p>YES.....1 NO.....2 NEXT BIRTH ←</p>	<p>YES.....1 NO.....2</p>

<p>AGE IN YEARS</p> <div style="border: 1px solid black; width: 30px; height: 20px; margin: 0 auto;"></div>	<p>YES.....1 NO.....2 ↓ 220</p>	<p>DAYS.....1 MONTHS.....2 YEARS.....3</p>	<table border="1" style="width: 100%; height: 100%; text-align: center;"> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> </table>									<p>YES.....1 NO.....2 NEXT BIRTH ←</p>	<p>YES.....1 NO.....2</p>

<p>AGE IN YEARS</p> <div style="border: 1px solid black; width: 30px; height: 20px; margin: 0 auto;"></div>	<p>YES.....1 NO.....2 ↓ 220</p>	<p>DAYS.....1 MONTHS.....2 YEARS.....3</p>	<table border="1" style="width: 100%; height: 100%; text-align: center;"> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> </table>									<p>YES.....1 NO.....2 NEXT BIRTH ←</p>	<p>YES.....1 NO.....2</p>

<p>AGE IN YEARS</p> <div style="border: 1px solid black; width: 30px; height: 20px; margin: 0 auto;"></div>	<p>YES.....1 NO.....2 ↓ 220</p>	<p>DAYS.....1 MONTHS.....2 YEARS.....3</p>	<table border="1" style="width: 100%; height: 100%; text-align: center;"> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> <tr><td> </td><td> </td></tr> </table>									<p>YES.....1 NO.....2 NEXT BIRTH ←</p>	<p>YES.....1 NO.....2</p>

IF THE ANSWER IS YES GO BACK AND MAKE THE NECESSARY CHANGES.

222	FROM 1998 SUBTRACT YEAR OF LAST BIRTH. IS THE DIFFERENCE 4 YEARS OR MORE ?	YES.....1 NO.....2 → 224			
223A	Have you had any live births since the birth of (NAME OF LAST BIRTH) ?	YES.....1 NO.....2 → 224			
223B	GO BACK AND MAKE THE NECESSARY CORRECTIONS.				
224	COMPARE 208 WITH NUMBER OF BIRTHS IN BIRTH HISTORY ABOVE: <table style="width: 100%; border: none;"> <tr> <td style="width: 30%; border: none;"> NUMBERS ARE SAME <input type="checkbox"/> </td> <td style="width: 30%; border: none;"> NUMBERS ARE DIFFERENT <input type="checkbox"/> </td> <td style="width: 40%; border: none;"> <input type="checkbox"/> → (PROBE, RECONCILE, AND MAKE NECESSARY CORRECTIONS) </td> </tr> </table> <p style="margin-left: 20px;">↓</p> <p>CHECK AND TICK:</p> <p>FOR EACH BIRTH: YEAR OF BIRTH (215) IS RECORDED.....</p> <p>FOR EACH BIRTH AFTER 1993: MONTH OF BIRTH (215) IS RECORDED.....</p> <p>FOR EACH LIVING CHILD: CURRENT AGE (217) IS RECORDED.....</p> <p>(IF ANY) FOR EACH DEAD CHILD:</p> <p> AGE AT DEATH (219) IS RECORDED.....</p> <p> FOR AGE AT DEATH 12 MONTHS OR 1 YR.: PROBED TO DETERMINE EXACT NUMBER OF MONTHS (219)</p>		NUMBERS ARE SAME <input type="checkbox"/>	NUMBERS ARE DIFFERENT <input type="checkbox"/>	<input type="checkbox"/> → (PROBE, RECONCILE, AND MAKE NECESSARY CORRECTIONS)
NUMBERS ARE SAME <input type="checkbox"/>	NUMBERS ARE DIFFERENT <input type="checkbox"/>	<input type="checkbox"/> → (PROBE, RECONCILE, AND MAKE NECESSARY CORRECTIONS)			
225	CHECK 215 AND ENTER THE NUMBER OF BIRTHS SINCE JANUARY 1993. IF NONE, RECORD '0'.	<input type="text"/>			
226	<table style="border: none;"> <tr> <td style="font-size: 2em; font-weight: bold; padding-right: 10px;">C</td> <td>FOR EACH BIRTH SINCE JANUARY 1993 ENTER 'D' IN THE MONTH OF BIRTH IN COLUMN 1 OF THE CALENDAR AND 'H' IN EACH OF THE 8 PRECEDING MONTHS. WRITE NAME OF CHILD TO THE LEFT OF THE 'D' CODE.</td> </tr> </table>		C	FOR EACH BIRTH SINCE JANUARY 1993 ENTER 'D' IN THE MONTH OF BIRTH IN COLUMN 1 OF THE CALENDAR AND 'H' IN EACH OF THE 8 PRECEDING MONTHS. WRITE NAME OF CHILD TO THE LEFT OF THE 'D' CODE.	
C	FOR EACH BIRTH SINCE JANUARY 1993 ENTER 'D' IN THE MONTH OF BIRTH IN COLUMN 1 OF THE CALENDAR AND 'H' IN EACH OF THE 8 PRECEDING MONTHS. WRITE NAME OF CHILD TO THE LEFT OF THE 'D' CODE.				
227	Are you pregnant now ?	YES.....1 NO.....2 UNSURE.....8 → 230A			
228	How many months pregnant are you ? <table style="border: none;"> <tr> <td style="font-size: 2em; font-weight: bold; padding-right: 10px;">C</td> <td>RECORD NUMBER OF COMPLETED MONTHS. ENTER 'H's IN COLUMN 1 OF THE CALENDAR, BEGINNING WITH THE MONTH OF INTERVIEW AND FOR TOTAL NUMBER OF COMPLETED MONTHS.</td> </tr> </table>	C	RECORD NUMBER OF COMPLETED MONTHS. ENTER 'H's IN COLUMN 1 OF THE CALENDAR, BEGINNING WITH THE MONTH OF INTERVIEW AND FOR TOTAL NUMBER OF COMPLETED MONTHS.	MONTHS..... <input style="width: 40px;" type="text"/>	
C	RECORD NUMBER OF COMPLETED MONTHS. ENTER 'H's IN COLUMN 1 OF THE CALENDAR, BEGINNING WITH THE MONTH OF INTERVIEW AND FOR TOTAL NUMBER OF COMPLETED MONTHS.				

229	At the time you became pregnant, did you want to become pregnant then, did you want to wait until later, or did you not want to have any more children at all ?	THEN.....1 LATER.....2 NOT WANT MORE CHILDREN.....3
230A	Have you ever had a pregnancy that ended in a miscarriage ?	YES.....1 NO.....2 → 230C
230B	In all, how many miscarriages have you had ?	NUMBER OF MISCARRIAGES..... <input style="width: 20px; height: 20px;" type="text"/>
230C	Have you ever had a pregnancy that ended in an induced abortion ?	YES.....1 NO.....2 → 230E
230D	In all, how many induced abortions have you had ?	NO. OF INDUCED ABORTIONS... <input style="width: 20px; height: 20px;" type="text"/>
230E	Have you ever had a pregnancy that ended in a still birth ?	YES.....1 NO.....2 → 230G
230F	In all, how many still births have you had ?	NUMBER OF STILL BIRTHS..... <input style="width: 20px; height: 20px;" type="text"/>
230G	CALCULATE THE TOTAL NUMBER OF PREGNANCIES. TOTAL NUMBER OF PREGNANCIES ENDING IN MISCARRIAGES, INDUCED ABORTIONS, OR STILL BIRTHS: SUM THE ANSWERS TO 230B, 230D, AND 230F. _____ TOTAL NUMBER OF PREGNANCIES ENDING IN LIVE BIRTHS: SUM THE NUMBER OF SINGLE BIRTHS IN THE BIRTH HISTORY. + _____ ADD TO THAT SUM THE NUMBER OF MULTIPLE BIRTHS. + _____ TOTAL NUMBER OF COMPLETED PREGNANCIES: = _____	TOTAL NUMBER OF COMPLETED PREGNANCIES..... <input style="width: 20px; height: 20px;" type="text"/>

230H	<p>CHECK 230G:</p> <p>Just to make sure that I have this right. You have had in TOTAL _____ completed pregnancies. Is that correct ?</p> <p>YES <input type="checkbox"/> NO <input type="checkbox"/></p> <p style="text-align: right;">PROBE AND CORRECT AS NECESSARY.</p>	
230I	<p>CHECK 230B, 230D AND 230F:</p> <p>HAD AT LEAST ONE INDUCED ABORTION, MISCARRIAGE OR STILLBIRTH <input type="checkbox"/></p> <p>HAD NO INDUCED ABORTIONS, MISCARRIAGES, OR STILLBIRTHS <input type="checkbox"/></p>	<p>→234</p>
231A	<p>Now I would like to ask about any recent induced abortions, miscarriages, or still births which you have had. When did the last such pregnancy ended ?</p>	<p>MONTH.....<input type="text"/><input type="text"/><input type="text"/><input type="text"/></p> <p>YEAR.....<input type="text"/><input type="text"/><input type="text"/><input type="text"/></p>
231B	<p>Was this an induced abortion, a miscarriage, or a still birth ?</p>	<p>INDUCED ABORTION.....1</p> <p>MISCARRIAGE.....2</p> <p>STILL BIRTH.....3 →232</p>
231C	<p>Whose decision was to end your pregnancy with an induced abortion ?</p>	<p>DOCTOR.....01</p> <p>HERSELF.....02</p> <p>HUSBAND.....03</p> <p>WOMAN AND HUSBAND TOGETHER.....04</p> <p>OTHER _____ 96</p> <p style="text-align: center;">(SPECIFY)</p>
231D	<p>What was the main reason that your pregnancy to end with an induced abortion ?</p>	<p>HEALTH REASONS</p> <p>RELATED TO MOTHER.....01</p> <p>RELATED TO CHILD.....02</p> <p>RELATED TO BOTH.....03</p> <p>DID NOT WANT TO HAVE ANOTHER CHILD.....04</p> <p>PREVIOUS PREGNANCY JUST ENDED.....05</p> <p>OTHER _____ 96</p> <p style="text-align: center;">(SPECIFY)</p>

231E

Where did the operation of induced abortion take place ?

(NAME OF PLACE)

PUBLIC SECTOR

STATE/SAMPLE HOSPITAL.....11
 MATERNITY HOUSE.....12
 MOTHER-CHILD HEALTH AND FAMILY
 PLANNING (MCHFP) CENTRE...13
 SSK HOSPITAL/DISPENSARY.....16
 OTHER PUBLIC SECTOR

19

(SPECIFY)

PRIVATE SECTOR

PRIVATE HOSPITAL.....21
 PRIVATE POLYCLINIC.....22
 PRIVATE DOCTOR.....23

OTHER PRIVATE SECTOR

29

(SPECIFY)

UNIVERSITY HOSPITAL.....31

OTHER _____ 96

(SPECIFY)

232

CHECK 231A:

LAST PREGNANCY
 ENDED SINCE
 JANUARY 1993

LAST PREGNANCY
 ENDED BEFORE
 JANUARY 1993

→234

233

How many months pregnant were you when the last pregnancy ended ?

MONTHS.....

C RECORD NUMBER OF COMPLETED MONTHS FOR ALL WASTED PREGNANCIES
 IN COLUMNS 1 AND 2.

- PROBE TO DETERMINE HOW THE PREGNANCY ENDED (INDUCED ABORTION, MISCARRIAGE, STILL BIRTH).

- IN COLUMN 1 ENTER THE APPROPRIATE CODE IN THE MONTH AND YEAR THE PREGNANCY TERMINATED.

CODES: F - SPONTANEOUS ABORTION
 K - INDUCED ABORTION
 J - STILL BIRTH

- DETERMINE THE NUMBER OF COMPLETED MONTHS AND ENTER 'H' FOR THE REMAINING NUMBER OF COMPLETED MONTHS.

- IF THE PREGNANCY ENDED WITH INDUCED ABORTION, ENTER THE CODE FOR THE PLACE OF INDUCED ABORTION IN COLUMN 2 OF CALENDAR IN THE MONTH THAT THE PREGNANCY TERMINATED.

CODES: C - STATE/SAMPLE HOSPITAL H - PRIVATE HOSPITAL
 D - MATERNITY HOUSE J - PRIVATE POLYCLINIC
 E - MCHFP CENTRE K - PRIVATE DOCTOR
 P - SSK HOSPITAL/DISPENSARY L - OTHER PRIVATE SECTOR
 G - OTHER PUBLIC SECTOR N - UNIVERSITY HOSPITAL

THEN ASK FOR DATES AND DURATIONS OF ANY OTHER PREGNANCIES BACK TO JANUARY 1993.

REPEAT THE PROCEDURES AS DESCRIBED ABOVE FOR THESE PREGNANCIES.

ILLUSTRATIVE QUESTIONS:

- How did this pregnancy end ? (Was it an induced abortion, miscarriage, or still birth etc.)
- What was the total duration of this pregnancy ? How many months pregnant were you ?
- Where did you have this induced abortion ?

<p>234</p>	<p>When did your last menstrual period start ?</p> <p>_____</p> <p>(DATE, IF GIVEN)</p> <p>RECORD THE ANSWER AS GIVEN.</p>	<p>DAYS AGO.....1</p> <p>WEEKS AGO.....2</p> <p>MONTHS AGO.....3</p> <p>YEARS AGO.....4</p> <table border="1" data-bbox="1234 283 1307 514"> <tr><td></td><td></td></tr> <tr><td></td><td></td></tr> <tr><td></td><td></td></tr> <tr><td></td><td></td></tr> </table> <p>CURRENTLY PREGNANT.....993</p> <p>IN MENOPAUSE.....994</p> <p>BEFORE LAST BIRTH.....995</p> <p>NEVER MENSTRUATED.....996</p>								
<p>235</p>	<p>Between the first day of a woman's period and the first day of her next period, are there certain times when she has a greater chance of becoming pregnant than other times ?</p>	<p>YES.....1</p> <p>NO.....2</p> <p>DON'T KNOW.....8 → 250</p>								
<p>236</p>	<p>During which times of the monthly cycle does a woman have the greatest chance of becoming pregnant ?</p>	<p>DURING HER PERIOD.....1</p> <p>RIGHT AFTER HER PERIOD HAS ENDED.....2</p> <p>IN THE MIDDLE OF THE CYCLE.....3</p> <p>JUST BEFORE HER PERIOD BEGINS...4</p> <p>OTHER _____ 7</p> <p>(SPECIFY)</p> <p>DON'T KNOW.....8</p>								

SECTION 2B. MARRIAGE

250	<p>Are you currently married ?</p> <p>ACCEPT THOSE LIVING TOGETHER AS BEING MARRIED.</p>	<p>YES, CURRENTLY MARRIED.....1 →253</p> <p>NO, CURRENTLY NOT MARRIED.....2</p>
252	<p>What is your current marital status ? Are you widowed, divorced, or separated ?</p>	<p>WIDOWED.....1</p> <p>DIVORCED.....2 →255</p> <p>SEPARATED.....3</p>
253	<p>Is your husband living with you now, or is he staying elsewhere because of work, military service, a journey abroad ?</p>	<p>LIVING WITH HER.....1</p> <p>STAYING ELSEWHERE.....2</p>
254	<p>ENTER THE LINE NUMBER OF HER HUSBAND FROM HOUSEHOLD SCHEDULE. IF NOT IN THE LIST, WRITE '96'.</p>	<div style="border: 1px solid black; width: 20px; height: 20px; margin: 0 auto;"></div>
255	<p>Did you marry only once or more than once?</p>	<p>ONCE.....1</p> <p>MORE THAN ONCE.....2</p>
257	<p>CHECK 255:</p> <p>MARRIED ONCE <input type="checkbox"/> MARRIED MORE THAN ONCE <input type="checkbox"/></p> <p>↓ ↓</p> <p>In what month and year did you marry (started living with) your husband ?</p> <p>Now, let's talk about your first husband. In what month and year did you marry (started living with) your husband ?</p>	<p>MONTH..... <div style="border: 1px solid black; width: 20px; height: 20px; display: inline-block;"></div></p> <p>DON'T KNOW MONTH.....98</p> <p>YEAR..... <div style="border: 1px solid black; padding: 2px;">1 9</div> →259</p> <p>DON'T KNOW YEAR.....9998</p>
258	<p>How old were you when you started living with your (first) husband ?</p>	<p>AGE..... <div style="border: 1px solid black; width: 20px; height: 20px; display: inline-block;"></div></p>
259	<p>How old was your (first) husband when you started living with him ?</p> <p>IF THE WOMAN DOES NOT KNOW HER (FIRST) HUSBAND'S AGE AT MARRIAGE, ASK HOW MANY YEARS DIFFERENCE IS THERE BETWEEN HER AND HER (FIRST) HUSBAND AND ESTIMATE HER (FIRST) HUSBAND'S MARRIAGE AGE.</p>	<p>AGE..... <div style="border: 1px solid black; width: 20px; height: 20px; display: inline-block;"></div></p> <p>DON'T KNOW.....98</p>

262	<p>C CHECK 250, IF CURRENTLY MARRIED ENTER 'X' IN THE MONTH OF INTERVIEW IN COLUMN 3 OF CALENDAR.</p> <p>THEN, DETERMINE MONTHS MARRIED OR IN UNION SINCE JANUARY 1993. ENTER 'X' IN COLUMN 3 OF CALENDAR FOR EACH MONTH MARRIED OR IN UNION, AND ENTER 'O' FOR EACH MONTH NOT MARRIED/NOT IN UNION.</p> <p>FOR WOMEN WITH MORE THAN ONE UNION: PROBE FOR THE DATE WHEN CURRENT UNION STARTED AND, IF APPROPRIATE, PROBE FOR THE STARTING AND TERMINATION DATES OF ANY PREVIOUS UNIONS.</p> <p>FOR WOMEN NOT CURRENTLY IN UNION: PROBE FOR DATE WHEN LAST UNION STARTED AND FOR TERMINATION DATE AND, IF ANY, FOR THE STARTING AND TERMINATION DATES OF ANY PREVIOUS UNIONS.</p> <p>NOTE : AFTER YOU HAVE COMPLETED THESE, ALL THE BOXES IN COLUMN 3 FROM JANUARY 1993 TO INTERVIEW MONTH SHOULD BE FILLED.</p>					
263	<p>Now I want to ask some questions about your marriage(s).</p> <p>CHECK 255: IF MARRIED ONLY ONCE, ASK ABOUT HER CURRENT/LAST HUSBAND. IF MARRIED MORE THAN ONCE, FIRST ASK ABOUT HER CURRENT/LAST HUSBAND AND THEN ABOUT HER FIRST HUSBAND.</p> <p>WRITE NAME(S) OF HER HUSBAND(S) AND USE THESE NAMES WHEN ASKING QUESTIONS.</p>					
	<table style="width: 100%; border: none;"> <tr> <td style="border: none; width: 50%; text-align: center;">CURRENT/LAST HUSBAND</td> <td style="border: none; width: 50%; text-align: center;">FIRST HUSBAND</td> </tr> <tr> <td style="border: none; text-align: center;">NAME _____</td> <td style="border: none; text-align: center;">NAME _____</td> </tr> </table>	CURRENT/LAST HUSBAND	FIRST HUSBAND	NAME _____	NAME _____	
CURRENT/LAST HUSBAND	FIRST HUSBAND					
NAME _____	NAME _____					
265A	<p>Did you have a civil marriage ceremony with your husband ?</p> <p>YES.....1 NO.....2</p>	<p>YES.....1 NO.....2</p>				
265B	<p>Did you have a religious ceremony with your husband ?</p> <p>YES.....1 NO.....2</p>	<p>YES.....1 NO.....2</p>				
265C	<p>CHECK 265A AND 265B:</p> <p>CIVIL AND RELIGIOUS CEREMONIES...1</p> <p>ONLY CIVIL CEREMONY.....2 → 268</p> <p>ONLY RELIGIOUS CEREMONY OR NO CEREMONY...3 → 267E</p>	<p>CIVIL AND RELIGIOUS CEREMONIES...1</p> <p>ONLY CIVIL CEREMONY.....2 } → 268</p> <p>ONLY RELIGIOUS CEREMONY OR NO CEREMONY...3 }</p>				
266	<p>Which marriage ceremony took place earlier?</p> <p>CIVIL.....1 RELIGIOUS.....2</p>	<p>CIVIL.....1 RELIGIOUS.....2</p>				
267A	<p>How much time elapsed between two ceremonies ?</p> <p>RECORD '00' DAYS IF BOTH TOOK PLACE ON THE SAME DAY.</p> <p>IF LESS THAN ONE MONTH RECORD AS DAY, IF LESS THAN TWO YEARS RECORD AS MONTH, OTHERWISE RECORD AS YEAR.</p> <p>YEARS.....1</p> <p>MONTHS.....2</p> <p>DAYS.....3</p>	<p>YEARS.....1</p> <p>MONTHS.....2</p> <p>DAYS.....3</p>				

		CURRENT/LAST HUSBAND		FIRST HUSBAND	
		NAME _____		NAME _____	
267B	CHECK 266:	RELIGIOUS IS EARLIER	CIVIL IS EARLIER	RELIGIOUS IS EARLIER	CIVIL IS EARLIER
		→268		→268	

267C

What is (was) the main reason that you have religious ceremony before the civil ceremony ?

RECORD ALL MENTIONED.

TRADITION.....A
 PREGNANCY/BIRTH.....B
 CHILD GOING SCHOOL...C
 CONFIDENCE.....D
 NEIGH. PRESSURE.....E
 REACHED ELIGIBLE
 AGE TO HAVE CIVIL
 CEREMONY.....P
 TO START THEIR
 MARITAL
 LIFE.....G

OTHER _____ U
 (SPECIFY)

TRADITION.....A
 PREGNANCY/BIRTH.....B
 CHILD GOING SCHOOL...C
 CONFIDENCE.....D
 NEIGH. PRESSURE.....E
 REACHED ELIGIBLE
 AGE TO HAVE CIVIL
 CEREMONY.....P
 TO START THEIR
 MARITAL
 LIFE.....G

OTHER _____ U
 (SPECIFY)

267D

SKIP TO 268.

SKIP TO 268.

267E

ASK IF CURRENTLY MARRIED (Q. 250=1).

In the future do you and/or your husband intend to have a civil ceremony ?

YES, ONLY WOMAN.....1
 YES, ONLY HUSBAND...2
 YES, BOTH.....3
 NO, NONE OF THEM....4
 OTHER _____ 7
 (SPECIFY)
 DON'T KNOW.....8

268

How was your marriage with your husband arranged ? Have you decided together or was it arranged by your families ?

BY OURSELVES.....1
 BY FAMILIES.....2
 270←
 ESCAPED/ABDUCTED....3
 OTHER _____ 7
 (SPECIFY)
 272←

BY OURSELVES.....1
 BY FAMILIES.....2
 270←
 ESCAPED/ABDUCTED....3
 OTHER _____ 7
 (SPECIFY)
 272←

269

Did you have to seek consent of your family to get married ?

YES.....1
 NO.....2
 272←

YES.....1
 NO.....2
 272←

270

Did your family seek your consent on your marriage ?

YES.....1
 NO.....2

YES.....1
 NO.....2

		CURRENT/LAST HUSBAND	FIRST HUSBAND
		NAME _____	NAME _____
271	Do you think that your family would insist on your marriage, even if you do not agree to marry ?	YES.....1 NO.....2 DON'T KNOW.....8	YES.....1 NO.....2 DON'T KNOW.....8
272	Did your husband or his family pay bridesmoney to your family ? IF YES: Was it given in cash or in kind ?	NO.....1 YES, IN CASH.....2 YES, IN KIND.....3 YES, BOTH.....4 OTHER _____ 7 (SPECIFY)	NO.....1 YES, IN CASH.....2 YES, IN KIND.....3 YES, BOTH.....4 OTHER _____ 7 (SPECIFY)
273A	When you first started to live with your husband, was there anyone else living with you in your household at that time ?	YES.....1 NO.....2 275A ←	YES.....1 NO.....2 275A ←
273B	Who else was with you ? Who else? RECORD ALL MENTIONED.	WOMAN'S MOTHER/FATHER.....A SIBLING(S).....B GRANDMOTHER/PATHER.C OTHER RELATIVES...D CHILDREN.....E HER HUSBAND'S MOTHER/FATHER.....F SIBLING(S).....G GRANDMOTHER/PATHER.H OTHER RELATIVES...I CHILDREN.....J NOT RELATIVES OF HER OR HER HUSBAND.....K OTHER _____ U (SPECIFY) OTHER _____ V (SPECIFY)	WOMAN'S MOTHER/FATHER.....A SIBLING(S).....B GRANDMOTHER/PATHER.C OTHER RELATIVES...D CHILDREN.....E HER HUSBAND'S MOTHER/FATHER.....F SIBLING(S).....G GRANDMOTHER/PATHER.H OTHER RELATIVES...I CHILDREN.....J NOT RELATIVES OF HER OR HER HUSBAND.....K OTHER _____ U (SPECIFY) OTHER _____ V (SPECIFY)
274	Approximately, how many years did you live with these person(s).	YEARS..... <input type="text"/> <input type="text"/> STILL LIVING WITH...96	YEARS..... <input type="text"/> <input type="text"/> STILL LIVING WITH...96
275A	Do (did) you have any relativeness with your husband ?	YES.....1 NO.....2 278 ←	YES.....1 NO.....2 277 ←

		CURRENT/LAST HUSBAND	FIRST HUSBAND
		NAME _____	NAME _____
276A	What is (was) his relationship to you ?	FATHER'S BROTHER'S SON..01 FATHER'S SISTER'S SON..02 MOTHER'S SISTER'S SON..03 MOTHER'S BROTHER'S SON..04 OTHER _____ 96 (SPECIFY)	FATHER'S BROTHER'S SON..01 FATHER'S SISTER'S SON..02 MOTHER'S SISTER'S SON..03 MOTHER'S BROTHER'S SON..04 OTHER _____ 96 (SPECIFY)
276B	What was the main reason to marry with a relative ?	PROTECTING FAMILY ASSET.....01 NOT TO ALLOW NONRELATIVES INTO FAMILY.....02 ADVANTAGE OF KNOWING HIM BEFORE.....03 LOVED.....04 ARRANGED DURING INFANCY.....05 OTHER _____ 96 (SPECIFY)	PROTECTING FAMILY ASSET.....01 NOT TO ALLOW NONRELATIVES INTO FAMILY.....02 ADVANTAGE OF KNOWING HIM BEFORE.....03 LOVED.....04 ARRANGED DURING INFANCY.....05 OTHER _____ 96 (SPECIFY)
277	How was this marriage ended ? Were you divorced or widowed ?		DIVORCED.....1 WIDOWED.....2
278		IF MARRIED MORE THAN ONCE .PROCEED WITH HER FIRST HUSBAND IN Q. 265A. IF NOT SKIP TO Q. 301.	SKIP TO 301.

SECTION 3. CONTRACEPTION

Now I would like to talk about family planning. There are various methods that a married couple can use to avoid pregnancy.

- CIRCLE CODE '1' IN IN Q. 301 FOR EACH METHOD MENTIONED SPONTANEOUSLY.
- THEN PROCEED DOWN COLUMN 302, READING THE NAME AND DESCRIPTION OF EACH METHOD NOT MENTIONED SPONTANEOUSLY AND ASK WHETHER SHE HAS HEARD THE METHOD. CIRCLE CODE '2' IF METHOD IS RECOGNIZED AND CODE '3' IF NOT RECOGNIZED.
- THEN FOR EACH METHOD WITH CODE '1' OR '2' CIRCLED IN 301 OR 302, ASK 303.
- AFTER ASKING ABOUT ALL METHODS PROCEED TO 304.

301 Which ways or methods have you heard ?	SPON- TANEOUS YES	302 Have you ever heard this method ?		303 Have you ever used this method ?
		PROBED YES	NO	
01 PILL Woman can avoid a pregnancy by taking a pill every day.	1	2	3	YES.....1 NO.....2
02 IUD Women can have the so called spiral or IUD placed in them by a doctor or a nurse.	1	2	3	YES.....1 NO.....2
03 INJECTABLES Woman can have an injection by a doctor or a nurse which stops them from becoming pregnant for a certain period of time.	1	2	3	YES.....1 NO.....2
04 NORPLANT Woman can have small rods placed in their arm and this can prevent pregnancy for several years.	1	2	3	YES.....1 NO.....2
05 DIAPHRAGM, FOAM, JELLY Woman can place a sponge, suppository, diaphragm, jelly, or cream inside themselves before intercourse.	1	2	3	YES.....1 NO.....2
06 CONDOM Men can put a rubber sheath on their penis during sexual intercourse.	1	2	3	YES.....1 NO.....2
07 TUBAL LIGATION Women can have an operation of tubal ligation to avoid having any more children.	1	2	3	Have you ever had such an operation to avoid having any more children ? YES.....1 NO.....2
08 MALE STERILIZATION Men can have an operation called vasectomy so that their wives would not get pregnant.	1	2	3	Has (had) your husband ever have such an operation ? YES.....1 NO.....2

301 Which ways or methods have you heard ?	SPON- TANEOUS YES	302 Have you ever heard this method ? PROBED YES NO		303 Have you ever used this method ?
09 RHYTHM Some couples can avoid having sexual intercourse on certain days of the month when the woman is more likely to become pregnant.	1	2	3	YES.....1 NO.....2
10 WITHDRAWAL Some men pull out during sexual intercourse before climax.	1	2	3	YES.....1 NO.....2
11 Have you heard of any other method that women or men can use to avoid pregnancy ?	1	(SPECIFY)		YES.....1 NO.....2
		(SPECIFY)		YES.....1 NO.....2
		(SPECIFY)		YES.....1 NO.....2
304 CHECK 303: NOT A SINGLE "YES" (NEVER USED) <input type="checkbox"/> AT LEAST ONE "YES" (EVER USED) <input type="checkbox"/> → 308				
305 Have you ever used anything or tried in any way to delay or avoid getting pregnant ?				YES.....1 → 307 NO.....2
306 C ENTER "0" IN COLUMN 1 OF CALENDAR IN EACH BLANK MONTH. → 331				
307 What have you used or done ? CORRECT 303 AND 304 (AND 302 IF NECESSARY).				
308 Now I would like to ask you about the first time that you did something or used a method to avoid getting pregnant. What was the first method you ever used ?	PILL.....01 IUD.....02 INJECTABLES.....03 NORPLANT.....04 DIAPHRAGM/FOAM/JELLY.....05 CONDOM.....06 TUBAL LIGATION.....07 MALE STERILIZATION.....08 RHYTHM.....09 WITHDRAWAL.....10 OTHER _____ 96 (SPECIFY)			

309	Did you have any children at that time ? IF YES: How many living children did you have at that time ? IF NONE, RECORD "00".	NUMBER OF CHILDREN..... <input style="width: 20px; height: 20px;" type="text"/> <input style="width: 20px; height: 20px;" type="text"/>
310	CHECK 303: NOT HAD TUBAL LIGATION <input type="checkbox"/> HAD TUBAL LIGATION <input type="checkbox"/>	→314A
311	CHECK 227: NOT PREGNANT OR UNSURE <input type="checkbox"/> PREGNANT <input type="checkbox"/>	→324B
312	CHECK 250: CURRENTLY MARRIED <input type="checkbox"/> NOT MARRIED <input type="checkbox"/>	→324B
313	Are you currently doing something or using any method to delay or avoid getting pregnant ?	YES.....1 NO.....2 →325
314	Which method are you using ?	PILL.....01 IUD.....02 INJECTABLES.....03 NORPLANT.....04 →324A DIAPHRAGM/FOAM/JELLY.....05 CONDOM.....06 TUBAL LIGATION.....07 MALE STERILIZATION.....08 RHYTHM.....09 →323 WITHDRAWAL.....10 OTHER _____ 96 →324A (SPECIFY)
314A	CIRCLE '07' FOR TUBAL LIGATION.	
318	Where did the tubal ligation operation take place ? _____ (NAME OF PLACE)	PUBLIC SECTOR GOVERNMENT/SAMPLE HOSPITAL...11 MATERNITY HOUSE.....12 MOTHER-CHILD HEALTH AND FAMILY PLANNING (MCHFP) CENTRE...13 SSK HOSPITAL/DISPENSARY.....16 OTHER PUBLIC SECTOR _____ 19 (SPECIFY) PRIVATE SECTOR PRIVATE HOSPITAL.....21 PRIVATE POLYCLINIC.....22 PRIVATE DOCTOR.....23 PRIVATE NURSE/MIDWIPE.....24 OTHER PRIVATE SECTOR _____ 29 (SPECIFY) UNIVERSITY HOSPITAL.....31 OTHER _____ 96 (SPECIFY) DON'T KNOW.....98

319	Do you regret that you or your husband had the operation not to have any (more) children ?	YES.....1 NO.....2 → 321
-----	--	-----------------------------

320	Why do you regret the operation ?	WOMAN WANTS ANOTHER CHILD.....01 PARTNER WANTS ANOTHER CHILD....02 SIDE EFFECTS.....03 CHILD DIED.....04 OTHER _____ 96 (SPECIFY)
-----	-----------------------------------	--

321	In what month and year was this operation performed ?	MONTH..... <input type="text"/> <input type="text"/> YEAR..... <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>
-----	---	---

<p>322 CHECK 321:</p> <p>HAD TUBAL LIGATION OPERATION BEFORE <input type="checkbox"/> JANUARY 1993</p> <p>ENTER CODE FOR TUBAL LIGATION IN MONTH OF INTERVIEW IN COLUMN 1 OF THE CALENDAR AND EACH MONTH BACK TO JANUARY 1993.</p> <p>THEN SKIP TO 329A.</p>	C	<p>HAD TUBAL LIGATION OPERATION AFTER <input type="checkbox"/> JANUARY 1993</p> <p>ENTER CODE FOR TUBAL LIGATION IN MONTH OF INTERVIEW IN COLUMN 1 OF THE CALENDAR AND IN EACH MONTH BACK TO THE DATE OF THE OPERATION.</p> <p>THEN SKIP TO 324B.</p>
--	---	---

323	How do you determine the times not to have sexual intercourse ?	BASED ON CALENDAR.....01 BASED ON BODY TEMPERATURE.....02 BASED ON CERVICAL MUCUS.....03 BASED ON BODY TEMPERATURE AND CERVICAL MUCUS.....04 NO SPECIFIC SYSTEM.....05 OTHER _____ 96 (SPECIFY)
-----	---	---

324A	<p>C ENTER METHOD CODE FROM 314 IN CURRENT MONTH IN COLUMN 1 OF CALENDAR. THEN DETERMINE WHEN SHE STARTED USING THIS METHOD. ENTER METHOD CODE IN EACH MONTH OF USE.</p> <p>ILLUSTRATIVE QUESTIONS:</p> <p style="padding-left: 40px;">When did you start using this method continuously ?</p> <p style="padding-left: 40px;">How long have you been using this method continuously ?</p>	
------	--	--

324B	<p>C CHECK COLUMN 3 OF CALENDAR:</p> <p style="padding-left: 40px;">IN COLUMN 1 OF CALENDAR ENTER "N" FOR MONTHS WOMAN NOT MARRIED.</p>	
------	--	--

324C

C CHECK COLUMN 1 OF CALENDAR:

THERE ARE EMPTY BOXES

ALL BOXES ARE FILLED

→326A

325

I would like to ask you some questions about the times you may have used a method to avoid getting pregnant during the last few years.

C START WITH THE MOST RECENT USE. USE CALENDAR TO PROBE FOR EARLIER PERIODS OF USE AND NONUSE BACK TO JANUARY 1993. USE NAMES OF CHILDREN, DATES OF BIRTH, AND STARTING AND ENDING DATES OF PREGNANCIES AS REFERENCE POINTS.

IN COLUMN 1, ENTER CODE IN EACH MONTH OF METHOD USE OR '0' FOR NONUSE.

ILLUSTRATIVE QUESTIONS FOR COLUMN 1:

- *When was the last time you used a method ? Which method was that ?
- *When did you start using that method ? How long after the birth of (NAME) ?
- *How long did you use the method then ?

IN COLUMN 2, ENTER CODES FOR DISCONTINUATION NEXT TO LAST MONTH OF USE. TO DO THIS, DETERMINE THE LAST MONTH OF METHOD USE. IN COLUMN 2 ENTER THE CODE FOR DISCONTINUATION.

ASK WHY SHE STOPPED USING THE METHOD. IF A PREGNANCY FOLLOWED, ASK WHETHER SHE BECAME PREGNANT UNINTENTIONALLY WHILE USING THE METHOD OR DELIBERATELY STOPPED TO GET PREGNANT.

ILLUSTRATIVE QUESTIONS FOR COLUMN 2:

- *Why did you stop using the (METHOD) ?
- *Did you become pregnant while using (METHOD), or did you stop to get pregnant, or did you stop for some other reason ?

IF DELIBERATELY STOPPED TO BECOME PREGNANT, ASK:

- "How many months did it take you to get pregnant after you stopped using (METHOD) ?
- AND ENTER '0' IN EACH SUCH MONTH IN COLUMN 1.

NOTE: PAY PARTICULAR ATTENTION FOR LONG PERIODS OF NONUSE. THERE MAY BE SOME PERIODS OF METHOD USE OR A PREGNANCY DURING THESE PERIODS.

NOTE: AFTER COMPLETING THIS SECTION, ALL THE BOXES IN COLUMN 1 OF CALENDAR MUST BE FILLED.

326A

CHECK 314:

CIRCLE THE CODE OF CURRENTLY USED METHOD.

NOT ASKED.....	00	→331
PILL.....	01	
IUD.....	02	
INJECTABLES.....	03	
NORPLANT.....	04	
DIAPHRAGM/FOAM/JELLY.....	05	
CONDOM.....	06	
TUBAL LIGATION.....	07	
MALE STERILIZATION.....	08	→329A
RHYTHM.....	09	
WITHDRAWAL.....	10	
OTHER METHOD.....	96	

326B	Would you like to use a different method of family planning than the one you are currently using ?	YES.....1 NO.....2	→327
326C	Which method would you prefer to use ?	PILL.....01 IUD.....02 INJECTABLES.....03 NORPLANT.....04 DIAPHRAGM/FOAM/JELLY.....05 CONDOM.....06 TUBAL LIGATION.....07 MALE STERILIZATION.....08 RHYTHM.....09 WITHDRAWAL.....10 ANY METHOD.....77 NOT SURE.....88 OTHER _____ 96 (SPECIFY)	
326D	What is the reason that you do not use (METHOD MENTIONED IN 326C) ?	DOCTOR DOES NOT ADVISE.....01 EXPENSIVE.....02 NOT AVAILABLE/ACCESS PROBLEMS..03 TOO FAR AWAY/HARD TO FIND HERE.04 DON'T KNOW HOW TO OBTAIN.....05 DON'T KNOW HOW TO USE IT.....06 HUSBAND OBJECTS.....07 RELIGIOUS REASONS.....08 OTHER _____ 96 (SPECIFY) DON'T KNOW.....98	→327
326E	Why ?	TOO YOUNG.....01 TOO OLD.....02 HAD MANY BIRTHS BEFORE.....03 DOES NOT HAVE ENOUGH CHILDREN YET.....04 HEALTH REASONS _____ 05 (SPECIFY) OTHER _____ 96 (SPECIFY)	

327	<p>CHECK 314:</p> <p>CIRCLE CODE OF CURRENTLY USED METHOD.</p>	<table border="0"> <tr><td>NOT ASKED.....</td><td>00</td><td>→331</td></tr> <tr><td>PILL.....</td><td>01</td><td></td></tr> <tr><td>IUD.....</td><td>02</td><td></td></tr> <tr><td>INJECTABLES.....</td><td>03</td><td></td></tr> <tr><td>NORPLANT.....</td><td>04</td><td></td></tr> <tr><td>DIAPHRAGM/FOAM/JELLY.....</td><td>05</td><td></td></tr> <tr><td>CONDOM.....</td><td>06</td><td></td></tr> <tr><td>TUBAL LIGATION.....</td><td>07</td><td></td></tr> <tr><td>MALE STERILIZATION.....</td><td>08</td><td>→329A</td></tr> <tr><td>RHYTHM.....</td><td>09</td><td></td></tr> <tr><td>WITHDRAWAL.....</td><td>10</td><td></td></tr> <tr><td>OTHER METHOD.....</td><td>96</td><td>→332</td></tr> </table>	NOT ASKED.....	00	→331	PILL.....	01		IUD.....	02		INJECTABLES.....	03		NORPLANT.....	04		DIAPHRAGM/FOAM/JELLY.....	05		CONDOM.....	06		TUBAL LIGATION.....	07		MALE STERILIZATION.....	08	→329A	RHYTHM.....	09		WITHDRAWAL.....	10		OTHER METHOD.....	96	→332
NOT ASKED.....	00	→331																																				
PILL.....	01																																					
IUD.....	02																																					
INJECTABLES.....	03																																					
NORPLANT.....	04																																					
DIAPHRAGM/FOAM/JELLY.....	05																																					
CONDOM.....	06																																					
TUBAL LIGATION.....	07																																					
MALE STERILIZATION.....	08	→329A																																				
RHYTHM.....	09																																					
WITHDRAWAL.....	10																																					
OTHER METHOD.....	96	→332																																				

328	<p>Where did you obtain (METHOD) ?</p> <p>_____</p> <p>(NAME OF PLACE)</p>	<table border="0"> <tr><td colspan="2">PUBLIC SECTOR</td></tr> <tr><td>GOVERNMENT/SAMPLE HOSPITAL.....</td><td>11</td></tr> <tr><td>MATERNITY HOUSE.....</td><td>12</td></tr> <tr><td>MOTHER-CHILD HEALTH AND FAMILY PLANNING (MCHFP) CENTRE.....</td><td>13</td></tr> <tr><td>HEALTH CENTRE.....</td><td>14</td></tr> <tr><td>HEALTH HOUSE.....</td><td>15</td></tr> <tr><td>SSK HOSPITAL/DISPENSARY.....</td><td>16</td></tr> <tr><td>OTHER PUBLIC SECTOR</td><td>_____ 19</td></tr> <tr><td></td><td>(SPECIFY)</td></tr> <tr><td colspan="2">PRIVATE SECTOR</td></tr> <tr><td>PRIVATE HOSPITAL.....</td><td>21</td></tr> <tr><td>PRIVATE POLYCLINIC.....</td><td>22</td></tr> <tr><td>PRIVATE DOCTOR.....</td><td>23</td></tr> <tr><td>PRIVATE MIDWIFE/NURSE.....</td><td>24</td></tr> <tr><td>PHARMACY.....</td><td>25</td></tr> <tr><td>OTHER PRIVATE SECTOR</td><td>_____ 29</td></tr> <tr><td></td><td>(SPECIFY)</td></tr> <tr><td>UNIVERSITY HOSPITAL.....</td><td>31</td></tr> <tr><td>COMMUNITY VOLUNTEERS/ ASSOCIATION/FOUNDATION.....</td><td>41</td></tr> <tr><td>MARKET/SHOP.....</td><td>51</td></tr> <tr><td>FRIEND/RELATIVE/NEIGHBOUR.....</td><td>52</td></tr> <tr><td>TRAD.MIDWIFE/MIDWIFE GRAN.....</td><td>53</td></tr> <tr><td>OTHER _____</td><td>96</td></tr> <tr><td></td><td>(SPECIFY)</td></tr> </table>	PUBLIC SECTOR		GOVERNMENT/SAMPLE HOSPITAL.....	11	MATERNITY HOUSE.....	12	MOTHER-CHILD HEALTH AND FAMILY PLANNING (MCHFP) CENTRE.....	13	HEALTH CENTRE.....	14	HEALTH HOUSE.....	15	SSK HOSPITAL/DISPENSARY.....	16	OTHER PUBLIC SECTOR	_____ 19		(SPECIFY)	PRIVATE SECTOR		PRIVATE HOSPITAL.....	21	PRIVATE POLYCLINIC.....	22	PRIVATE DOCTOR.....	23	PRIVATE MIDWIFE/NURSE.....	24	PHARMACY.....	25	OTHER PRIVATE SECTOR	_____ 29		(SPECIFY)	UNIVERSITY HOSPITAL.....	31	COMMUNITY VOLUNTEERS/ ASSOCIATION/FOUNDATION.....	41	MARKET/SHOP.....	51	FRIEND/RELATIVE/NEIGHBOUR.....	52	TRAD.MIDWIFE/MIDWIFE GRAN.....	53	OTHER _____	96		(SPECIFY)
PUBLIC SECTOR																																																		
GOVERNMENT/SAMPLE HOSPITAL.....	11																																																	
MATERNITY HOUSE.....	12																																																	
MOTHER-CHILD HEALTH AND FAMILY PLANNING (MCHFP) CENTRE.....	13																																																	
HEALTH CENTRE.....	14																																																	
HEALTH HOUSE.....	15																																																	
SSK HOSPITAL/DISPENSARY.....	16																																																	
OTHER PUBLIC SECTOR	_____ 19																																																	
	(SPECIFY)																																																	
PRIVATE SECTOR																																																		
PRIVATE HOSPITAL.....	21																																																	
PRIVATE POLYCLINIC.....	22																																																	
PRIVATE DOCTOR.....	23																																																	
PRIVATE MIDWIFE/NURSE.....	24																																																	
PHARMACY.....	25																																																	
OTHER PRIVATE SECTOR	_____ 29																																																	
	(SPECIFY)																																																	
UNIVERSITY HOSPITAL.....	31																																																	
COMMUNITY VOLUNTEERS/ ASSOCIATION/FOUNDATION.....	41																																																	
MARKET/SHOP.....	51																																																	
FRIEND/RELATIVE/NEIGHBOUR.....	52																																																	
TRAD.MIDWIFE/MIDWIFE GRAN.....	53																																																	
OTHER _____	96																																																	
	(SPECIFY)																																																	

329	<p>Do you know another place where you could have obtained (METHOD) ?</p>	<table border="0"> <tr><td>YES.....</td><td>1</td><td></td></tr> <tr><td>NO.....</td><td>2</td><td>→337</td></tr> </table>	YES.....	1		NO.....	2	→337
YES.....	1							
NO.....	2	→337						
329A	<p>At the time of tubal ligation operation, did you know another place where you could have the operation ?</p>							

330

People select the place where they get family planning services for various reasons.

What was the main reason you went to (NAME OF PLACE IN Q.328 OR Q.318) to obtain (METHOD) instead of some other place you know about ?

RECORD RESPONSE AND CIRCLE CODE.

ACCESS-RELATED REASONS

- CLOSER TO HOME.....11
- CLOSER TO MARKET/WORK.....12
- AVAILABILITY OF TRANSPORT....13

SERVICE-RELATED REASONS

- STAFF MORE COMPETENT/
FRIENDLY.....21
- CLEANER FACILITY.....22
- OFFERS MORE PRIVACY.....23
- SHORTER WAITING TIME.....24
- LONGER/SUITABLE HOURS
OF OPERATION.....25
- USE OTHER SERVICES AT
FACILITY.....26

- LOWER COST/CHEAPER.....31
- WANTED ANONYMITY.....41
- HAD TUBAL LIGATION
DURING DELIVERY.....51
- OTHER _____ 96
(SPECIFY)
- DON'T KNOW.....98

→337

331

CHECK 227:

NOT PREGNANT
OR UNSURE

CURRENTLY
PREGNANT

→332

331A

What is the main reason you are not using a method of contraception to avoid pregnancy ?

NOT MARRIED.....11

FERTILITY-RELATED REASONS

- NOT HAVING SEX.....21
- INFREQUENT SEX.....22
- MENOPAUSAL/HYSTERECTOMY.....23
- SUBFECUND/INFECUND.....24
- POSTPARTUM/BREASTFEEDING....25
- WANTS (MORE) CHILDREN.....26

OPPOSITION TO USE

- WOMAN OPPOSED.....31
- HUSBAND OPPOSED.....32
- OTHERS OPPOSED.....33
- RELIGIOUS REASONS.....34

LACK OF KNOWLEDGE

- KNOWS NO METHOD.....41
- KNOWS NO SOURCE.....42

METHOD-RELATED REASONS

- HEALTH CONCERNS.....51
- SIDE EFFECTS.....52
- LACK OF ACCESS/TOO FAR.....53
- COST TOO MUCH.....54
- INCONVENIENT TO USE.....55
- INTERFERES WITH BODY'S
NORMAL PROCESSES.....56

- OTHER _____ 96
(SPECIFY)
- DON'T KNOW.....98

332	Do you know of a place where you can obtain a method of family planning ?	YES.....1 NO.....2 → 337
333	Where is that ? <hr style="width: 20%; margin: 10px auto;"/> (NAME OF PLACE) <hr style="width: 20%; margin: 10px auto;"/> (NAME OF PLACE)	PUBLIC SECTOR GOVERNMENT/SAMPLE HOSPITAL...A MATERNITY HOUSE.....B MOTHER-CHILD HEALTH AND FAMILY PLANNING (MCHPP) CENTRE...C HEALTH CENTRE.....D HEALTH HOUSE.....E SSK HOSPITAL/DISPENSARY.....F OTHER PUBLIC SECTOR _____ G (SPECIFY) PRIVATE SECTOR PRIVATE HOSPITAL.....H PRIVATE POLYCLINIC.....I PRIVATE DOCTOR.....J PRIVATE MIDWIPE/NURSE.....K PHARMACY.....L OTHER PRIVATE SECTOR _____ M (SPECIFY) UNIVERSITY HOSPITAL.....N COMMUNITY VOLUNTEERS/ ASSOCIATION/FOUNDATION.....O MARKET/SHOP.....P FRIEND/RELATIVE/NEIGHBOUR.....R TRAD. MIDWIPE/MIDWIPE GRAN.....S OTHER _____ U (SPECIFY)
337	Do you think a woman's chance of becoming pregnant is influenced by breastfeeding ?	YES.....1 NO.....2 DON'T KNOW.....8 → 401
338	Do you think a woman's chance of becoming pregnant is increased or decreased by breastfeeding ?	INCREASED.....1 → 401 DECREASED.....2 DEPENDS.....3 DON'T KNOW.....8
339	CHECK 208: ONE OR MORE BIRTHS <input type="checkbox"/> NO BIRTHS <input type="checkbox"/>	→ 601
340	Have you ever relied on breastfeeding as a method of avoiding pregnancy ?	YES.....1 NO.....2 → 401

341	CHECK 227 AND 314:	NOT PREGNANT OR UNSURE AND HAD NO TUBAL LIGATION	<input type="checkbox"/>	EITHER PREGNANT OR HAD TUBAL LIGATION	<input type="checkbox"/>	→401
-----	--------------------	--	--------------------------	--	--------------------------	------

342	Are you currently relying on breastfeeding to avoid getting pregnant ?	YES.....1 NO.....2
-----	---	-----------------------

SECTION 4A. PREGNANCY AND BREASTFEEDING

401	CHECK 225:	ONE OR MORE LIVE BIRTHS SINCE JAN. 1993	<input type="checkbox"/>	NO LIVE BIRTHS SINCE JAN. 1993	<input type="checkbox"/>	→601
-----	------------	---	--------------------------	--------------------------------	--------------------------	------

402 ENTER THE LINE NUMBER, NAME, AND SURVIVAL STATUS OF EACH BIRTH SINCE JANUARY 1993 IN THE TABLE ASK THE QUESTIONS ABOUT ALL OF THESE BIRTHS. BEGIN WITH THE LAST BIRTH. (IF THERE ARE MORE THAN 2 BIRTHS USE ADDITIONAL QUESTIONNAIRES- DO NOT USE THE LAST BIRTH COLUMN IN THE ADDITIONAL QUESTIONNAIRE).

I would like to ask you some more questions about the health of all your children born in the past five years. We will talk about one child at a time.

		LAST BIRTH		NEXT TO LAST BIRTH	
403	LINE NUMBER FROM Q212.	LINE NUMBER.....	<input type="text"/>	LINE NUMBER.....	<input type="text"/>

		NAME		NAME	
404	FROM Q212	_____		_____	
	FROM Q216	ALIVE <input type="checkbox"/>	DEAD <input type="checkbox"/>	ALIVE <input type="checkbox"/>	DEAD <input type="checkbox"/>

405	At the time you became pregnant with (NAME), did you want to become pregnant then, did you want to wait until later, or did you want no (more) children at all ?	THEN.....1	<input type="checkbox"/>	THEN.....1	<input type="checkbox"/>
		LATER.....2	<input type="checkbox"/>	LATER.....2	<input type="checkbox"/>
		NO MORE.....3	<input type="checkbox"/>	NO MORE.....3	<input type="checkbox"/>
		407A ←		407A ←	

406	How much longer would you like to have waited ?	MONTHS.....1	<input type="text"/>	MONTHS.....1	<input type="text"/>
		YEARS.....2	<input type="text"/>	YEARS.....2	<input type="text"/>
		DON'T KNOW.....998		DON'T KNOW.....998	

407A	When you were pregnant with (NAME) did you see anyone for antenatal care for this pregnancy ?	HEALTH PROFESSIONAL		HEALTH PROFESSIONAL	
	IF YES: Whom did you see ?	DOCTOR.....A		DOCTOR.....A	
	Anyone else ?	NURSE/MIDWIPE.....B		NURSE/MIDWIPE.....B	
	PROBE FOR THE TYPE OF PERSON AND RECORD ALL PERSONS SEEN.	OTHER PERSON		OTHER PERSON	
		TRAD. MIDWIPE.....D		TRAD. MIDWIPE.....D	
		OTHER _____ U		OTHER _____ U	
		(SPECIFY)		(SPECIFY)	
		NO ONE.....Y	<input type="checkbox"/>	NO ONE.....Y	<input type="checkbox"/>
		407C ←		407C ←	

		LAST BIRTH NAME _____	NEXT TO LAST BIRTH NAME _____
--	--	--------------------------	----------------------------------

407B Where did you go for antenatal care ?

RECORD ALL MENTIONED.

(NAME OF PLACE)

(NAME OF PLACE)

PUBLIC SECTOR

GOVT./SAMPLE HOSP.....A
MATERNITY HOUSE.....B
MCHFP CENTER.....C
HEALTH CENTER.....D
HEALTH HOUSE.....E
SSK HOSP/DISPANSERY....F
OTHER PUBLIC SECTOR

(SPECIFY) G

PRIVATE

PRIVATE HOSP.....H
PRIVATE POLYCLINIC....I
PRIVATE DOCTOR.....J
PRIVATE NURSE/MIDWIPE..K
OTHER PRIVATE MEDICAL

(SPECIFY) M

UNIVERSITY HOSPITAL.....N

OTHER _____ U
(SPECIFY)

PUBLIC SECTOR

GOVT./SAMPLE HOSP.....A
MATERNITY HOUSE.....B
MCHFP CENTER.....C
HEALTH CENTER.....D
HEALTH HOUSE.....E
SSK HOSP/DISPANSERY....F
OTHER PUBLIC SECTOR

(SPECIFY) G

PRIVATE

PRIVATE HOSP.....H
PRIVATE POLYCLINIC....I
PRIVATE DOCTOR.....J
PRIVATE NURSE/MIDWIPE..K
OTHER PRIVATE MEDICAL

(SPECIFY) M

UNIVERSITY HOSPITAL.....N

OTHER _____ U
(SPECIFY)

		SKIP TO 408.	SKIP TO 408.
--	--	--------------	--------------

407C During your pregnancy with (NAME) why did not you receive antenatal care ?

RECORD ALL MENTIONED.

NO NEED.....A
ACCESSIBILITY PROBLEMS.....B
DISTRUST OF INSTITUTIONS OR PERSONNEL.....C
PROBLEMS IN USING HEALTH INSTITUTIONS.....D
TRADITIONS ETC.....E
MONETARY REASONS.....F
POOR SERVICE.....G
DON'T KNOW WHERE.....H
OTHER _____ U
(SPECIFY)
DON'T KNOW.....X

NO NEED.....A
ACCESSIBILITY PROBLEMS.....B
DISTRUST OF INSTITUTIONS OR PERSONNEL.....C
PROBLEMS IN USING HEALTH INSTITUTIONS.....D
TRADITIONS ETC.....E
MONETARY REASONS.....F
POOR SERVICE.....G
DON'T KNOW WHERE.....H
OTHER _____ U
(SPECIFY)
DON'T KNOW.....X

407D		SKIP TO 409F.	SKIP TO 409F.
------	--	---------------	---------------

		LAST BIRTH		NEXT TO LAST BIRTH	
		NAME _____		NAME _____	
408	How many months pregnant were you with (NAME) when you first received antenatal care ?	MONTH..... <input type="text"/>	<input type="text"/>	MONTH..... <input type="text"/>	<input type="text"/>
		DON'T KNOW.....98		DON'T KNOW.....98	
409A	During your pregnancy with (NAME) when you go for the first time for antenatal care did you go because there was a problem or was it an ordinary control ?	THERE WAS A PROBLEM.....01		THERE WAS A PROBLEM.....01	
		ORDINARY CONTROL.....02		ORDINARY CONTROL.....02	
		OTHER _____ 96		OTHER _____ 96	
		(SPECIFY)		(SPECIFY)	
		DON'T KNOW.....98		DON'T KNOW.....98	
409B	How many times did you receive antenatal care during your pregnancy with (NAME) ?	NO. OF TIMES..... <input type="text"/>	<input type="text"/>	NO. OF TIMES..... <input type="text"/>	<input type="text"/>
		DON'T KNOW.....98		DON'T KNOW.....98	
409D	In any of your antenatal checks, were you;	YES NO		YES NO	
	Weighted ?	WEIGHTED.....1	2	WEIGHTED.....1	2
	Measured ?	MEASURED.....1	2	MEASURED.....1	2
	Blood pressure measured ?	BLOOD PRESSURE.....1	2	BLOOD PRESSURE.....1	2
	Blood test ?	BLOOD TEST.....1	2	BLOOD TEST.....1	2
	Urine test ?	URINE TEST.....1	2	URINE TEST.....1	2
	Abdomen measured ?	ABDOMEN MEASURED.....1	2	ABDOMEN MEASURED.....1	2
	Listened to baby's heartbeat ?	LISTENED TO BABY.....1	2	LISTENED TO BABY.....1	2
	Ultrasound ?	ULTRASOUND.....1	2	ULTRASOUND.....1	2
	Internal examination ?	INTERNAL EXAMINATION.....1	2	INTERNAL EXAMINATION.....1	2
409E	Have you been given information about the following subjects in any of your antenatal checks ?	YES NO		YES NO	
	Diet ?	DIET.....1	2	DIET.....1	2
	Danger signs of pregnancy ?	DANGER SIGNS.....1	2	DANGER SIGNS.....1	2
	Breastfeeding ?	BREASTFEEDING.....1	2	BREASTFEEDING.....1	2
	Family planning ?	FAMILY PLANNING.....1	2	FAMILY PLANNING.....1	2
	Delivery ?	DELIVERY.....1	2	DELIVERY.....1	2
	Postnatal care ?	POSTNATAL CARE.....1	2	POSTNATAL CARE.....1	2
409F	Have you taken any of the following when you were pregnant with (NAME) ?	YES NO		YES NO	
	Iron tablets ?	IRON TABLETS.....1	2	IRON TABLETS.....1	2
	Vitamins ?	VITAMINS.....1	2	VITAMINS.....1	2
	Folic acid ?	FOLIC ACID.....1	2	FOLIC ACID.....1	2
	Other medication ?	OTHER MEDICATION.....1	2	OTHER MEDICATION.....1	2

		LAST BIRTH	NEXT TO LAST BIRTH
		NAME _____	NAME _____
410	When you were pregnant with (NAME) were you give an injection in the arm to prevent the baby from getting tetanus, that is, convulsions after birth ?	YES.....1 NO.....2 412 ← DON'T KNOW.....8	YES.....1 NO.....2 412 ← DON'T KNOW.....8
411A	During this pregnancy how many times did you get this injection ?	NO. OF TIMES..... <input type="text"/> DON'T KNOW.....8	NO. OF TIMES..... <input type="text"/> DON'T KNOW.....8
411B	In which month of your pregnancy did you get the first tetanus injection ?	MONTH..... <input type="text"/> <input type="text"/> DON'T KNOW.....98	MONTH..... <input type="text"/> <input type="text"/> DON'T KNOW.....98
412	Where did you give birth to (NAME) ? _____ (NAME OF PLACE)	HOME WOMAN'S HOME.....01 OTHER HOME.....02 PUBLIC SECTOR GOVT. /SAMPLE HOSP...11 MATERNITY HOUSE.....12 MCHFP CENTER.....13 HEALTH CENTER.....14 HEALTH HOUSE.....15 SSK HOSP/DISPANSERY...16 OTHER PUBLIC SECTOR _____ 19 (SPECIFY)	HOME WOMAN'S HOME.....01 OTHER HOME.....02 PUBLIC SECTOR GOVT. /SAMPLE HOSP...11 MATERNITY HOUSE.....12 MCHFP CENTER.....13 HEALTH CENTER.....14 HEALTH HOUSE.....15 SSK HOSP/DISPANSERY...16 OTHER PUBLIC SECTOR _____ 19 (SPECIFY)
		PRIVATE SECTOR PRIVATE HOSPITAL.....21 PRIVATE POLYCLINIC...22 PRIVATE DOCTOR.....23 PRIVATE NURSE/MIDWIFE.24 OTHER PRIVATE SECTOR _____ 29 (SPECIFY)	PRIVATE SECTOR PRIVATE HOSPITAL.....21 PRIVATE POLYCLINIC...22 PRIVATE DOCTOR.....23 PRIVATE NURSE/MIDWIFE.24 OTHER PRIVATE SECTOR _____ 29 (SPECIFY)
		UNIVERSITY HOSPITAL....31 OTHER _____ 96 (SPECIFY)	UNIVERSITY HOSPITAL....31 OTHER _____ 96 (SPECIFY)

	LAST BIRTH NAME _____	NEXT TO LAST BIRTH NAME _____
--	--------------------------	----------------------------------

413 Who assisted with the delivery of (NAME)? Anyone else ? PROBE FOR THE TYPE OF PERSON AND RECORD ALL PERSONS ASSISTING.	HEALTH PROFESSIONAL DOCTOR.....A NURSE/MIDWIFE.....B OTHER PERSON TRADITIONAL MIDWIFE....D RELATIVE/FRIENDS.....E OTHER _____ U (SPECIFY) NO ONE.....Y	HEALTH PROFESSIONAL DOCTOR.....A NURSE/MIDWIFE.....B OTHER PERSON TRADITIONAL MIDWIFE....D RELATIVE/FRIENDS.....E OTHER _____ U (SPECIFY) NO ONE.....Y
--	--	--

413A How many months did your pregnancy to (NAME) last ?	MONTH..... <input type="text"/> <input type="text"/>	AY..... <input type="text"/> <input type="text"/>
	AT TIME.....94 BEFORE ITS TIME.....95	AT TIME.....94 BEFORE ITS TIME.....95

413B CHECK 412: BIRTH IN A HEALTH INSTITUTION ?	YES NO <input type="checkbox"/> → 414 <input type="checkbox"/>	YES NO <input type="checkbox"/> → 414 <input type="checkbox"/>
--	---	---

413C What was the main reason for not having done (NAME) birth in a health institution ?	NO REASON.....00 ACCESSIBILITY PROBLEMS...01 DISTRUST OF HEALTH FACILITY OR PERSONNEL.....02 HAPPENED SUDDENLY.....03 PROBLEMS IN USING HEALTH INSTITUTION....04 EXPENSIVE.....05 TRADITIONS ETC.....06 OTHER _____ 96 (SPECIFY) DON'T KNOW.....98	NO REASON.....00 ACCESSIBILITY PROBLEMS...01 DISTRUST OF HEALTH FACILITY OR PERSONNEL.....02 HAPPENED SUDDENLY.....03 PROBLEMS IN USING HEALTH INSTITUTION....04 EXPENSIVE.....05 TRADITIONS ETC.....06 OTHER _____ 96 (SPECIFY) DON'T KNOW.....98
--	---	---

414 Around the time of the birth of (NAME) did you have any of the following problems :	YES NO	YES NO
	LABOUR MORE THAN 12 HOURS...1 2	LABOUR MORE THAN 12 HOURS...1 2
Excessive bleeding that you think was life threatening ?	EXCESSIVE BLEEDING.....1 2	EXCESSIVE BLEEDING.....1 2
A high fever with bad smelling vaginal discharge ?	FEVER/BAD SMELLING VAG. DISCHARGE.....1 2	FEVER/BAD SMELLING VAG. DISCHARGE.....1 2
Convulsions not caused by fever ?	CONVULSIONS.....1 2	CONVULSIONS.....1 2
Episiotomy ?	EPISIOTOMY.....1 2	EPISIOTOMY.....1 2

		LAST BIRTH		NEXT TO LAST BIRTH	
		NAME _____		NAME _____	
414A	CHECK 412: BIRTH IN A HEALTH INSTITUTION ?	YES <input type="checkbox"/>	NO <input type="checkbox"/> → 415B	YES <input type="checkbox"/>	NO <input type="checkbox"/> → 415B
415	Was (NAME) delivered by caesarean section?	YES.....1 NO.....2 415B ←	YES.....1 NO.....2 415B ←	YES.....1 NO.....2 415B ←	YES.....1 NO.....2 415B ←
415A	What was the reason for having a caesarean ?	_____ <input type="checkbox"/>		_____ <input type="checkbox"/>	
415B	Did you have any of the following complications within 40 days period after the birth of (NAME) ?	YES NO		YES NO	
	Excessive (vaginal) bleeding ?	EXCESSIVE BLEEDING....1 2	EXCESSIVE BLEEDING....1 2	EXCESSIVE BLEEDING....1 2	EXCESSIVE BLEEDING....1 2
	Very high fever ?	VERY HIGH FEVER.....1 2	VERY HIGH FEVER.....1 2	VERY HIGH FEVER.....1 2	VERY HIGH FEVER.....1 2
	Convulsions ?	CONVULSIONS.....1 2	CONVULSIONS.....1 2	CONVULSIONS.....1 2	CONVULSIONS.....1 2
	Bad smelling (vaginal) discharge ?	VAGINAL DISCHARGE....1 2	VAGINAL DISCHARGE....1 2	VAGINAL DISCHARGE....1 2	VAGINAL DISCHARGE....1 2
	Lower abdominal pain ?	LOWER ABDOMINAL PAIN..1 2	LOWER ABDOMINAL PAIN..1 2	LOWER ABDOMINAL PAIN..1 2	LOWER ABDOMINAL PAIN..1 2
	Severe back pain ?	SEVERE BACK PAIN.....1 2	SEVERE BACK PAIN.....1 2	SEVERE BACK PAIN.....1 2	SEVERE BACK PAIN.....1 2
	Painful urination ?	PAINFUL URINATION....1 2	PAINFUL URINATION....1 2	PAINFUL URINATION....1 2	PAINFUL URINATION....1 2
	Giddiness ?	GIDDINESS.....1 2	GIDDINESS.....1 2	GIDDINESS.....1 2	GIDDINESS.....1 2
	Severe headache ?	SEVERE HEADACHE.....1 2	SEVERE HEADACHE.....1 2	SEVERE HEADACHE.....1 2	SEVERE HEADACHE.....1 2
	Swollen and painful breast ?	SWOLLEN BREAST.....1 2	SWOLLEN BREAST.....1 2	SWOLLEN BREAST.....1 2	SWOLLEN BREAST.....1 2
415C	CHECK 415B: EXPERIENCED ANY OF THE COMPLICATIONS ?	YES <input type="checkbox"/>	NO <input type="checkbox"/> → 415E	YES <input type="checkbox"/>	NO <input type="checkbox"/> → 416
415D	Did you seek any treatment or advice from a health personnel on any of these complications ? IF YES: From whom ?	HEALTH PERSONNEL DOCTOR.....A NURSE/MIDWIFE.....B OTHER _____ U (SPECIFY) NO ADVICE/ NO TREATMENT.....Y		HEALTH PERSONNEL DOCTOR.....A NURSE/MIDWIFE.....B OTHER _____ U (SPECIFY) NO ADVICE/ NO TREATMENT.....Y	

	LAST BIRTH	NEXT TO LAST BIRTH
	NAME _____	NAME _____

415E	<p>Do you have any of the following complications now ?</p> <p>Uterine prolapse ? IF YES: Did you get treatment for it ?</p> <p>Urinary incontinence ? IF YES: Did you get treatment for it ?</p> <p>Vaginal discharge ? IF YES: Did you get treatment for it ?</p> <p>Urinary infection ? IF YES: Did you get treatment for it ?</p> <p>Menstrual disorder ? IF YES: Did you get treatment for it ?</p>	<p>NO.....0</p> <p>YES, GOT TREATMENT.....1</p> <p>YES, NO TREATMENT.....2</p> <p>NO.....0</p> <p>YES, GOT TREATMENT.....1</p> <p>YES, NO TREATMENT.....2</p> <p>NO.....0</p> <p>YES, GOT TREATMENT.....1</p> <p>YES, NO TREATMENT.....2</p> <p>NO.....0</p> <p>YES, GOT TREATMENT.....1</p> <p>YES, NO TREATMENT.....2</p> <p>NO.....0</p> <p>YES, GOT TREATMENT.....1</p> <p>YES, NO TREATMENT.....2</p>	
------	--	--	--

416	<p>When (NAME) was born, was he/she very large, larger than average, average, smaller than average or very small ?</p>	<p>VERY LARGE.....1</p> <p>LARGER THAN AVERAGE.....2</p> <p>AVERAGE.....3</p> <p>SMALLER THAN AVERAGE.....4</p> <p>VERY SMALL.....5</p> <p>DON'T KNOW.....8</p>	<p>VERY LARGE.....1</p> <p>LARGER THAN AVERAGE.....2</p> <p>AVERAGE.....3</p> <p>SMALLER THAN AVERAGE.....4</p> <p>VERY SMALL.....5</p> <p>DON'T KNOW.....8</p>
-----	--	---	---

417	<p>Was (NAME) weighted at birth ?</p>	<p>YES.....1</p> <p>NO.....2</p> <p style="text-align: center;">419 ←</p>	<p>YES.....1</p> <p>NO.....2</p> <p style="text-align: center;">420 ←</p>
-----	---------------------------------------	---	---

418	<p>How much did (NAME) weigh ?</p> <p>RECORD WEIGHT FROM HEALTH CARD, IF AVAILABLE.</p>	<p style="text-align: center;">GRAMS</p> <p>FROM CARD...1 <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/></p> <p>FROM RECALL.2 <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/></p> <p>DON'T KNOW.....99998</p>	<p style="text-align: center;">GRAMS</p> <p>FROM CARD...1 <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/></p> <p>FROM RECALL.2 <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/></p> <p>DON'T KNOW.....99998</p>
-----	---	--	--

419	<p>Has your period returned since the birth of (NAME) ?</p>	<p>YES.....1</p> <p style="text-align: center;">421 ←</p> <p>NO.....2</p> <p style="text-align: center;">422 ←</p>	
-----	---	--	--

		LAST BIRTH		NEXT TO LAST BIRTH	
		NAME _____		NAME _____	
420	Did your period return between the birth of (NAME) and your next pregnancy ?			YES.....1	NO.....2
				424 ←	
421	For how many months after the birth of (NAME) did you not have a period ?	MONTH..... <input type="text"/> <input type="text"/>	MONTH..... <input type="text"/> <input type="text"/>	DON'T KNOW.....98	DON'T KNOW.....98
422	CHECK 227: RESPONDENT CURRENTLY PREGNANT ?	NOT PREGNANT <input type="checkbox"/>	PREGNANT OR UNSURE <input type="checkbox"/>		
		424			
423	Have you resumed sexual relations since the birth of (NAME) ?			YES.....1	NO.....2
				425 ←	
424	For how many months after the birth of (NAME) did you not have sexual relations ?	MONTH..... <input type="text"/> <input type="text"/>	MONTH..... <input type="text"/> <input type="text"/>	DON'T KNOW.....98	DON'T KNOW.....98
425	Did you ever breastfeed (NAME) ?	YES.....1	YES.....1	NO.....2	NO.....2
		431 ←		431 ←	
426	How long after birth did you first put (NAME) to the breast ? IF LESS THAN 1 HOUR, RECORD '00' HOURS . IF LESS THAN 24 HOURS, RECORD HOURS, OTHERWISE, RECORD DAYS.	IMMEDIATELY.....000	IMMEDIATELY.....000	HOURS.....1 <input type="text"/> <input type="text"/>	HOURS.....1 <input type="text"/> <input type="text"/>
		DAYS.....2 <input type="text"/> <input type="text"/>	DAYS.....2 <input type="text"/> <input type="text"/>		
427	CHECK 404: CHILD ALIVE ?	ALIVE <input type="checkbox"/>	DEAD <input type="checkbox"/>	ALIVE <input type="checkbox"/>	DEAD <input type="checkbox"/>
		429		429	
428	Are you still breastfeeding ?			YES.....1	YES.....1
				432 ←	

		LAST BIRTH NAME _____	NEXT TO LAST BIRTH NAME _____
429	For how many months did you breastfeed (NAME) ?	MONTH..... <input type="text"/> <input type="text"/> DON'T KNOW.....98	MONTH..... <input type="text"/> <input type="text"/> DON'T KNOW.....98
430	Why did you stop breastfeeding (NAME)?	MOTHER ILL/WEAK.....01 CHILD ILL/WEAK.....02 CHILD DIED.....03 NIPPLE/BREAST PROBLEM..04 NOT ENOUGH MILK.....05 MOTHER WORKING.....06 CHILD REPUSED.....07 WEANING AGE/AGE TO STOP.08 BECAME PREGNANT.....09 STARTED USING CONTRACEPTION.....10 OTHER _____ 96 (SPECIFY)	MOTHER ILL/WEAK.....01 CHILD ILL/WEAK.....02 CHILD DIED.....03 NIPPLE/BREAST PROBLEM..04 NOT ENOUGH MILK.....05 MOTHER WORKING.....06 CHILD REPUSED.....07 WEANING AGE/AGE TO STOP.08 BECAME PREGNANT.....09 STARTED USING CONTRACEPTION.....10 OTHER _____ 96 (SPECIFY)
431	CHECK 404: CHILD ALIVE ?	ALIVE <input type="checkbox"/> ↓ 434 DEAD <input type="checkbox"/> ↓ GO BACK TO 405 IN NEXT COLUMN OR, IF NO MORE BIRTHS GO TO 440.	ALIVE <input type="checkbox"/> ↓ 434 DEAD <input type="checkbox"/> ↓ GO BACK TO 405 IN ADDIT. QUEST OR, IF NO MORE BIRTHS GO TO 440.
432	How many times did you breastfeed (NAME) lastnight between sunset and sunrise ? IF ANSWER IS NOT NUMERIC, PROBE FOR APPROXIMATE NUMBER.	NUMBER OF NIGHTTIME FEEDINGS..... <input type="text"/> <input type="text"/>	NUMBER OF NIGHTTIME FEEDINGS..... <input type="text"/> <input type="text"/>
433	How many times did you breastfeed (NAME) yesterday during the daylight hours ? IF ANSWER IS NOT NUMERIC, PROBE FOR APPROXIMATE NUMBER.	NUMBER OF DAYTIME FEEDINGS..... <input type="text"/> <input type="text"/>	NUMBER OF DAYTIME FEEDINGS..... <input type="text"/> <input type="text"/>
434	Did (NAME) drink anything from a bottle with a nipple yesterday or last night ?	YES.....1 NO.....2 DON'T KNOW.....8	YES.....1 NO.....2 DON'T KNOW.....8

		LAST BIRTH	NEXT TO LAST BIRTH
		NAME _____	NAME _____
435	At any time yesterday or last night, was (NAME) given any of the following ?	Y N DK	Y N DK
	Plain water ?	PLAIN WATER.....1 2 8	PLAIN WATER.....1 2 8
	Sugar water ?	SUGAR WATER.....1 2 8	SUGAR WATER.....1 2 8
	Juice ?	JUICE.....1 2 8	JUICE.....1 2 8
	Tea ?	TEA.....1 2 8	TEA.....1 2 8
	Baby formula ?	BABY FORMULA.....1 2 8	BABY FORMULA.....1 2 8
	Yoghurt ?	YOGHURT.....1 2 8	YOGHURT.....1 2 8
	Pudding ?	PUDDING.....1 2 8	PUDDING.....1 2 8
	Juice of cooked meal ?	JUICE OF COOKED ME..1 2 8	JUICE OF COOKED ME..1 2 8
	Turkish delight ?	TURKISH DELIGHT....1 2 8	TURKISH DELIGHT....1 2 8
	Bottled milk ?	BOTTLED MILK.....1 2 8	BOTTLED MILK.....1 2 8
	Fresh milk ?	FRESH MILK.....1 2 8	FRESH MILK.....1 2 8
	Other liquids ?	OTHER LIQUIDS.....1 2 8	OTHER LIQUIDS.....1 2 8
	Any other solid or semi solid foods ?	SOLID/SEMI SOLID FOODS.....1 2 8	SOLID/SEMI SOLID FOOD.....1 2 8
439		GO BACK TO 405 IN NEXT COLUMN; OR, IF NO MORE BIRTHS, GO TO 440.	GO BACK TO 405 IN ADDITIONAL QUESTIONNAIRE; OR IF NO MORE BIRTHS, GO TO 440.

SECTION 4B. IMMUNIZATION AND HEALTH

440 ENTER LINE NUMBER, NAME AND SURVIVAL STATUS OF EACH BIRTH SINCE JANUARY 1993 IN THE TABLE.
ASK THE QUESTIONS ABOUT ALL OF THESE BIRTHS. BEGIN WITH THE LAST BIRTH. (IF THERE ARE MORE THAN 2 BIRTHS USE ADDITIONAL QUESTIONNAIRES- DO NOT USE THE LAST BIRTH COLUMN IN THE ADDITIONAL QUESTIONNAIRE)

	LAST BIRTH	NEXT TO LAST BIRTH
441	LINE NUMBER FROM Q212. LINE NO..... <input type="text"/> <input type="text"/>	LINE NO..... <input type="text"/> <input type="text"/>

442	FROM Q212.	NAME _____	NAME _____
	FROM Q216.	ALIVE <input type="checkbox"/> DEAD <input type="checkbox"/> (GO TO 442 IN NEXT COLUMN; OR, IF NO MORE BIRTHS, GO TO 601)	ALIVE <input type="checkbox"/> DEAD <input type="checkbox"/> (GO TO 442 IN ADDI. QUEST. OR, IF NO MORE BIRTHS, GO TO 601)

443	Do you have a card where (NAME'S) vaccinations are written down ? IF YES: May I see it please ?	YES, SEEN.....1- <input type="checkbox"/> 445 ←	YES, SEEN.....1- <input type="checkbox"/> 445 ←
		YES, NOT SEEN.....2- <input type="checkbox"/> 447 ←	YES, NOT SEEN.....2- <input type="checkbox"/> 447 ←
		NO CARD.....3	NO CARD.....3

444	Did you ever have a vaccination card for (NAME) ?	YES.....1- <input type="checkbox"/> 447 ←	YES.....1- <input type="checkbox"/> 447 ←
		NO.....2- <input type="checkbox"/>	NO.....2- <input type="checkbox"/>

445	(1) COPY VACCINATION DATES FOR EACH VACCINE FROM THE CARD. PAY ATTENTION TO APPOINTMENT DAYS AND THE CONSISTENCY OF VACCINATION DATES. (2) WRITE '44' IN THE DAY COLUMN IF CARD SHOWS THAT A VACCINATION WAS GIVEN BUT NO DATE IS RECORDED.	DAY MO YEAR	DAY MO YEAR
		BCG	BCG
		Polio 1	P1.
		Polio 2	P2.
		Polio 3	P3.
		DPT 1	D1.
		DPT 2	D2.
		DPT 3	D3.
		MEASLES	MEA
		Hepatitis B 1	H1.
		Hepatitis B 2	H2.
		Hepatitis B 3	H3.

		LAST BIRTH	NEXT TO LAST BIRTH
		NAME _____	NAME _____
446	<p>Has (NAME) received any vaccination that are not recorded on this card ?</p> <p>RECORD 'YES' IF ONLY RESPONDENT MENTIONS BCG, POLIO 1-3, DPT 1-3 MEASLES AND/OR HEPATITIS B 1-3.</p>	<p>YES.....1 (PROBE FOR VACCINATIONS ← AND WRITE '66' IN THE CORRESPONDING DAY COLUMN IN 445).....</p> <p>NO.....2 DON'T KNOW.....8 454 ←</p>	<p>YES.....1 (PROBE FOR VACCINATIONS ← AND WRITE '66' IN THE CORRESPONDING DAY COLUMN IN 445).....</p> <p>NO.....2 DON'T KNOW.....8 454 ←</p>
447	<p>Did (NAME) ever receive any vaccinations to prevent him/her from getting infectious diseases ?</p>	<p>YES.....1 NO.....2 454 ← DON'T KNOW.....8</p>	<p>YES.....1 NO.....2 454 ← DON'T KNOW.....8</p>
448	<p>Please tell me if (NAME) received any of the following vaccinations:</p>		
448A	<p>BCG: A vaccination agains tuberculosis, that is an injection in the left arm or shoulder that caused a scar ?</p>	<p>YES.....1 NO.....2 DON'T KNOW.....8</p>	<p>YES.....1 NO.....2 DON'T KNOW.....8</p>
448B	<p>Polio vaccination: That is drops in the mouth ?</p>	<p>YES.....1 NO.....2 448E ← DON'T KNOW.....8</p>	<p>YES.....1 NO.....2 448E ← DON'T KNOW.....8</p>
448C	<p>How many times ?</p>	<p>NUMBER OF TIMES..... <input type="text"/></p>	<p>NUMBER OF TIMES..... <input type="text"/></p>
448E	<p>DPT vaccination: This vaccination includes diphtheria, whooping-cough and tetanus. And it s usually given at the same time as polio drops.</p>	<p>YES.....1 NO.....2 448G ← DON'T KNOW.....8</p>	<p>YES.....1 NO.....2 448G ← DON'T KNOW.....8</p>
448F	<p>How many times ?</p>	<p>NUMBER OF TIMES..... <input type="text"/></p>	<p>NUMBER OF TIMES..... <input type="text"/></p>
448G	<p>Measles vaccination ?</p>	<p>YES.....1 NO.....2 DON'T KNOW.....8</p>	<p>YES.....1 NO.....2 DON'T KNOW.....8</p>
448H	<p>How many times ?</p>	<p>NUMBER OF TIMES..... <input type="text"/></p>	<p>NUMBER OF TIMES..... <input type="text"/></p>
448I	<p>Hepatitis B vaccination ?</p>	<p>YES.....1 NO.....2 DON'T KNOW.....8</p>	<p>YES.....1 NO.....2 DON'T KNOW.....8</p>
448J	<p>How many times ?</p>	<p>NUMBER OF TIMES..... <input type="text"/></p>	<p>NUMBER OF TIMES..... <input type="text"/></p>

		LAST BIRTH	NEXT TO LAST BIRTH																																																																																																				
		NAME _____	NAME _____																																																																																																				
454	Has (NAME) had diarrhea in the last 15 days?	YES.....1 NO.....2 464 ← DON'T KNOW.....8	YES.....1 NO.....2 464 ← DON'T KNOW.....8																																																																																																				
455	Was there any blood in the stools ?	YES.....1 NO.....2 DON'T KNOW.....8	YES.....1 NO.....2 DON'T KNOW.....8																																																																																																				
456	On the worst day of diarrhea, how many bowel movements did (NAME) have ?	NUMBER OF BOWEL MOVEMENTS..... <input type="text"/> <input type="text"/> DON'T KNOW.....98	NUMBER OF BOWEL MOVEMENTS..... <input type="text"/> <input type="text"/> DON'T KNOW.....98																																																																																																				
457	Was (NAME) given the same amount to drink as before the diarrhea, or more or less ?	SAME.....1 MORE.....2 LESS.....3 DON'T KNOW.....8	SAME.....1 MORE.....2 LESS.....3 DON'T KNOW.....8																																																																																																				
458	Was (NAME) given the same amount of food to eat as before the diarrhea, or more or less ?	SAME.....1 MORE.....2 LESS.....3 DON'T KNOW.....8	SAME.....1 MORE.....2 LESS.....3 DON'T KNOW.....8																																																																																																				
459	When (NAME) had diarrhea was he/she given any of the following to drink ?	<table border="0"> <tr> <td></td> <td></td> <td>Y</td> <td>N</td> <td>DK</td> </tr> <tr> <td>A fluid made from a special packet called ORS ?</td> <td>ORS PKT.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Home made sugar-salt-water solution ?</td> <td>HOME MADE ORS.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Milk or infant formula ?</td> <td>MILK/INFANT FORM.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Soup ?</td> <td>SOUP.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Ayran (yoghurt based drink) ?</td> <td>AYRAN.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Water ?</td> <td>WATER.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Tea ?</td> <td>TEA.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Rice-pudding ?</td> <td>RICE PUDDING.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Other liquid ?</td> <td>OTHER LIQUID.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> </table>			Y	N	DK	A fluid made from a special packet called ORS ?	ORS PKT.....	1	2	8	Home made sugar-salt-water solution ?	HOME MADE ORS.....	1	2	8	Milk or infant formula ?	MILK/INFANT FORM.....	1	2	8	Soup ?	SOUP.....	1	2	8	Ayran (yoghurt based drink) ?	AYRAN.....	1	2	8	Water ?	WATER.....	1	2	8	Tea ?	TEA.....	1	2	8	Rice-pudding ?	RICE PUDDING.....	1	2	8	Other liquid ?	OTHER LIQUID.....	1	2	8	<table border="0"> <tr> <td></td> <td></td> <td>Y</td> <td>N</td> <td>DK</td> </tr> <tr> <td>A fluid made from a special packet called ORS ?</td> <td>ORS PKT.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Home made sugar-salt-water solution ?</td> <td>HOME MADE ORS.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Milk or infant formula ?</td> <td>MILK/INFANT FORM.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Soup ?</td> <td>SOUP.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Ayran (yoghurt based drink) ?</td> <td>AYRAN.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Water ?</td> <td>WATER.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Tea ?</td> <td>TEA.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Rice-pudding ?</td> <td>RICE PUDDING.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Other liquid ?</td> <td>OTHER LIQUID.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> </table>			Y	N	DK	A fluid made from a special packet called ORS ?	ORS PKT.....	1	2	8	Home made sugar-salt-water solution ?	HOME MADE ORS.....	1	2	8	Milk or infant formula ?	MILK/INFANT FORM.....	1	2	8	Soup ?	SOUP.....	1	2	8	Ayran (yoghurt based drink) ?	AYRAN.....	1	2	8	Water ?	WATER.....	1	2	8	Tea ?	TEA.....	1	2	8	Rice-pudding ?	RICE PUDDING.....	1	2	8	Other liquid ?	OTHER LIQUID.....	1	2	8
		Y	N	DK																																																																																																			
A fluid made from a special packet called ORS ?	ORS PKT.....	1	2	8																																																																																																			
Home made sugar-salt-water solution ?	HOME MADE ORS.....	1	2	8																																																																																																			
Milk or infant formula ?	MILK/INFANT FORM.....	1	2	8																																																																																																			
Soup ?	SOUP.....	1	2	8																																																																																																			
Ayran (yoghurt based drink) ?	AYRAN.....	1	2	8																																																																																																			
Water ?	WATER.....	1	2	8																																																																																																			
Tea ?	TEA.....	1	2	8																																																																																																			
Rice-pudding ?	RICE PUDDING.....	1	2	8																																																																																																			
Other liquid ?	OTHER LIQUID.....	1	2	8																																																																																																			
		Y	N	DK																																																																																																			
A fluid made from a special packet called ORS ?	ORS PKT.....	1	2	8																																																																																																			
Home made sugar-salt-water solution ?	HOME MADE ORS.....	1	2	8																																																																																																			
Milk or infant formula ?	MILK/INFANT FORM.....	1	2	8																																																																																																			
Soup ?	SOUP.....	1	2	8																																																																																																			
Ayran (yoghurt based drink) ?	AYRAN.....	1	2	8																																																																																																			
Water ?	WATER.....	1	2	8																																																																																																			
Tea ?	TEA.....	1	2	8																																																																																																			
Rice-pudding ?	RICE PUDDING.....	1	2	8																																																																																																			
Other liquid ?	OTHER LIQUID.....	1	2	8																																																																																																			
460	Was anything (else) given to treat the diarrhea ?	YES.....1 NO.....2 462 ← DON'T KNOW.....8	YES.....1 NO.....2 462 ← DON'T KNOW.....8																																																																																																				

	LAST BIRTH	NEXT TO LAST BIRTH
	NAME _____	NAME _____

461	What was given to treat the diarrhea ?	PILL OR SYRUP.....A	PILL OR SYRUP.....A
	Anything else ?	INJECTION.....B	INJECTION.....B
	RECORD ALL MENTIONED.	(I.V) INTRAVENOUS.....C	(I.V) INTRAVENOUS.....C
		HOME REMEDIES/ HERBAL MEDICINES.....D	HOME REMEDIES/ HERBAL MEDICINES.....D
		OTHER _____ U	OTHER _____ U
		(SPECIFY)	(SPECIFY)

462	Did you seek advice or treatment for the diarrhea ?	YES.....1	YES.....1
		NO.....2	NO.....2
		464 ←	464 ←

463	Where did you seek advice or treatment ?	PUBLIC SECTOR	PUBLIC SECTOR
	Anywhere else ?	GOVT./SAMPLE HOSP.....A	GOVT./SAMPLE HOSP.....A
	RECORD ALL MENTIONED.	MATERNITY HOUSE.....B	MATERNITY HOUSE.....B
	_____	MCHFP CENTER.....C	MCHFP CENTER.....C
	(NAME OF PLACE)	HEALTH CENTER.....D	HEALTH CENTER.....D
		HEALTH HOUSE.....E	HEALTH HOUSE.....E
		SSK HOSP/DISPANSERY...F	SSK HOSP/DISPANSERY...F
		OTHER PUBLIC SECTOR	OTHER PUBLIC SECTOR
		_____ G	_____ G
		(SPECIFY)	(SPECIFY)
		PRIVATE SECTOR	PRIVATE SECTOR
		PRIVATE HOSP.....H	PRIVATE HOSP.....H
		PRIVATE CLINIC.....I	PRIVATE CLINIC.....I
		PRIVATE DOCTOR.....J	PRIVATE DOCTOR.....J
		PRIVATE NURSE/MIDWIFE..K	PRIVATE NURSE/MIDWIFE..K
		PHARMACY.....L	PHARMACY.....L
		OTHER PRIVATE SECTOR	OTHER PRIVATE SECTOR
		_____ M	_____ M
		(SPECIFY)	(SPECIFY)
		UNIVERSITY HOSPITAL.....N	UNIVERSITY HOSPITAL.....N
		COMM.VOLUNTEERS/ASSOC./	COMM.VOLUNTEERS/ASSOC./
		FOUNDATIONS.....O	FOUNDATIONS.....O
		RELATIVE/FRIENDS/NEIGHB..R	RELATIVE/FRIENDS/NEIGHB..R
		OTHER _____ U	OTHER _____ U
		(SPECIFY)	(SPECIFY)

464	GO BACK TO 442 IN NEXT COLUMN; OR, IF NO MORE BIRTHS GO TO 601.	GO BACK TO 442 IN ADDITIONAL QUESTIONNAIRE; OR, IF NO MORE BIRTHS GO TO 601.
-----	---	--

SECTION 6. FERTILITY PREFERENCES

601	CHECK 250:	CURRENTLY MARRIED <input type="checkbox"/>	NOT CURRENTLY MARRIED <input type="checkbox"/>	→612
601A	CHECK 314:	WOMAN NOT HAD TUBAL LIGATION OR HUSBAND NOT STERILIZED <input type="checkbox"/>	WOMAN HAD TUBAL LIGATION OR HUSBAND STERILIZED <input type="checkbox"/>	→612

602	CHECK 227:	NOT PREGNANT OR UNSURE <input type="checkbox"/> Now I have some questions about the future. Would you like to have (a/another) child or would you prefer not to have any (more) children ?	CURRENTLY PREGNANT <input type="checkbox"/> Now I have some questions about the future. After the child you are expecting, would you like to have another child or would you prefer not to have any more children ?	HAVE (A/ANOTHER) CHILD.....1 NO MORE/NONE.....2 →604 SAYS SHE CAN'T GET PREGNANT.....3 →606 UNDECIDED/DON'T KNOW.....8 →604
-----	------------	---	--	--

602A	CHECK 227:	NOT PREGNANT OR UNSURE <input type="checkbox"/> How many more children would you like to have in the future ?	CURRENTLY PREGNANT <input type="checkbox"/> How many more children would you like to have in the future, not counting the one you are currently pregnant with ?	NUMBER..... <input style="width: 40px;" type="text"/> OTHER ANSWERS _____ 96 (SPECIFY)
------	------------	--	--	--

603	CHECK 227:	NOT PREGNANT OR UNSURE <input type="checkbox"/> How long would you like to wait from now before the birth of (a/another) child ?	CURRENTLY PREGNANT <input type="checkbox"/> After the child you are expecting now, how long would you like to wait before the birth of another child ?	MONTHS.....1 <input style="width: 40px;" type="text"/> YEARS.....2 <input style="width: 40px;" type="text"/> SOON/NOW.....993 →606 SAYS SHE CAN'T GET PREGNANT.....994 OTHER _____ 996 (SPECIFY) DON'T KNOW.....998
-----	------------	---	---	---

604	CHECK 227:	NOT PREGNANT OR UNSURE <input type="checkbox"/>	CURRENTLY PREGNANT <input type="checkbox"/>	→607
-----	------------	--	--	------

605	If you become pregnant in the next few weeks, would you be happy, unhappy, or would it not matter very much ?	HAPPY.....1 UNHAPPY.....2 WOULD NOT MATTER.....3
-----	---	--

606	CHECK 313: USING A METHOD ?	NOT ASKED <input type="checkbox"/>	NOT CURRENTLY USING <input type="checkbox"/>	CURRENTLY USING <input type="checkbox"/>	→612
-----	-----------------------------	------------------------------------	--	---	------

607	Do you think you will use a method to delay or avoid pregnancy within the next 12 months ?	YES.....1 →609 NO.....2 DON'T KNOW.....8
-----	--	--

608	Do you think you will use a method to delay or avoid pregnancy at any time in the future ?	YES.....1 NO.....2 DON'T KNOW.....8 →610
-----	--	--

609	Which method would you prefer to use ?	PILL.....01 IUD.....02 INJECTABLES.....03 NORPLANT.....04 DIAPHRAGM/FOAM/JELLY.....05 CONDOM.....06 TUBAL LIGATION.....07 MALE STERILIZATION.....08 →612 RHYTHM.....09 WITHDRAWAL.....10 OTHER _____ 96 (SPECIFY) UNSURE.....98
-----	--	---

610

What is the main reason that you think you will never use a method ?

PERTILITY-RELATED REASONS

- INFREQUENT SEX.....22
- MENOPAUSAL/HYSTERECTOMY.....23
- SUBFECUND/INFECUND.....24
- WANTS MORE CHILDREN.....26

OPPOSITION TO USE

- WOMAN OPPOSED.....31
- HUSBAND OPPOSED.....32
- OTHERS OPPOSED.....33
- RELIGIOUS REASONS.....34

LACK OF KNOWLEDGE

- KNOWS NO METHOD.....41
- KNOWS NO SOURCE.....42

METHOD-RELATED REASONS

- HEALTH CONCERNS.....51
- SIDE EFFECTS.....52
- LACK OF ACCESS/TOO PAR.....53
- COST TOO MUCH.....54
- INCONVENIENT TO USE.....55
- INTERFERES WITH BODY'S NORMAL PROCESSES.....56

OTHER _____ 96

(SPECIFY)

DON'T KNOW.....98

612

CHECK 216:

HAS LIVING CHILDREN

NO LIVING CHILDREN

If you could go back to the time you did not have any children and could choose exactly the number of children to have in your whole life, how many would that be ?

If you could choose exactly the number of children to have in your whole life, how many would that be ?

PROBE FOR A NUMERIC RESPONSE.

NUMBER.....

OTHER _____ 96

(SPECIFY)

→614

613

How many of these children would you like to be boys, how many would you like to be girls and for how many would it not matter ?

BOYS

NUMBER.....

OTHER _____ 96

(SPECIFY)

GIRLS

NUMBER.....

OTHER _____ 96

(SPECIFY)

EITHER

NUMBER.....

OTHER _____ 96

(SPECIFY)

614	Do you approve of couples using a method to avoid getting pregnant ?	APPROVE.....1 DISAPPROVE...2 NO OPINION...3																								
615	Is it acceptable or not acceptable to you for information on family planning to be provided:	<table border="1"> <thead> <tr> <th></th> <th>ACCEPT- ABLE</th> <th>NOT ACCEPT- ABLE</th> <th>DO NOT KNOW</th> </tr> </thead> <tbody> <tr> <td>On the radio ?</td> <td>RADIO.....1</td> <td>2</td> <td>8</td> </tr> <tr> <td>On the television ?</td> <td>TELEVISION....1</td> <td>2</td> <td>8</td> </tr> <tr> <td>In newspapers/magazines ?</td> <td>NEWSP./MAGZ....1</td> <td>2</td> <td>8</td> </tr> <tr> <td>In secondary schools ?</td> <td>SECONDARY.....1</td> <td>2</td> <td>8</td> </tr> <tr> <td>In high schools ?</td> <td>HIGH.....1</td> <td>2</td> <td>8</td> </tr> </tbody> </table>		ACCEPT- ABLE	NOT ACCEPT- ABLE	DO NOT KNOW	On the radio ?	RADIO.....1	2	8	On the television ?	TELEVISION....1	2	8	In newspapers/magazines ?	NEWSP./MAGZ....1	2	8	In secondary schools ?	SECONDARY.....1	2	8	In high schools ?	HIGH.....1	2	8
	ACCEPT- ABLE	NOT ACCEPT- ABLE	DO NOT KNOW																							
On the radio ?	RADIO.....1	2	8																							
On the television ?	TELEVISION....1	2	8																							
In newspapers/magazines ?	NEWSP./MAGZ....1	2	8																							
In secondary schools ?	SECONDARY.....1	2	8																							
In high schools ?	HIGH.....1	2	8																							
616	In the last few months have you heard about family planning:	<table border="1"> <thead> <tr> <th></th> <th>YES</th> <th>NO</th> </tr> </thead> <tbody> <tr> <td>On the radio ?</td> <td>RADIO.....1</td> <td>2</td> </tr> <tr> <td>On the television ?</td> <td>TELEVISION....1</td> <td>2</td> </tr> <tr> <td>In newspaper or magazine?</td> <td>NEWSPAPER OR MAGAZINE.....1</td> <td>2</td> </tr> <tr> <td>From a poster ?</td> <td>POSTER.....1</td> <td>2</td> </tr> <tr> <td>From brochures or leaflets ?</td> <td>BROCHURES/LEAFLETS.....1</td> <td>2</td> </tr> <tr> <td>From billboards or cloth posters ?</td> <td>BILLBOARDS/CLOTH POSTERS....1</td> <td>2</td> </tr> <tr> <td>Any other place ?</td> <td>ANOTHER PLACE.....1</td> <td>2</td> </tr> </tbody> </table>		YES	NO	On the radio ?	RADIO.....1	2	On the television ?	TELEVISION....1	2	In newspaper or magazine?	NEWSPAPER OR MAGAZINE.....1	2	From a poster ?	POSTER.....1	2	From brochures or leaflets ?	BROCHURES/LEAFLETS.....1	2	From billboards or cloth posters ?	BILLBOARDS/CLOTH POSTERS....1	2	Any other place ?	ANOTHER PLACE.....1	2
	YES	NO																								
On the radio ?	RADIO.....1	2																								
On the television ?	TELEVISION....1	2																								
In newspaper or magazine?	NEWSPAPER OR MAGAZINE.....1	2																								
From a poster ?	POSTER.....1	2																								
From brochures or leaflets ?	BROCHURES/LEAFLETS.....1	2																								
From billboards or cloth posters ?	BILLBOARDS/CLOTH POSTERS....1	2																								
Any other place ?	ANOTHER PLACE.....1	2																								
618	In the last few months have you discussed the practice of family planning with friends, neighbours, or relatives ?	YES.....1 NO.....2 → 620																								
619	With whom ? Anyone else ? RECORD ALL MENTIONED.	HUSBAND/PARTNER.....A MOTHER.....B FATHER.....C SISTER(S).....D BROTHER(S).....E DAUGHTER.....F SON.....G MOTHER-IN-LAW.....H FRIEND/RELATIVE.....I NEIGHBOUR.....J OTHER _____ U (SPECIFY)																								
620	CHECK 250: CURRENTLY <input type="checkbox"/> MARRIED NOT CURRENTLY <input type="checkbox"/> MARRIED	→ 701																								
621	Spouses/partners do not always agree on everything. Now I want to ask you about your husband's/partner's views on family planning. Do you think that your husband/partner approves of couples using a method to avoid pregnancy ?	APPROVES.....1 DISAPPROVES...2 DON'T KNOW...8																								

622B	CHECK 313 AND 314: USING A METHOD ?	
	CURRENTLY USING A METHOD	CURRENTLY NOT USING A METHOD/NOT ASKED
	<input type="checkbox"/>	<input type="checkbox"/> → 623

622C	Before you started to use (METHOD), did you or your husband/partner talk about which method you are going to use ?	YES.....1 NO.....2 DOESN'T REMEMBER.....8
------	--	---

622D	After you started to use (METHOD) did you and your husband/partner talk about the method ?	YES.....1 NO.....2 DOESN'T REMEMBER.....8
------	--	---

622E	CHECK 314:	
	CIRCLE METHOD CODE:	
		PILL.....01 IUD.....02 INJECTABLES.....03 NORPLANT.....04 DIAPHRAGM/FOAM/JELLY.....05 CONDOM.....06 → 623 TUBAL LIGATION.....07 MALE STERILIZATION.....08 → 623 RHYTHM.....09 WITHDRAWAL.....10 → 623 OTHER METHOD.....96

622F	Did your husband/partner support you about the use of (METHOD) ?	SUPPORTED.....1 DIDN'T SUPPORT.....2 INDIFFERENT/UNINTERESTED.....3 DON'T KNOW.....8
------	--	---

623	Do you think your husband/partner wants the same number of children that you want, or does he want more or fewer than you want ?	SAME NUMBER.....1 MORE CHILDREN.....2 FEWER CHILDREN.....3 DON'T KNOW.....8
-----	--	--

628A	Do you think that the use of family planning is against religion ?	YES.....1 → 628c SOME METHODS ARE.....2 NO.....3 NOT RELIGIOUS/NO RELIGION.....4 → 628c DON'T KNOW.....8
------	--	--

628B	<p>Which method(s) do you think (are) is against religion ?</p> <p>Any other ?</p> <p>RECORD ALL MENTIONED.</p>	<p>PILL.....A</p> <p>IUD.....B</p> <p>INJECTABLES.....C</p> <p>NORPLANT.....D</p> <p>DIAPHRAGM/FOAM/JELLY.....E</p> <p>CONDOM.....F</p> <p>TUBAL LIGATION.....G</p> <p>MALE STERILIZATION.....H</p> <p>RHYTHM.....I</p> <p>WITHDRAWAL.....J</p> <p>OTHER METHOD _____ U (SPECIFY)</p> <p>INDUCED ABORTION.....M</p>
628C	<p>Does your husband have any objections to any family planning method or to family planning in general on religious grounds ?</p>	<p>YES.....1</p> <p>THINKS SOME METHODS ARE AGAINST RELIGION.....2</p> <p>NO.....3</p> <p>DON'T KNOW.....8</p>
630	<p>Did you ever reinforce or try to persuade any of your friends or relatives for the use of family planning methods ?</p>	<p>YES.....1</p> <p>NO.....2</p>

SECTION 7A. HUSBAND'S BACKGROUND

701	CHECK 205 : CURRENTLY MARRIED <input type="checkbox"/> CURRENTLY NOT MARRIED <input type="checkbox"/>	→703
702	How old is your husband ?	AGE..... <input type="text"/>
703	Did your (last) husband ever attend school ?	YES.....1 NO.....2 →706A
704	What was the highest level of school he attended ?	PRIMARY1 SECONDARY.....2 HIGH.....3 UNIVERSITY.....4 DON'T KNOW.....8 →706A
705	What was the highest grade he completed at that level ?	GRADE..... <input type="text"/> DON'T KNOW.....8
705A	Did he graduate from that school ?	YES.....1 NO.....2 DON'T KNOW.....8
706A	What is/was your (last) husband's occupation ? That is, what kind of work does/did he mainly do ?	<input type="text"/> <hr/> <hr/> <hr/>
706B	Did/does he pay social security when doing this job ? IF YES : According to which schedule ?	NO.....0 SSK.....1 EMEKLİ SANDIĞI.....2 BAĞ-KUR.....3 OTHER _____ 7 (SPECIFY) DON'T KNOW.....8

706C	CHECK 706A, CIRCLE THE APPROPRIATE WORK CODE	AGRICULTURE.....01 ANIMAL HUSBANDRY.....02 AGRICULTURE AND A.HUSBANDRY...03 FORESTRY.....04 FISHERY.....05 MINING.....06 MANUFACTURE INDUSTRY (FOOD)...07 MANUFACTURE INDUSTRY (TEXTILE).08 MANUFACTURE INDUSTRY (METAL)...09 MANUFACTURE INDUSTRY (OTHER)...10 ELECTRICITY/GAS/WATER.....11 CONSTRUCTION.....12 WHOLESALE/RETAIL TRADE.....13 HOTEL AND RESTAURANT.....14 TRANSPORTATION/COMMUNICATION..15 HEALTH.....16 EDUCATION/CULTURE.....17 PUBLIC ADMINISTRATION.....18 TOURISM.....19 BANKING/INSURANCE.....20 OTHER SOCIAL SERVICES.....21 PERSONAL SERVICES.....22 OTHER _____ 96 (SPECIFY)
------	--	---

706D	CHECK 706A, CIRCLE THE APPROPRIATE CODE FOR THE POSITION AT WORK	EMPLOYER EMPLOYING TEN OR MORE THAN TEN PERSONS.....01 SMALL EMPLOYER EMPLOYING LESS THAN TEN PERSONS.....02 SALARIED (GOVERNMENT OFFICIAL).03 WAGED (PUBLIC SECTOR).....04 WAGED (PRIVATE SECTOR).....05 FOR HIS OWN.....06 UNPAID FAMILY LABOURER.....07 OTHER _____ 96 (SPECIFY)
------	--	---

706E	CHECK 706A :	
	AGRICULTURE (01) OR AGRICULTURE AND ANIMAL HUSBANDRY (03) <input type="checkbox"/>	OTHER WORK CODES <input type="checkbox"/> → 706G

706F	Whom the land that your (last) husband works/worked on belongs/belonged to ?	HIS LAND.....01 FAMILY LAND.....02 RENTED LAND.....03 SOMEONE ELSE'S LAND.....04 STATE LAND.....05 OTHER _____ 96 (SPECIFY)
------	--	---

706G	<p>Is/was your (last) husband covered by health insurance ? Does/did he have health insurance ?</p> <p>IF YES : According to which schedule ?</p>	<p>NO.....0</p> <p>SSK.....1</p> <p>EMEKLİ SANDIĞI.....2</p> <p>BAĞ-KUR.....3</p> <p>PRIVATE INSURANCE.....4</p> <p>GREEN CARD.....5</p> <p>OTHER _____ 7 (SPECIFY)</p> <p>DON'T KNOW.....8</p>
------	---	---

708A	<p>What is/was your (last) husband's mother tongue ?</p> <p>RECORD ONLY ONE RESPONSE.</p>	<p>TURKISH.....01</p> <p>KURDISH AND DIALECTS (KURMANCI, GORANI, ZAZACA ETC).02</p> <p>ARABIC.....03</p> <p>GREEK.....04</p> <p>ARMENIAN.....05</p> <p>HEBREW(LADINO).....06</p> <p>CIRCASSIAN.....07</p> <p>GEORGIAN.....08</p> <p>LAZ LANGUAGE.....09</p> <p>PERSIAN.....10</p> <p>BULGARIAN.....11</p> <p>RUMANIAN.....12</p> <p>SERBIAN.....13</p> <p>ENGLISH.....14</p> <p>GERMAN.....15</p> <p>OTHER _____ 96 (SPECIFY)</p>
------	---	---

708B	<p>In addition to his mother tongue, which language(s) does/did your (last) husband speak ?</p> <p>RECORD ALL MENTIONED.</p>	<p>TURKISH.....A</p> <p>KURDISH AND ITS DIALECTS (KURMANCI, GORANI, ZAZACA ETC).B</p> <p>ARABIC.....C</p> <p>GREEK.....D</p> <p>ARMENIAN.....E</p> <p>HEBREW(LADINO).....F</p> <p>CIRCASSIAN.....G</p> <p>GEORGIAN.....H</p> <p>LAZ LANGUAGE.....I</p> <p>PERSIAN.....J</p> <p>BULGARIAN.....K</p> <p>RUMANIAN.....L</p> <p>SERBIAN.....M</p> <p>ENGLISH.....N</p> <p>GERMAN.....O</p> <p>OTHER _____ U (SPECIFY)</p> <p>KNOWS NO OTHER LANGUAGE.....Y</p>
------	--	--

708C	What is (was) the mother tongues of your (last) husband's mother and father ? USE THE CODES IN 708A.	HUSBAND'S MOTHER..... HUSBAND'S FATHER.....
------	---	--

708D	What is/was your (last) husband's religion ? IF THE RESPONSE IS "MUSLIM" THEN PROBE FOR SECT. CODE ACCORDING TO THE GIVEN RESPONSE.	MUSLIM SUNNI.....01 ALAWI.....02 OTHER _____ 03 (SPECIFY) CHRISTIAN06 JEWISH.....07 NO RELIGION.....10 OTHER _____ 96 (SPECIFY)
------	--	--

708E	Have you lived in only one or in more than one settlements since January 1993 ?	ONE SETTLEMENT.....1 MORE THAN ONE SETTLEMENT.....2 → 708G
------	---	---

708F

C ENTER (IN COLUMN 4 OF CALENDAR) THE APPROPRIATE CODE FOR SETTLEMENT ('1' PROVINCE CENTER, '2' DISTRICT CENTER, '3' SUB-DISTRICT/VILLAGE, '4' ABROAD).


ENTER (IN COLUMN 5 OF THE CALENDAR) THE PROVINCE CODE FOR THE SETTLEMENT.

BEGIN IN THE MONTH OF INTERVIEW AND CONTINUE WITH ALL PRECEDING MONTHS BACK TO JANUARY 1993 BY FILLING COLUMNS 4 AND 5.

THEN SKIP TO 709.

708G	In what month and year did you move to (NAME OF THE SETTLEMENT OF CURRENT RESIDENCE) ? C ENTER (IN COLUMNS 4 AND 5 OF THE CALENDAR) "X" IN THE MONTH AND YEAR OF THE MOVE. FOR COLUMN 4: FOR THE FOLLOWING MONTHS ENTER THE APPROPRIATE CODE FOR SETTLEMENT. ('1' PROVINCE CENTER, '2' DISTRICT CENTER, '3' SUB-DISTRICT/VILLAGE, '4' ABROAD). FOR COLUMN 5: FOR THE FOLLOWING MONTHS ENTER THE PROVINCE CODE THAT THE SETTLEMENT IS CURRENTLY LOCATED. CONTINUE PROBING FOR PREVIOUS SETTLEMENTS AND RECORD MOVES, TYPES OF SETTLEMENTS AND PROVINCE CODES ACCORDINGLY. RECORD ALSO THE MOVES BETWEEN THE SAME TYPE OF SETTLEMENTS OR BETWEEN THE TYPES OF SETTLEMENTS IN THE SAME PROVINCE. ILLUSTRATIVE QUESTIONS: > Where did you live before..... ? > In what month and year did you arrive here ? > Is that place a province center, a district center, a subdistrict/village or abroad? > In which province islocated ?
------	---

SECTION 7B. WOMAN'S WORK AND STATUS

709	<p>Now I would like to ask you questions about working.</p> <p>Aside from your own housework, are you currently working?</p>	<p>YES.....1 → 712</p> <p>NO.....2</p>	
710	<p>As you know, some women sell small things, sell goods at the market place, work on the family farm or business, look after children, work as cleaning ladies etc. Are you doing any of these at the moment, or any other work of similar nature?</p>	<p>YES.....1 → 712</p> <p>NO.....2</p>	
711	<p>Have you worked in any job in the last 12 months ?</p>	<p>YES.....1</p> <p>NO.....2 → 740</p>	
712	<p>What type of work are/were you doing ? What kind of job are/were you in ?</p>	<p>_____</p> <p>_____</p>	
712A	<p>Do/did you pay social security when doing this job ?</p> <p>IF YES: According to which schedule ?</p>	<p>NO.....0</p> <p>SSK.....1</p> <p>EMEKLİ SANDIĞI.....2</p> <p>BAĞ-KUR.....3</p> <p>OTHER _____ 7</p> <p>(SPECIFY)</p> <p>DON'T KNOW.....8</p>	

712B	CHECK 712, CIRCLE THE APPROPRIATE WORK CODE.	AGRICULTURE.....01 ANIMAL HUSBANDRY.....02 AGRICULTURE AND A.HUSBANDRY...03 FORESTRY.....04 FISHERY.....05 MINING.....06 MANUFACTURE INDUSTRY (FOOD)...07 MANUFACTURE INDUSTRY (TEXTILE).08 MANUFACTURE INDUSTRY (METAL)...09 MANUFACTURE INDUSTRY (OTHER)...10 ELECTRICITY/GAS/WATER.....11 CONSTRUCTION.....12 WHOLESALE/RETAIL TRADE.....13 HOTEL AND RESTAURANT.....14 COMMUNICATION/TRANSPORTATION...15 HEALTH.....16 EDUCATION/CULTURE.....17 PUBLIC ADMINISTRATION.....18 TOURISM.....19 BANKING/INSURANCE.....20 OTHER SOCIAL SERVICES.....21 PERSONAL SERVICES.....22 OTHER _____ 96 (SPECIFY)
------	--	--

712C	CHECK 712, CIRCLE THE APPROPRIATE CODE FOR THE POSITION AT WORK	EMPLOYER EMPLOYING TEN OR MORE THAN TEN PERSONS.....01 SMALL EMPLOYER EMPLOYING LESS THAN TEN PERSONS.....02 SALARIED (GOVERNMENT OFFICIAL).03 WAGED (PUBLIC SECTOR).....04 WAGED (PRIVATE SECTOR).....05 FOR HIS OWN.....06 UNPAID FAMILY LABOURER.....07 OTHER _____ 96 (SPECIFY)
------	---	---

713	CHECK 712 : AGRICULTURE (01) OR ARRICULTURE AND ANIMAL HUSBANDRY (03)	OTHER WORK CODES
	<input type="checkbox"/>	<input type="checkbox"/> → 716

714	Whom does the land you work on belong to ?	HER LAND.....01 FAMILY LAND.....02 RENTED LAND.....03 SOMEONE ELSE'S LAND.....04 STATE LAND.....05 OTHER _____ 96 (SPECIFY)
-----	--	---

716	Do you usually work throughout the year, or do you work seasonally, or only once in a while ?	THROUGHOUT THE YEAR.....1 →718 SEASONALLY/PART OF THE YEAR....2 ONCE IN A WHILE.....3 →719
717	During the last 12 months, how many months did you work ?	NUMBER OF MONTHS..... <input type="text"/>
718	How many days a week do (did) you usually work (in the months that you worked) ?	NUMBER OF DAYS..... <input type="text"/> →720
719	During the last 12 months, approximately, how many days did you work ?	NUMBER OF DAYS..... <input type="text"/>
720	Do/did you earn cash for your work ? PROBE: Do/did you make money for your work ?	YES.....1 NO.....2 →723
721A	How much do/did you usually earn for this work ? PROBE: Is this by the day, by the week, or by the month ?	PER HOUR.....1 <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> PER DAY.....2 <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> PER WEEK.....3 <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> PER MONTH.....4 <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> PER YEAR.....5 <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> OTHER _____ 999996 →722 (SPECIFY)
721B	CIRCLE THE APPROPRIATE CODE FOR THE GIVEN AMOUNT IN 721A .	MILLION.....1 THOUSAND.....2
721C	(In the times that you work) Generally what amount of the family expenses is met by your earnings? The whole, more than the half, the half, less than the half, or does your earning have no contribution to the family expenses ?	THE WHOLE.....1 MORE THAN THE HALP.....2 THE HALP.....3 LESS THAN THE HALP.....4 NO CONTRIBUTION.....5 DON'T KNOW.....8

722 CHECK 250 :

<p>CURRENTLY MARRIED <input type="checkbox"/></p> <p>Who mainly decides how the money you earn will be used: you, your husband, you and your husband jointly, or someone else ?</p>	<p>CURRENTLY NOT MARRIED <input type="checkbox"/></p> <p>Who mainly decides how the money you earn will be used: you, someone else, or you and someone else jointly ?</p>
<p>HERSELF DECIDES.....1 HUSBAND DECIDES.....2 JOINTLY WITH HUSBAND.....3 SOMEONE ELSE DECIDES.....4 JOINTLY WITH SOMEONE ELSE.....5</p>	

723 Do you usually work at home or away from home ?

HOME.....1
AWAY.....2

724 CHECK 217 AND 218: IS A CHILD LIVING WITH HER WHOSE AGE IS 5 OR LESS ?

YES NO →740

725 Who usually takes care of (NAME OF THE YOUNGEST CHILD AT HOME) while you are working ?

<p>WOMAN.....01 HUSBAND.....02 OLDER FEMALE CHILD.....03 OLDER MALE CHILD.....04 OTHER RELATIVES.....05 NEIGHBOURS.....06 FRIENDS.....07 SERVANTS/HIRED HELP.....08 INSTITUTIONAL CHILDCARE.....10 HAS NOT WORKED SINCE LAST BIRTH.....95 OTHER _____ 96 (SPECIFY)</p>

740 CURRENTLY NOT WORKING (709=2 AND 710=2) OR CURRENTLY WORKING (709=1 OR 710=1) OR WORKED DURING THE LAST 12 MONTHS (711=1) →742

741 You told that currently you are not working. What is the main reason that you are not working ?

<p>HOLIDAY/ON VACATION.....01 LOOKS AFTER CHILD(REN).....02 HOUSEWORK.....03 SICK/HANDICAPPED.....04 CAN'T FIND/LOOKING FOR JOB.....05 HUSBAND/ELDERS DON'T WANT.....06 NO NEED FOR WORKING.....07 DOES NOT WANT WORKING ANYMORE..08 NO TALENT/EDUCATION.....09 DISCHARGED.....10 OTHER _____ 96 (SPECIFY)</p>

741B	CHECK 711: WORKED DURING THE LAST 12 MONTHS	NOT WORKED DURING THE LAST 12 MONTHS	
	<input type="checkbox"/>	<input type="checkbox"/>	→746

742	How long have/had you been working in your current/last job ?	YEAR.....	<input type="text"/> <input type="text"/>
-----	---	-----------	---

746	Have you ever worked before you got married ?	YES.....1 NO.....2	→757
-----	---	-----------------------	------

747	What was the last job you worked before you got married ? What type of work were you doing ?	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>	
-----	---	---	--

748	Did you pay social security when doing this job ?	NO.....0 SSK.....1 EMEKLİ SANDIĞI.....2 BAĞ-KUR.....3 OTHER _____ 7 (SPECIFY) DON'T KNOW.....8	
	IF YES: According to which schedule ?		

749	CHECK 747 , CIRCLE THE APPROPRIATE WORK CODE	AGRICULTURE.....01 ANIMAL HUSBANDRY.....02 AGRICULTURE AND A.HUSBANDRY...03 FORESTRY.....04 FISHERY.....05 MINING.....06 MANUFACTURE INDUSTRY (FOOD)...07 MANUFACTURE INDUSTRY (TEXTILE).08 MANUFACTURE INDUSTRY (METAL)...09 MANUFACTURE INDUSTRY (OTHER)...10 ELECTRICITY/GAS/WATER.....11 CONSTRUCTION.....12 WHOLESALE/RETAIL TRADE.....13 HOTEL AND RESTAURANT.....14 COMMUNICATION/TRANSPORTATION..15 HEALTH.....16 EDUCATION/CULTURE.....17 PUBLIC ADMINISTRATION.....18 TOURISM.....19 BANKING/INSURANCE.....20 OTHER SOCIAL SERVICES.....21 PERSONAL SERVICES.....22 OTHER _____ 96 (SPECIFY)	
-----	--	---	--

750	CHECK 747. CIRCLE THE APPROPRIATE CODE FOR THE POSITION AT WORK	EMPLOYER EMPLOYING TEN OR MORE THAN TEN PERSONS.....01 SMALL EMPLOYER EMPLOYING LESS THAN TEN PERSONS.....02 SALARIED (GOVERNMENT OFFICIAL).03 WAGED (PUBLIC SECTOR).....04 WAGED (PRIVATE SECTOR).....05 FOR HIS OWN.....06 UNPAID FAMILY LABOURER.....07 OTHER _____ 96 (SPECIFY)
-----	---	---

751	Were you working at the time you got married ?	YES.....1 NO.....2 → 757
-----	--	-----------------------------

752	After you got married, did you go on working in the same job, or start to work in another job, or did you stop working ?	YES, THE SAME JOB.....1 YES, DIFFERENT JOB.....2 → 757 STOPPED WORKING.....3
-----	--	--

753	What was the main reason that you stop working ?	GOT PREGNANT/CHILD CARE.....01 HOUSEWORK.....02 SICK/HANDICAPPED.....03 COULDN'T FIND JOB.....04 MOVED/MIGRATED.....05 HUSBAND/ELDERS DIDN'T WANT.....06 NO NEED FOR WORKING.....07 DIDN'T WANT TO WORK.....08 WORKED UNPAID.....09 DISCHARGED.....10 OTHER _____ 96 (SPECIFY)
-----	--	---

757	What is the main source of income providing your and your family's subsistence?	HUSBAND'S EARNINGS.....01 HER AND HER HUSBAND'S JOINTLY..02 HER EARNINGS.....03 HUSBAND'S FAMILY'S INCOME.....04 HER FAMILY'S INCOME.....05 ALIMONY/SUPPORT FOR CHILD.....06 HER PENSION.....07 HUSBAND'S PENSION.....08 OTHER _____ 96 (SPECIFY)
-----	---	--

758	<p>Are you covered by any health insurance ?</p> <p>IF YES : According to which schedule ?</p>	<p>NO.....0</p> <p>SSK.....1</p> <p>EMEKLİ SANDIĞI.....2</p> <p>BAG-KUR.....3</p> <p>PRIVATE INSURANCE.....4</p> <p>GREEN CARD.....5</p> <p>OTHER _____ 7</p> <p>(SPECIFY)</p> <p>DON'T KNOW.....8</p>
-----	--	--

766	<p>Woman can sometimes make things that may annoy or make angry her husband. Now I will talk about some situations. Can you please tell me if the husband can have the right for beating his wife in those situations?</p> <p>If she burns the food ?</p> <p>If she neglects the care of children ?</p> <p>If she argues with her husband ?</p> <p>If she talks with other men ?</p> <p>If she spends the money needlessly ?</p> <p>If she refuses to have sexual intercourse ?</p>	<table style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th style="text-align: center;">YES</th> <th style="text-align: center;">NO</th> <th style="text-align: center;">DK</th> </tr> </thead> <tbody> <tr> <td>BURN FOOD.....1</td> <td style="text-align: center;">2</td> <td></td> <td style="text-align: center;">8</td> </tr> <tr> <td>NEGLECT CHILD CARE....1</td> <td style="text-align: center;">2</td> <td></td> <td style="text-align: center;">8</td> </tr> <tr> <td>ARGUE WITH HUSBAND....1</td> <td style="text-align: center;">2</td> <td></td> <td style="text-align: center;">8</td> </tr> <tr> <td>TALK OTHER MEN.....1</td> <td style="text-align: center;">2</td> <td></td> <td style="text-align: center;">8</td> </tr> <tr> <td>SPEND NEEDLESSLY.....1</td> <td style="text-align: center;">2</td> <td></td> <td style="text-align: center;">8</td> </tr> <tr> <td>REPUSE INTERCOURSE....1</td> <td style="text-align: center;">2</td> <td></td> <td style="text-align: center;">8</td> </tr> </tbody> </table>		YES	NO	DK	BURN FOOD.....1	2		8	NEGLECT CHILD CARE....1	2		8	ARGUE WITH HUSBAND....1	2		8	TALK OTHER MEN.....1	2		8	SPEND NEEDLESSLY.....1	2		8	REPUSE INTERCOURSE....1	2		8
	YES	NO	DK																											
BURN FOOD.....1	2		8																											
NEGLECT CHILD CARE....1	2		8																											
ARGUE WITH HUSBAND....1	2		8																											
TALK OTHER MEN.....1	2		8																											
SPEND NEEDLESSLY.....1	2		8																											
REPUSE INTERCOURSE....1	2		8																											

767	<p>Now I will read you a few sentences. I would like to learn what you think about the ideas in these sentences. Do you agree or disagree with the following:</p> <p>The important decisions in the family should be made by the male members of the family.</p> <p>Men are usually wiser than women.</p> <p>A woman should not argue with her husband even if she does not share the same views with him.</p> <p>It is always better for the male child to have education than the female child.</p>	<table style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th style="text-align: center;">AGREES</th> <th style="text-align: center;">DISAGREES</th> <th style="text-align: center;">HAS NO IDEA</th> </tr> </thead> <tbody> <tr> <td>The important decisions in the family should be made by the male members of the family.</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> <td style="text-align: center;">8</td> </tr> <tr> <td>Men are usually wiser than women.</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> <td style="text-align: center;">8</td> </tr> <tr> <td>A woman should not argue with her husband even if she does not share the same views with him.</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> <td style="text-align: center;">8</td> </tr> <tr> <td>It is always better for the male child to have education than the female child.</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> <td style="text-align: center;">8</td> </tr> </tbody> </table>		AGREES	DISAGREES	HAS NO IDEA	The important decisions in the family should be made by the male members of the family.	1	2	8	Men are usually wiser than women.	1	2	8	A woman should not argue with her husband even if she does not share the same views with him.	1	2	8	It is always better for the male child to have education than the female child.	1	2	8
	AGREES	DISAGREES	HAS NO IDEA																			
The important decisions in the family should be made by the male members of the family.	1	2	8																			
Men are usually wiser than women.	1	2	8																			
A woman should not argue with her husband even if she does not share the same views with him.	1	2	8																			
It is always better for the male child to have education than the female child.	1	2	8																			

SECTION 8. SEXUALLY TRANSMITTED DISEASES AND AIDS

801A	<p>Now I would like to talk you about sexually transmitted diseases.</p> <p>Have you ever heard of sexually transmitted diseases ?</p>	<p>YES.....1</p> <p>NO.....2 →801L</p>
------	--	--

801B	<p>Which diseases have you heard ?</p> <p>RECORD ALL MENTIONED.</p>	<p>SYPHILIS.....A</p> <p>BEL SOĞUKLUĞU.....B</p> <p>AIDS.....C</p> <p>WART/SORE IN GENITAL ORGANS.....D</p> <p>FUNGUS.....E</p> <p>OTHER _____ U (SPECIFY)</p> <p>OTHER _____ V (SPECIFY)</p> <p>DOESN'T KNOW THE NAMES.....X</p>
------	---	---

801K	<p>CHECK 801B :</p> <p>NOT MENTIONED AIDS <input type="checkbox"/></p>	<p>MENTIONED AIDS <input type="checkbox"/> →802A</p>
------	--	--

801L	<p>Have you ever heard of an illness called AIDS ?</p>	<p>YES.....1</p> <p>NO.....2 →812</p>
------	--	---------------------------------------

802A	<p>From which sources of information have you learned most about AIDS ?</p> <p>Any other sources ?</p> <p>RECORD ALL MENTIONED.</p>	<p>RADIO.....A</p> <p>TV.....B</p> <p>NEWSPAPERS/MAGAZINES.....C</p> <p>PAMPHLETS/POSTERS.....D</p> <p>HEALTH WORKERS.....E</p> <p>MOSQUES.....F</p> <p>SCHOOLS/TEACHERS.....G</p> <p>FRIENDS/RELATIVES.....I</p> <p>WORK PLACE.....J</p> <p>OTHER _____ U (SPECIFY)</p>
------	---	--

802B	<p>Through what ways a person is transmitted AIDS ?</p> <p>Any other ways ?</p> <p>RECORD ALL MENTIONED.</p>	<p>SEXUAL RELATION.....A</p> <p>SEXUAL RELATION WITH MORE THAN ONE PARTNER.....B</p> <p>SEXUAL RELATION WITH A PROSTITUTE.....C</p> <p>NOT USING CONDOM.....D</p> <p>HOMOSEXUAL RELATION.....E</p> <p>BLOOD TRANSFUSION.....F</p> <p>INJECTION.....G</p> <p>KISSING.....H</p> <p>MOSQUITO BITE.....I</p> <p>OTHER _____ U (SPECIFY)</p> <p>OTHER _____ V (SPECIFY)</p> <p>DON'T KNOW.....X</p>
------	--	--

803	Is there anything a person can do to avoid getting AIDS or the virus that causes AIDS ?	YES.....1 NO.....2 DON'T KNOW.....8	→807
804	What can a person do ? Any other ways ? RECORD ALL MENTIONED.	SAFE SEX.....A ABSTAIN FROM SEX.....B USE CONDOMS.....C HAVE ONLY ONE SEX PARTNER.....D AVOID SEX WITH PROSTITUTES.....E AVOID SEX WITH HOMOSEXUALS.....F AVOID BLOOD TRANSFUSIONS.....G AVOID INJECTIONS.....H AVOID KISSING.....I AVOID MOSQUITO BITES.....J SEEK PROTECTION FROM TRADITIONAL HEALER.....K OTHER _____ U (SPECIFY) OTHER _____ V (SPECIFY) DON'T KNOW.....X	
807	Is it possible for a healthy-looking person to have the AIDS virus ?	YES.....1 NO.....2 DON'T KNOW.....8	
808A	Do you think that persons with AIDS almost never die from the disease, sometimes die, or almost always die from the disease ?	ALMOST NEVER.....1 SOMETIMES.....2 ALMOST ALWAYS.....3 DON'T KNOW.....8	
808B	Is there a medical treatment for AIDS ?	YES.....1 NO.....2 DON'T KNOW.....8	
808C	Is AIDS transmitted from mother to child ?	YES.....1 NO.....2 DON'T KNOW.....8	

812	RECORD THE TIME.	HOUR..... MINUTES.....	<table border="1" style="width: 20px; height: 20px; margin-left: auto; margin-right: auto;"> <tr><td style="width: 10px; height: 10px;"></td><td style="width: 10px; height: 10px;"></td></tr> <tr><td style="width: 10px; height: 10px;"></td><td style="width: 10px; height: 10px;"></td></tr> </table>				
813	PRESENCE OF OTHERS DURING THE INTERVIEW : CIRCLE ALL APPROPRIATE ALTERNATIVES.	NO ONE.....A CHILDREN UNDER 10.....B HUSBAND.....C MOTHER-IN-LAW.....D HER MOTHER.....E OTHER MEN.....F OTHER WOMEN.....G					
814	WAS THE INTERVIEW INTERRUPTED ? IF YES, FOR HOW MANY MINUTES APPROXIMATELY ?	NO.....000 YES.....1	<table border="1" style="width: 20px; height: 20px; margin-left: auto; margin-right: auto;"> <tr><td style="width: 10px; height: 10px;"></td><td style="width: 10px; height: 10px;"></td></tr> </table>				
815	WHAT IS THE RELIABILITY OF THE RESPONSES, IN YOUR OPINION ?	POOR.....1 FAIR.....2 GOOD.....3 VERY GOOD.....4					
816	WHAT LANGUAGE WAS USED DURING THE INTERVIEW ?	TURKISH.....1 KURDISH.....2 ARABIC.....3 OTHER _____ 7 (SPECIFY)	 →921				
817	WAS AN INTERPRETER USED DURING THE INTERVIEW ?	YES.....1 NO.....2					

SECTION 9. HEIGHT AND WEIGHT

921

- WRITE THE NAME OF WOMEN IN 923. RECORD HER HEIGHT AND WEIGHT IN 926 AND 928.
- IN COLUMNS 2 AND 3, RECORD THE CHILDREN BORN SINCE JANUARY 1993 AND STILL ALIVE. IN 922, RECORD THE LINE NUMBERS OF THE CHILDREN IN THE BIRTH HISTORY, IN 923, RECORD THEIR NAMES AND IN 924, RECORD THEIR DATES OF BIRTH. IF THERE ARE MORE THAN 2 CHILDREN THEN USE AN ADDITIONAL QUESTIONNAIRE.
- ASK WHETHER THE CHILDREN HAVE IDENTITY CERTIFICATES AND SAY YOU WOULD LIKE TO SEE THEM. (924B) RECORD THE DATES OF BIRTH IN THE IDENTITY CERTIFICATES (924C), COMPARE IT WITH THE INFORMATION IN 215 AND DETERMINE THE ONE THAT IS TRUE. (924D AND 924E). WHATEVER THE RESULT OF THE COMPARISON DO NOT MAKE ANY CORRECTIONS ON THE QUESTIONNAIRE PAPER ANYWAY.
- MEASURE THE HEIGHT AND WEIGHT OF THE LIVING CHILDREN BORN SINCE JANUARY 1993 AND RECORD THE FINDINGS IN THE RELATED QUESTIONS. WHILE DOING THE MEASUREMENTS, CHECK FOR BCG SCAR (TUBERCULOSIS INJECTION SCAR) ON TOP OF LEFT SHOULDER AND RECORD IT TO 925.

	1	WOMAN	2	YOUNGEST LIVING CHILD	3	NEXT-TO-YOUNGEST LIVING CHILD
--	---	-------	---	-----------------------	---	-------------------------------

922	LINE NO FROM Q.212		LINE NO.....		LINE NO.....	
-----	--------------------	--	--------------	--	--------------	--

923	NAME FROM Q.212 FOR CHILDREN		(NAME)		(NAME)		(NAME)
-----	---------------------------------	--	--------	--	--------	--	--------

924A	DATE OF BIRTH CHECK FROM Q.215 AND ASK FOR DAY OF BIRTH.		DAY.....		DAY.....		MONTH.....		MONTH.....		YEAR..		YEAR..	
------	---	--	----------	--	----------	--	------------	--	------------	--	--------	--	--------	--

924B	ASK IF THE CHILD HAVE IDENTITY CERTIFICATE. IF YES, THEN SAY YOU'D LIKE TO SEE IT.		YES, SEEN.....1		YES, SEEN.....1		YES, NOT SEEN...2		YES, NOT SEEN...2		NO.....3		NO.....3		DOSSN'T KNOW...8		DOSSN'T KNOW...8
------	---	--	-----------------	--	-----------------	--	-------------------	--	-------------------	--	----------	--	----------	--	------------------	--	------------------

924C	RECORD THE DATE OF BIRTH IN THE IDENTITY CERTIFICATE.		DAY.....		DAY.....		MONTH.....		MONTH.....		YEAR..		YEAR..	
------	---	--	----------	--	----------	--	------------	--	------------	--	--------	--	--------	--

924D	CHECK Q.215 AND COMPARE IT WITH THE DATE OF BIRTH IN Q.924C.		SAME.....1		SAME.....1		DIFFERENT.....2		DIFFERENT.....2
------	--	--	------------	--	------------	--	-----------------	--	-----------------

924E	DETERMINE THE TRUE INFORMATION FOR DATE OF BIRTH. DO NOT MAKE ANY CORRECTION.		ANSWER IN Q.215...1		ANSWER IN Q.215...1		ID. CERTIFICATE...2		ID. CERTIFICATE...2
------	---	--	---------------------	--	---------------------	--	---------------------	--	---------------------

		1 WOMAN	2 YOUNGEST LIVING CHILD	3 NEXT-TO-YOUNGEST LIVING CHILD
925	BCG (TUBERCULOSIS INJECTION) SCAR ON TOP OF LEFT SHOULDER		SCAR SEEN.....1 NO SCAR.....2	SCAR SEEN.....1 NO SCAR.....2
926	HEIGHT (in centimetres)			
927	WAS HEIGHT/LENGTH OF THE CHILD MEASURED LYING DOWN OR STANDING UP ?		LYING.....1 STANDING.....2	LYING.....1 STANDING.....2
928	WEIGHT (in kilograms)			
929	DATE WEIGHTED AND MEASURED	DAY..... MONTH..... YEAR.. 1 9	DAY..... MONTH..... YEAR.. 1 9	DAY..... MONTH..... YEAR.. 1 9
930	RESULT	MEASURED.....1 NOT PRESENT.....3 REFUSED.....4 OTHER _____ 7 (SPECIFY)	CHILD MEASURED....1 CHILD SICK.....2 CHILD NOT PRESENT.....3 CHILD REFUSED....4 MOTHER REFUSED...5 OTHER _____ 7 (SPECIFY)	CHILD MEASURED....1 CHILD SICK.....2 CHILD NOT PRESENT.....3 CHILD REFUSED....4 MOTHER REFUSED...5 OTHER _____ 7 (SPECIFY)
931	NAME OF MEASURER			

INTERVIEWER'S OBSERVATIONS
(To be filled after completing interview)

COMMENTS ABOUT WOMAN

COMMENTS ON SPECIFIC QUESTIONS

ANY OTHER COMMENTS

SUPERVISOR'S OBSERVATIONS

NAME OF THE SUPERVISOR:
DATE:

EDITOR'S OBSERVATIONS

NAME OF THE EDITOR:
DATE:

CALENDAR

INSTRUCTIONS:

ONLY ONE CODE SHOULD APPEAR IN ANY BOX.
FOR COLUMNS 1, 3, 4, AND 5a-5b, ALL MONTHS SHOULD BE FILLED IN.

COLUMN 1:

BIRTHS AND PREGNANCIES

- D BIRTHS
- H PREGNANCIES
- K INDUCED ABORTION
- F SPONTANEOUS ABORTION
- J STILL BIRTH

CONTRACEPTIVE USE

- 0 NO METHOD
- 1 PILL
- 2 IUD
- 3 INJECTABLE
- 4 NORPLANT
- 5 DIAPHRAGM/FOAM/JELLY
- 6 CONDOM
- 7 TUBAL LIGATION
- 9 MALE STERILIZATION
- 9 RHYTHM
- G WITHDRAWAL
- U OTHER _____

(SPECIFY)

- N MONTHS OUT OF WEDLOCK

COLUMN 2:

DISCONTINUATION USE

- 0 INFREQUENT SEX/HUSBAND AWAY
- 1 BECAME PREGNANT WHILE USING
- 2 WANTED TO BECOME PREGNANT
- 3 HUSBAND DISAPPROVED
- 4 WANTED MORE EFFECTIVE METHOD
- 5 HEALTH CONCERNS
- 6 SIDE EFFECTS
- 7 LACK OF ACCESS/TOO FAR
- 8 COST TOO MUCH
- 9 INCONVENIENT TO USE
- Y FATALISTIC
- M DIFFICULT TO GET PREGNANT/MENOPAUSE
- B MARITAL DISSOLUTION/SEPARATION/WIDOWHOOD
- U OTHER _____

(SPECIFY)

- X DON'T KNOW

PLACE OF INDUCED ABORTION

- C GOVERNMENT/SAMPLE HOSPITAL
- D MATERNITY HOUSE
- E MCHFP CENTRE
- F SSK HOSPITAL/DISPENSARY
- G OTHER PUBLIC SECTOR
- H PRIVATE HOSPITAL
- J PRIVATE POLYCLINIC
- K PRIVATE DOCTOR
- L OTHER PRIVATE SECTOR
- N UNIVERSITY HOSPITAL

COLUMN 3: MARRIAGE/UNION

- X MARRIED
- O NOT MARRIED

COLUMN 4: MOVES AND TYPES OF COMMUNITIES

- X CHANGE OF COMMUNITY
- 1 PROVINCE CENTRE
- 2 DISTRICT CENTRE
- 3 SUBDISTRICT/VILLAGE
- 4 ABROAD

COLUMN 5a AND 5b: MOVES AND PROVINCES

- X CHANGE OF COMMUNITY
- 01-80 PROVINCE CODES
- 90 ABROAD

		1	2	3	4	5a	5b		
12 DEC	01							01	DEC
11 NOV	02							02	NOV
10 OCT	03							03	OCT
09 SEP	04							04	SEP
1 09 AUG	05							05	AUG 1
9 07 JUL	06							06	JUL 9
9 06 JUN	07							07	JUN 9
8 05 MAY	08							08	MAY 8
04 APR	09							09	APR
03 MAR	10							10	MAR
02 FEB	11							11	FEB
01 JAN	12							12	JAN
12 DEC	13							13	DEC
11 NOV	14							14	NOV
10 OCT	15							15	OCT
09 SEP	16							16	SEP
1 08 AUG	17							17	AUG 1
9 07 JUL	18							18	JUL 9
9 06 JUN	19							19	JUN 9
7 05 MAY	20							20	MAY 7
04 APR	21							21	APR
03 MAR	22							22	MAR
02 FEB	23							23	FEB
01 JAN	24							24	JAN
12 DEC	25							25	DEC
11 NOV	26							26	NOV
10 OCT	27							27	OCT
09 SEP	28							28	SEP
1 08 AUG	29							29	AUG 1
9 07 JUL	30							30	JUL 9
9 06 JUN	31							31	JUN 9
6 05 MAY	32							32	MAY 6
04 APR	33							33	APR
03 MAR	34							34	MAR
02 FEB	35							35	FEB
01 JAN	36							36	JAN
12 DEC	37							37	DEC
11 NOV	38							38	NOV
10 OCT	39							39	OCT
09 SEP	40							40	SEP
1 08 AUG	41							41	AUG 1
9 07 JUL	42							42	JUL 9
9 06 JUN	43							43	JUN 9
5 05 MAY	44							44	MAY 5
4 04 APR	45							45	APR
03 MAR	46							46	MAR
02 FEB	47							47	FEB
01 JAN	48							48	JAN
12 DEC	49							49	DEC
11 NOV	50							50	NOV
10 OCT	51							51	OCT
09 SEP	52							52	SEP
1 08 AUG	53							53	AUG 1
9 07 JUL	54							54	JUL 9
9 06 JUN	55							55	JUN 9
4 05 MAY	56							56	MAY 4
04 APR	57							57	APR
03 MAR	58							58	MAR
02 FEB	59							59	FEB
01 JAN	60							60	JAN
12 DEC	61							61	DEC
11 NOV	62							62	NOV
10 OCT	63							63	OCT
09 SEP	64							64	SEP
1 08 AUG	65							65	AUG 1
9 07 JUL	66							66	JUL 9
9 06 JUN	67							67	JUN 9
3 05 MAY	68							68	MAY 3
04 APR	69							69	APR
03 MAR	70							70	MAR
02 FEB	71							71	FEB
01 JAN	72							72	JAN

PROVINCE CODES

01 ADANA	21 DIYARBAKIR	41 KOCAELI	61 TRABZON
02 ADIYAMAN	22 EDİRNE	42 KONYA	62 TUNCELI
03 AFYON	23 ELAZIĞ	43 KÜTAHYA	63 ŞANLIURFA
04 AĞRI	24 ERZİNCAN	44 MALATYA	64 UŞAK
05 AMASYA	25 ERZURUM	45 MANİSA	65 VAN
06 ANKARA	26 ESKİŞEHİR	46 K.MARAŞ	66 YOZGAT
07 ANTALYA	27 GAZİANTEP	47 MARDİN	67 ZONGULDAK
08 ARTVİN	28 GİRESUN	48 MUĞLA	68 AKSARAY
09 AYDIN	29 GÜMÜŞHANE	49 MUŞ	69 BAYBURT
10 BALIKESİR	30 HAKKARİ	50 NEVŞEHİR	70 KARAMAN
11 BİLECİK	31 HATAY	51 NİĞDE	71 KIRIKKALE
12 BİNGÖL	32 İSPARTA	52 ORDU	72 BATMAN
13 BİTLİS	33 İÇEL	53 RİZE	73 ŞIRNAK
14 BOLU	34 İSTANBUL	54 SAKARYA	74 BARTIN
15 BURDUR	35 İZMİR	55 SAMSUN	75 ARDAHAN
16 BURSA	36 KARS	56 SİRT	76 İĞDIR
17 ÇANAKKALE	37 KASTAMONU	57 SİNOP	77 YALOVA
18 ÇANKIRI	38 KAYSERİ	58 SİVAS	78 KARABÜK
19 ÇORUM	39 KIRKLARELİ	59 TEKİRDAĞ	79 KİLİS
20 DENİZLİ	40 KIRŞEHİR	60 TOKAT	80 OSMANIYE
90 OTHER COUNTRY			

CONVERSION OF YEARS OF BIRTH FROM
RUMI CALENDAR TO MILADI CALENDAR
YEARS :

RUMİ YEAR + 584 = MILADI YEAR

HACETTEPE UNIVERSITY INSTITUTE OF POPULATION STUDIES
 1998 TURKISH DEMOGRAPHIC AND HEALTH SURVEY
 NEVER-MARRIED (SINGLE) WOMAN'S QUESTIONNAIRE

IDENTIFICATION			
CLUSTER NO.....	□	□	PROVINCE _____
HOUSEHOLD NO.....	□	□	DISTRICT _____
REGION.....	□	□	SUBDISTRICT _____
URBAN (1) / RURAL (2).....	□	□	VILLAGE _____
			QUARTER _____
			STREET _____ DOOR NO _____

NAME-SURNAME OF WOMAN _____	LINE NO OF WOMAN □ □
-----------------------------	----------------------------

INTERVIEWER VISITS				
	1	2	3	FINAL VISIT
DATE (DAY-MONTH)	_ _	_ _	_ _	□ □ □ □
INTERVIEWER'S NAME-SURNAME	_____	_____	_____	□ □ □ □
RESULT (*)	_	_	_	□
NEXT DAY-MONTH VISIT HOUR	_ _	_ _		TOTAL NO OF VISITS □

(*) RESULT CODES :

1 COMPLETED	5 PARTLY COMPLETED
2 NOT AT HOME	
3 POSTPONED	7 OTHER _____
4 REFUSED	(SPECIFY)

SUPERVISOR _____ □ □ □	FIELD EDITOR _____ □ □ □	KEYED BY _____ □ □ □
DAY-MONTH □ □ □ □	DAY-MONTH □ □ □ □	DAY-MONTH □ □ □ □

Age	Year of birth	
	Has not had birthday in 1998	Has already had birthday in 1998
	Does not know	
0	1997	--
1	1996	1997
2	1995	1996
3	1994	1995
4	1993	1994
5	1992	1993
6	1991	1992
7	1990	1991
8	1989	1990
9	1988	1989
10	1987	1988
11	1986	1987
12	1985	1986
13	1984	1985
14	1983	1984
15	1982	1983
16	1981	1982
17	1980	1981
18	1979	1980
19	1978	1979
20	1977	1978
21	1976	1977
22	1975	1976
23	1974	1975
24	1973	1974
25	1972	1973
26	1971	1972
27	1970	1971
28	1969	1970
29	1968	1969
30	1967	1968
31	1966	1967
32	1965	1966
33	1964	1965
34	1963	1964

Age	Year of birth	
	Has not had birthday in 1998	Has already had birthday in 1998
	Does not know	
35	1962	1963
36	1961	1962
37	1960	1961
38	1959	1960
39	1958	1959
40	1957	1958
41	1956	1957
42	1955	1956
43	1954	1955
44	1953	1954
45	1952	1953
46	1951	1952
47	1950	1951
48	1949	1950
49	1948	1949
50	1947	1948
51	1946	1947
52	1945	1946
53	1944	1945
54	1943	1944
55	1942	1943
56	1941	1942
57	1940	1941
58	1939	1940
59	1938	1939
60	1937	1938
61	1936	1937
62	1935	1936
63	1934	1935
64	1933	1934
65	1932	1933
66	1931	1932
67	1930	1931
68	1929	1930
69	1928	1929

SECTION 1. RESPONDENT'S BACKGROUND

101	RECORD THE TIME	HOUR - MINUTES..... <input type="text"/> <input type="text"/>
102A	First I would like to ask some questions about you and the place you lived. For most of the time until you were 12 years old, where did you live ? _____ (NAME OF THE PLACE) Was it a city centre, district centre, a subdistrict or village ? Or did you live abroad ?	PROVINCE CENTRE.....1 DISTRICT CENTRE.....2 SUBDISTRICT OR VILLAGE.....3 ABROAD.....4 → 103
102B	In which province is this place now ? RECORD THE NAME AND CODE OF THE PROVINCE.	NAME OF PROVINCE <input type="text"/> _____ PROVINCE CODE
103	How long have you been living continuously in (NAME OF PLACE OF INTERVIEW) ?	YEARS..... <input type="text"/> SINCE I WAS BORN.....95 VISITOR.....96 → 105
104A	Where did you live before you moved here ? _____ (NAME OF THE PLACE) Was that a city centre, district centre, a subdistrict or village ? Or did you live abroad ?	PROVINCE CENTRE.....1 DISTRICT CENTRE.....2 SUBDISTRICT OR VILLAGE.....3 ABROAD.....4 → 104C
104B	In which province is this place now ? RECORD THE NAME AND CODE OF THE PROVINCE.	NAME OF PROVINCE <input type="text"/> _____ PROVINCE CODE
104C	What was the main reason for you to move here ?	PERSONAL REASONS EDUCATION.....02 LOOKING FOR JOB.....03 CHANGE OF JOB/APPOINTMENT.....04 RETURN TO HOMELAND.....05 FAMILIAL REASONS TO JOIN PARENTS.....09 CHANGE OF JOB/APPOINTMENT..10 LOOKING FOR JOB.....11 OTHER _____ 96 (SPECIFY)

105	In what month and year were you born ?	MONTH..... <input type="text"/> <input type="text"/> DON'T KNOW MONTH.....98 YEAR..... <input type="text"/> 1 <input type="text"/> 9 <input type="text"/> <input type="text"/> DON'T KNOW YEAR.....9998
106	How old are you exactly ? What age have you completed ? CHECK ANSWERS TO 105 AND 106 USING AGE-BIRTH YEAR TABLE. IF INCONSISTENT PROBE AND CORRECT. AGE INFORMATION MUST BE RECORDED !	AGE IN COMPLETED YEARS..... <input type="text"/> <input type="text"/>
107	Have you ever attended school ?	YES.....1 NO.....2 →114
108	What is the highest level you have attended ?	PRIMARY.....1 SECONDARY.....2 HIGH SCHOOL.....3 UNIVERSITY.....4
109A	What is the highest grade you have completed at that level ?	GRADE..... <input type="text"/>
109B	Did you graduate from this school ?	YES.....1 NO.....2
110	CHECK 106 : AGE 24 OR BELOW <input type="checkbox"/> AGE 25 OR ABOVE <input type="checkbox"/> →113	
111	Are you currently attending school ?	YES.....1 →113 NO.....2

112	<p>What was the main reason you stopped attending school ?</p>	<p>TO CARE FOR YOUNGER CHILDREN...03 FAMILY NEEDED HELP ON FARM OR IN BUSINESS.....04 COULD NOT PAY SCHOOL FEES.....05 NEEDED TO EARN MONEY.....06 GRADUATED/HAD ENOUGH SCHOOLING.07 DID NOT PASS ENTRANCE EXAMS....08 DID NOT LIKE SCHOOL.....09 SCHOOL NOT ACCESSIBLE/TOO FAR..10 FATHER/PARENTS DID NOT SEND....11</p> <p>OTHER _____ 96 (SPECIFY) DON'T KNOW.....98</p>
-----	--	--

113	<p>CHECK 108:</p> <p>ATTENDED PRIMARY <input type="checkbox"/></p> <p>ATTENDED SECONDARY OR HIGHER <input type="checkbox"/></p>	<p>→115</p>
-----	---	-------------

114	<p>Can you read and understand a letter or newspaper easily, with difficulty, or not at all ?</p>	<p>EASILY.....1 WITH DIFFICULTY.....2 NOT AT ALL.....3 →116A</p>
-----	---	--

115	<p>How frequent do you read a newspaper or magazine ?</p>	<p>EVERY DAY/ALMOST EVERY DAY.....1 ONCE-TWICE A WEEK.....2 RARELY/SELDOM.....3 NEVER/ALMOST NEVER.....4</p>
-----	---	---

116A	<p>What is your mother tongue ?</p> <p>RECORD ONLY ONE RESPONSE.</p>	<p>TURKISH.....01 KURDISH AND DIALECTS (KURMANCI, GORANI, ZAZA ETC)...02 ARABIC.....03 GREEK.....04 ARMANIAN.....05 HEBREW(LADINO).....06 CIRCASSIAN.....07 GEORGIAN.....08 LAZ LANGUAGE.....09 PERSIAN.....10 BULGARIAN.....11 RUMANIAN.....12 SERBIAN.....13 ENGLISH.....14 GERMAN.....15</p> <p>OTHER _____ 96 (SPECIFY)</p>
------	--	---

116B	<p>In addition to your mother tongue, which language(s) can you speak ?</p> <p style="text-align: center;">RECORD ALL MENTIONED.</p>	<p>TURKISH.....A KURDISH AND DIALECTS (KURMANCI, GORANI, ZAZA ETC)....B ARABIC.....C GREEK.....D ARMANIAN.....E HEBREW.....F CIRCASSIAN.....G GEORGIAN.....H LAZ LANGUAGE.....I PERSIAN.....J BULGARIAN.....K RUMANIAN.....L SERBIAN.....M ENGLISH.....N GERMAN.....O</p> <p>OTHER _____ U (SPECIFY)</p> <p>KNOWS NO OTHER LANGUAGE.....Y</p>
------	--	---

116C	<p>What is (was) your mother's and father's mother tongue ?</p> <p style="text-align: center;">USE THE CODES IN 116A.</p>	<p>MOTHER..... <table border="1" style="display: inline-table; width: 40px; height: 20px; vertical-align: middle;"></table></p> <p>FATHER..... <table border="1" style="display: inline-table; width: 40px; height: 20px; vertical-align: middle;"></table></p>
------	---	---

117	<p>What is your religion ?</p> <p style="text-align: center;">IF THE ANSWER IS "MUSLIM" PROBE FOR RELIGIOUS SECT AND CIRCLE APPROPRIATE CODE.</p>	<p>MUSLIM SUNNI.....01 ALAWI.....02</p> <p>OTHER _____ 03 (SPECIFY)</p> <p>CHRISTIAN.....06 JEWISH.....07 NO RELIGION.....10</p> <p>OTHER _____ 96 (SPECIFY)</p>
-----	---	--

118	<p>CHECK Q.4 IN THE HOUSEHOLD QUESTIONNAIRE :</p> <p>THE WOMAN INTERVIEWED IS NOT A USUAL RESIDENT <input type="checkbox"/></p>	<p>THE WOMAN INTERVIEWED IS A USUAL RESIDENT <input type="checkbox"/></p> <p style="text-align: right;">→234</p>
-----	---	--

119A	<p>Now I would like to ask about the place in which you usually live.</p> <p>What is the name of the place in which you usually live ?</p> <p style="text-align: center;">_____</p> <p style="text-align: center;">(NAME OF PLACE)</p> <p>Is that a city centre, a district centre, a subdistrict or village, or are you living abroad ?</p>	<p>PROVINCE CENTRE.....1 DISTRICT CENTRE.....2 SUBDISTRICT OR VILLAGE.....3 ABROAD.....4 →120A</p>
------	--	---

119B	In which province is that located ?	NAME OF PROVINCE <input type="text"/> PROVINCE CODE <input type="text"/>
120A	Now I would like to ask about the household you usually live. How many persons do usually live in your house ?	NUMBER..... <input type="text"/>
120B	Does the house you usually live belong to a household member, is it rented from someone else, is it a lodging, or do you just live here without having to pay anything ?	OWNED BY A HOUSEHOLD MEMBER.....1 RENTED.....2 LODGING.....3 NO MONEY PAID.....4 OTHER _____ 7 (SPECIFY)
121A	What is the source of drinking water for your household ?	PIPED WATER PIPED WATER IN HOUSE/GARDEN..11 →122A PUBLIC PIPED WATER OUTSIDE HOUSE/GARDEN.....12 WELL WATER WELL IN RESIDENCE/GARDEN....21 →122A PUBLIC WELL.....22 SURFACE WATER SPRING WATER PIPED INTO HOUSE/GARDEN.....31 →122A SPRING/PUBLIC FOUNTAIN.....32 RIVER/STREAM/ POND/LAKE/DAM.....33 RAINWATER.....41 →122A TANKER TRUCK.....51 BOTTLED WATER/DEMIJOHN/PET WTR.61 →122A WATER STATION.....71 OTHER _____ 96 (SPECIFY)
121B	How long does it take you to go there, get water, and come back ?	MINUTES..... <input type="text"/> ON PREMISES.....996
122A	Is the toilet inside the house or outside ?	NO FACILITY/BUSH/FIELD.....0 →123 INSIDE.....1 OUSIDE.....2 INSIDE AND OUTSIDE.....3 OTHER _____ 7 (SPECIFY)

122B	<p>What type of toilet system do you have in your household ?</p> <p>ASK WHETHER THE TOILET IS CONNECTED TO DRAINAGE SYSTEM</p> <p>IF IT IS CONNECTED :</p> <p>Is the toilet used only by this household or shared with another household ?</p> <p>IF IT IS NOT CONNECTED:</p> <p>What is the facility for excrata disposal ?</p>	<p>CONNECTED TO DRAINAGE SYSTEM</p> <p>USED ONLY BY THIS HOUSEHOLD...11</p> <p>SHARED WITH ANOTHER HOUSEHOLD...12</p> <p>PIT</p> <p>OPEN PIT.....21</p> <p>CLOSED PIT.....22</p> <p>NO FACILITY.....31</p> <p>OTHER _____ 96</p> <p>(SPECIFY)</p>
123	<p>What is the source of heating in winter for your house ?</p>	<p>RADIATOR (CENTRAL HEATING).....1</p> <p>RADIATOR (PRIVATE).....2</p> <p>NATURAL GAS STOVE.....3</p> <p>STOVE (COAL/WOOD).....4</p> <p>OTHER _____ 7</p> <p>(SPECIFY)</p>
124	<p>How many rooms in your household are used for sleeping ?</p>	<p>ROOMS USED FOR SLEEPING.... <input type="text"/></p>
125	<p>What is the main material of the floor ?</p>	<p>NATURAL FLOOR</p> <p>EARTH.....11</p> <p>RUDIMENTARY</p> <p>WOOD PLANKS.....21</p> <p>FINISHED FLOOR</p> <p>PARQUET/POLISHED WOOD.....31</p> <p>KARO.....32</p> <p>CEMENT.....34</p> <p>CARPET/WALL TO WALL CARPET...35</p> <p>MARLEY.....36</p> <p>MOSAIC.....37</p> <p>OTHER _____ 96</p> <p>(SPECIFY)</p>
126	<p>I would like to get an estimate of the total income that enters your household each month. Is the total amount of money earned by the members of your household in a month:</p> <p>126A More than 100 million ?</p> <p>126B More than 300 million ?</p> <p>126C More than 500 million ?</p> <p>126D Less than 50 million ?</p>	<p>YES.....1</p> <p>NO.....2 → 126D</p> <p>YES.....1</p> <p>NO.....2 → 127</p> <p>YES.....1</p> <p>NO.....2 → 127</p> <p>YES.....1</p> <p>NO.....2</p>

127

Do you have the following in your household ?

	YES	NO
Refrigerator	REFRIGERATOR.....1	2
Gas or electric oven	GAS/ELECTRIC OVEN.....1	2
Dishwasher	DISHWASHER.....1	2
Washing machine	WASHING MACHINE.....1	2
Vacuum cleaner	VACUUM CLEANER.....1	2
Television	TELEVISION.....1	2
Video	VIDEO.....1	2
Camera	CAMERA.....1	2
Music set with CD player	MUSIC SET WITH CD PLAYER...1	2
Telephone	TELEPHONE.....1	2
Cellular Telephone	CELLULAR TELEPHONE.....1	2
Car (excluding tractors, taxis, etc.)	CAR.....1	2
Computer	COMPUTER.....1	2

SECTION 2A. FERTILITY

234	<p>Now I would like to talk to you about subjects concerning marriage life and fertility.</p> <p>When did your last menstrual period start ?</p> <p>_____</p> <p>(DATE, IF GIVEN)</p>	<p>DAYS AGO.....1 <table border="1" style="display: inline-table; vertical-align: middle;"><tr><td style="width: 20px; height: 20px;"></td><td style="width: 20px; height: 20px;"></td></tr></table></p> <p>WEEKS AGO.....2 <table border="1" style="display: inline-table; vertical-align: middle;"><tr><td style="width: 20px; height: 20px;"></td><td style="width: 20px; height: 20px;"></td></tr></table></p> <p>MONTHS AGO.....3 <table border="1" style="display: inline-table; vertical-align: middle;"><tr><td style="width: 20px; height: 20px;"></td><td style="width: 20px; height: 20px;"></td></tr></table></p> <p>YEARS AGO.....4 <table border="1" style="display: inline-table; vertical-align: middle;"><tr><td style="width: 20px; height: 20px;"></td><td style="width: 20px; height: 20px;"></td></tr></table></p> <p>IN MENOPAUSE.....994</p> <p>NEVER MENSTRUATED.....996</p>								

235	<p>Between the first day of a woman's period and the first day of her next period, are there certain times when she has a greater chance of becoming pregnant than other times ?</p>	<p>YES.....1</p> <p>NO.....2</p> <p>DON'T KNOW.....8 → 237</p>
-----	--	--

236	<p>During which times of the monthly cycle does a woman have the greatest chance of becoming pregnant ?</p>	<p>DURING HER PERIOD.....1</p> <p>RIGHT AFTER HER PERIOD HAS ENDED.....2</p> <p>IN THE MIDDLE OF THE CYCLE.....3</p> <p>JUST BEFORE HER PERIOD BEGINS...4</p> <p>OTHER _____ 7</p> <p>(SPECIFY)</p> <p>DON'T KNOW.....8</p>
-----	---	---

237	<p>CHECK 106:</p> <p>YOUNGER THAN AGE 25 <input type="checkbox"/> AGED 25 OR OLDER THAN 25 <input type="checkbox"/> → 280</p>	
-----	--	--

238	<p>Suppose you get married in the future. If all of your children are girls, would you give more births than you normally desired for the chance of having a boy ?</p>	<p>YES.....1</p> <p>NO.....2</p> <p>DON'T KNOW.....8</p>
-----	--	--

239	<p>Let's suppose just the opposite. If all of your children are boys, would you give more births than you normally desired for the chance of having a girl ?</p>	<p>YES.....1</p> <p>NO.....2</p> <p>DON'T KNOW.....8</p>
-----	--	--

SECTION 2b. MARRIAGE

280	<p>Now I would like to ask you some questions about marriage. Do you currently have any arrangement or any plan for marriage ?</p>	<p>NO.....0 YES ENGAGED.....1 PROMISED.....2 DATING, PLANS MARRIGE.....3 OTHER _____7</p>
281	<p>I would like to learn your thoughts about marriage. Do you think civil ceremony is necessary for starting a husband-wife relationship ?</p>	<p>NECESSARY.....1 NOT NECESSARY.....2</p>
282	<p>What about religious ceremony ? Is it necessary ?</p>	<p>NECESSARY.....1 NOT NECESSARY.....2</p>
283	<p>In Turkey some couples cohabit as married by having only religious ceremony, without having civil ceremony. Do you think this is legal ?</p>	<p>LEGAL.....1 NOT LEGAL.....2</p>
284	<p>Do you think having only religious ceremony may have disadvantages for the woman, for the man or for the children ? IF YES: What kind of disadvantage(s) it may have ?</p>	<p>WOMAN DOESN'T HAVE GUARANTEE....A NOT HAVING OFFICIAL CHARACTER...B PROBLEM IN REGISTERING CHILDREN.....C POSSIBILITY FOR POLYGAMY (SECOND WIFE).....D OTHER _____ U (SPECIFY) OTHER _____ V (SPECIFY) NO DISADVANTAGE.....Y</p>
285	<p>Who should decide for marriage ? Should the decision for marriage made by the couples or by the families ?</p>	<p>COUPLES DECIDE.....1 FAMILIES DECIDE.....2 OTHER _____7 DON'T KNOW.....8</p>
286	<p>In a situation where the couples decide to marry, is it necessary to seek also the families' consent ?</p>	<p>YES.....1 NO.....2 OTHER _____7 DON'T KNOW.....8</p>

287	In a situation where the family arranged the marriage, should the consent of the man or the woman in question be sought ?	YES.....1 NO.....2 OTHER _____ 7 (SPECIFY) DON'T KNOW.....8
288 CHECK 106 : YOUNGER THAN AGE 25 <input type="checkbox"/> AGED 25 OR OLDER THAN 25 <input type="checkbox"/> → 290		
289	If you get married, will your family elders demand bridesmoney for you from the other side ?	YES.....1 NO.....2 OTHER _____ 7 (SPECIFY) DON'T KNOW.....8
290	Is it good or bad for the husband and wife being close relatives, being cousins for example ?	GOOD.....1 BAD.....2 → 292 HAS BOTH GOOD AND BAD SIDES.....3 DON'T KNOW.....8 → 301
291	What may be the possible advantages of marrying a relative ?	PROTECTING FAMILY ASSET.....A ADVANTAGE OF KNOWING HIM/HER BEFORE.....B NOT TO ALLOW NON RELATIVES INTO FAMILY.....C OTHER _____ U (SPECIFY) OTHER _____ V (SPECIFY) NO ADVANTAGES.....Y
292	What may be the possible disadvantages of marrying a relative ?	DISABLED BORN CHILDREN.....A ONGOING FAMILY PRESSURE.....B NO CHANCE OF CHOICE.....C OTHER _____ U (SPECIFY) OTHER _____ V (SPECIFY) NO DISADVANTAGES.....Y

SECTION 3. CONTRACEPTION

Now I would like to talk about family planning. There are various methods that a married couple can use to avoid pregnancy.

- CIRCLE CODE '1' IN IN Q. 301 FOR EACH METHOD MENTIONED SPONTANEOUSLY.
- THEN PROCEED DOWN COLUMN 302, READING THE NAME AND DESCRIPTION OF EACH METHOD NOT MENTIONED SPONTANEOUSLY AND ASK WHETHER SHE HAS HEARD THE METHOD. CIRCLE CODE '2' IF METHOD IS RECOGNIZED AND CODE '3' IF NOT RECOGNIZED.
- AFTER ASKING ABOUT ALL METHODS PROCEED TO 331B.

301 Which ways or methods have you heard ?	SPONTANEOUS	302 Have you heard this method ?	
	YES	PROBED YES	NO
01 PILL Woman can avoid a pregnancy by taking a pill every day.	1	2	3
02 IUD Women can have the so called spiral or IUD placed in them by a doctor or a nurse.	1	2	3
03 INJECTABLES Woman can have an injection by a doctor or a nurse which stops them from becoming pregnant for a certain period of time.	1	2	3
04 NORPLANT Woman can have small rods placed in their arm and this can prevent pregnancy for several years.	1	2	3
05 DIAPHRAGM, FOAM, JELLY Woman can place a sponge, suppository, diaphragm, jelly or cream inside themselves before intercourse.	1	2	3
06 CONDOM Men can put a rubber sheath on their penis during sexual intercourse.	1	2	3
07 TUBAL LIGATION Women can have an operation of tubal ligation to avoid having any more children.	1	2	3
08 MALE STERILIZATION Men can have an operation called vasectomy so that their wives would not get pregnant.	1	2	3
09 RHYTHM Some couples can avoid having sexual intercourse on certain days of the month when the woman is more likely to become pregnant.	1	2	3
10 WITHDRAWAL Some men pull out during sexual intercourse before climax.	1	2	3
11 Have you heard of any other method that women or men can use to avoid pregnancy ?	1 _____ (SPECIFY) _____ (SPECIFY) _____ (SPECIFY)		3 ↓

331B	CHECK 301 AND 302:	
	RECOGNIZED ONE METHOD AT LEAST <input type="checkbox"/>	RECOGNIZED NONE OF THE METHODS <input type="checkbox"/> → 337

332	Do you know of a place where couples can obtain a method of family planning ?	YES.....1 NO.....2 → 337
-----	---	-----------------------------

333	Where is that ?	PUBLIC SECTOR GOVERNMENT/SAMPLE HOSPITAL....A MATERNITY HOUSE.....B MOTHER-CHILD HEALTH AND FAMILY PLANNING (MCHPP) CENTRE....C HEALTH CENTRE.....D HEALTH HOUSE.....E SSK HOSPITAL/DISPENSARY.....F OTHER PUBLIC SECTOR _____ G (SPECIFY) PRIVATE SECTOR PRIVATE HOSPITAL.....H PRIVATE POLYCLINIC.....I PRIVATE DOCTOR.....J PRIVATE MIDWIFE/NURSE.....K PHARMACY.....L OTHER PRIVATE SECTOR _____ M (SPECIFY) UNIVERSITY HOSPITAL.....N COMMUNITY VOLUNTEERS/ASSOCIATION/FOUNDATION.....O MARKET/SHOP.....P FRIEND/RELATIVE/NEIGHBOUR.....R TRAD. MIDWIFE/MIDWIFE GRAN.....S OTHER _____ U (SPECIFY)
-----	-----------------	--

337	Do you think a woman's chance of becoming pregnant is influenced by breastfeeding ?	YES.....1 NO.....2 DON'T KNOW.....8 → 602
-----	---	---

338	Do you think a woman's chance of becoming pregnant is increased or decreased by breastfeeding ?	INCREASED.....1 DECREASED.....2 DEPENDS.....3 DON'T KNOW.....8
-----	---	---

SECTION 6. FERTILITY PREFERENCES

602	Now I have some questions about the future. Would you like to have a child or would you prefer not to have any children ?	HAVE A CHILD.....1 NONE.....2 SAYS SHE WON'T MARRY / SHE CAN'T GET PREGNANT.....3 UNDECIDED/DON'T KNOW.....8
-----	---	---

→606A

602A	How many children would you like to have in the future ?	NUMBER..... <input type="text"/> <input type="text"/> OTHER ANSWERS _____ 96 (SPECIFY)
------	--	--

602B	How many of these children would you like to be boys, how many would you like to be girls and for how many would it not matter ?	<table style="width: 100%;"> <tr> <td style="text-align: right;">BOYS</td> </tr> <tr> <td>NUMBER..... <input type="text"/> <input type="text"/></td> </tr> <tr> <td>OTHER _____ 96 (SPECIFY)</td> </tr> <tr> <td style="text-align: right;">GIRLS</td> </tr> <tr> <td>NUMBER..... <input type="text"/> <input type="text"/></td> </tr> <tr> <td>OTHER _____ 96 (SPECIFY)</td> </tr> <tr> <td style="text-align: right;">EITHER</td> </tr> <tr> <td>NUMBER..... <input type="text"/> <input type="text"/></td> </tr> <tr> <td>OTHER _____ 96 (SPECIFY)</td> </tr> </table>	BOYS	NUMBER..... <input type="text"/> <input type="text"/>	OTHER _____ 96 (SPECIFY)	GIRLS	NUMBER..... <input type="text"/> <input type="text"/>	OTHER _____ 96 (SPECIFY)	EITHER	NUMBER..... <input type="text"/> <input type="text"/>	OTHER _____ 96 (SPECIFY)
BOYS											
NUMBER..... <input type="text"/> <input type="text"/>											
OTHER _____ 96 (SPECIFY)											
GIRLS											
NUMBER..... <input type="text"/> <input type="text"/>											
OTHER _____ 96 (SPECIFY)											
EITHER											
NUMBER..... <input type="text"/> <input type="text"/>											
OTHER _____ 96 (SPECIFY)											

606A	CHECK 301 AND 302:	
	HEARD AT LEAST ONE METHOD <input type="checkbox"/>	HEARD NONE OF THE METHODS <input type="checkbox"/>

→614

608	Do you think you will use a method to delay or avoid pregnancy after you got married ?	YES.....1 NO.....2 OTHER _____ 7 (SPECIFY) DON'T KNOW.....8
-----	--	---

→610

609 Which method would you prefer to use ?

PILL.....01
IUD.....02
INJECTABLES.....03
NORPLANT.....04
DIAPHRAGM/FOAM/JELLY.....05
CONDOM.....06
TUBAL LIGATION.....07
MALE STERILIZATION.....08
RHYTHM.....09
WITHDRAWAL.....10

OTHER _____ 96
(SPECIFY)
NOT SURE/DON'T KNOW.....98

→614

610 What is the main reason that you think you will never use a method ?

PERTILITY-RELATED REASONS

MENAPOUSAL/HYSTERECTOMY.....23
SUBFECUND/INFECOND.....24
WANTS MORE CHILDREN.....26

OPPOSITION TO USE
WOMAN OPPOSED.....31
OTHERS OPPOSED.....33
RELIGIOUS REASONS.....34

LACK OF KNOWLEDGE
KNOWS NO METHOD.....41
KNOWS NO SOURCE.....42

METHOD-RELATED REASONS
HEALTH CONCERNS.....51
SIDE EFFECTS.....52
LACK OF ACCESS/TOO FAR.....53
COST TOO MUCH.....54
INCONVENIENT TO USE.....55
INTERFERES WITH BODY'S
NORMAL PROCESSES.....56

OTHER _____ 96
(SPECIFY)
DON'T KNOW.....98

614 Do you approve of couples using a method to avoid getting pregnant ?

APPROVE.....1
DISAPPROVE.....2
NO OPINION.....3

615	Is it acceptable or not acceptable to you for information on family planning to be provided:			NOT	DO
			ACCEPT-	ACCEPT-	NOT
			ABLE	ABLE	KNOW
		On the radio ?	RADIO.....1	2	8
		On the television ?	TELEVISION.....1	2	8
		In newspapers/magazines ?	NEWSP./MAGZ....1	2	8
	In secondary schools ?	SECONDARY.....1	2	8	
	In high schools ?	HIGH.....1	2	8	

616	In the last few months have you heard about family planning?			YES	NO
		On the radio ?	RADIO.....1	2	
		On the television ?	TELEVISION.....1	2	
		In newspaper or magazine ?	NEWSPAPER OR MAGAZINE.....1	2	
		From a poster ?	POSTER.....1	2	
		From brochures or leaflets ?	BROCHURES/LEAFLETS.....1	2	
		From billboards or cloth posters ?	BILLBOARDS/CLOTH POSTERS....1	2	
		Any other place ?	ANOTHER PLACE.....1	2	

618	In the last few months have you discussed the practice of family planning with friends, neighbours, or relatives ?	YES.....11	
		NO.....22	→628A

619	With whom ? Anyone else ? RECORD ALL MENTIONED.	MOTHER.....B
		FATHER.....C
		SISTER(S).....D
		BROTHER(S).....E
		FRIEND/RELATIVE.....I
		NEIGHBOUR.....J
		OTHER _____ U
		(SPECIFY)

628A	Do you think that the use of family planning is against religion ?	YES.....1	→708E
		SOME METHODS ARE.....2	
		NO.....3	
		NOT RELIGIOUS/NO RELIGION.....4	→708E
		DON'T KNOW.....8	

628B	<p>Which method(s) do you think (are) is against religion ?</p> <p>Any other ?</p> <p>RECORD ALL MENTIONED.</p>	<p>PILL.....A</p> <p>IUD.....B</p> <p>INJECTABLES.....C</p> <p>NORPLANT.....D</p> <p>DIAPHRAGM/FOAM/JELLY.....E</p> <p>CONDOM.....F</p> <p>TUBAL LIGATION.....G</p> <p>MALE STERILIZATION.....H</p> <p>RHYTHM.....I</p> <p>WITHDRAWAL.....J</p> <p>SEXUAL ABSTINANCE.....K</p> <p>OTHER METHOD _____ U</p> <p>(SPECIFY)</p> <p>INDUCED ABORTION.....M</p>
------	---	---

628C	<p>Does(did)your mother have any objections to any family planning method or to family planning in general on religious grounds ?</p>	<p>YES.....1</p> <p>THINKS SOME METHODS ARE AGAINST RELIGION.....2</p> <p>DON'T BELIEVE/NO RELIGION.....4</p> <p>DON'T KNOW.....8</p>
------	---	---

SECTION 7A. MIGRATION

708E	Have you lived in only one settlement or more than one settlements since January 1993 ?	ONE SETTLEMENT.....1 MORE THAN ONE SETTLEMENT.....2 → 708G
------	---	---

708F	<p>C</p> <ul style="list-style-type: none"> ▪ ENTER IN COL.4 OF CALENDAR THE APPROPRIATE CODE FOR SETTLEMENT OF CURRENT RESIDENCE ("1" PROVINCE CENTER, "2" DISTRICT CENTER, "3" SUB-DISTRICT/VILLAGE,"4" ABROAD.) ▪ ENTER (IN COLUMN 5 OF THE CALENDAR) THE PROVINCE CODE FOR THE SETTLEMENT. ▪ BEGIN IN THE MONTH OF INTERVIEW AND CONTINUE WITH ALL PRECEDING MONTHS BACK TO JANUARY 1993. ▪ THEN SKIP TO 709A. 	
------	---	--

708G	<p>In what month and year did you move to (NAME OF THE SETTLEMENT OF CURRENT RESIDENCE)?</p> <p>C ENTER (IN COLUMNS 4 AND 5 OF THE CALENDAR) "X" IN THE MONTH AND YEAR OF THE MOVE.</p> <p>FOR COLUMN 4: FOR THE FOLLOWING MONTHS ENTER THE APPROPRIATE CODE FOR SETTLEMENT. ('1' PROVINCE CENTER, '2' DISTRICT CENTER, '3' SUB-DISTRICT/VILLAGE, '4' ABROAD).</p> <p>FOR COLUMN 5: FOR THE FOLLOWING MONTHS ENTER THE PROVINCE CODE THAT THE SETTLEMENT IS CURRENTLY LOCATED.</p> <p>CONTINUE PROBING FOR PREVIOUS SETTLEMENTS AND RECORD MOVES, TYPES OF SETTLEMENTS AND PROVINCE CODES ACCORDINGLY.</p> <p>RECORD ALSO MOVES BETWEEN THE SAME TYPE OF SETTLEMENTS OR BETWEEN THE TYPES OF SETTLEMENTS IN THE SAME PROVINCE.</p> <p>ILLUSTRATIVE QUESTIONS</p> <ul style="list-style-type: none"> > Where did you live before..... ? > In what month and year did you arrive there ? > Is that place a province center, a district center, a subdistrict/village or abroad? > In which province islocated ? 	
------	---	--

SECTION 7B. WOMAN'S WORK AND STATUS

709	<p>Now I would like to ask you questions about working.</p> <p>Aside from your own housework, are you currently working?</p>	<p>YES.....1 → 712</p> <p>NO.....2</p>			
710	<p>You say that you are not working. As you know, some women sell small things, sell goods at the market place, work on the family farm or business, look after children, work as cleaning ladies etc. Are you doing any of these at the moment, or any other work of similar nature ?</p>	<p>YES.....1 → 712</p> <p>NO.....2</p>			
711	<p>Have you worked in any job in the last 12 months ?</p>	<p>YES.....1 → 740</p> <p>NO.....2</p>			
712	<p>What type of work are/were you doing ? What kind of job are/were you in ?</p>	<p>_____ <table border="1" data-bbox="1169 976 1266 1039"><tr><td> </td><td> </td><td> </td></tr></table></p> <p>_____</p>			
712A	<p>Do/did you pay social security when doing this job ?</p> <p>IF YES: According to which schedule ?</p>	<p>NO.....0</p> <p>SSK.....1</p> <p>EMEKLİ SANDIĞI.....2</p> <p>BAĞ-KUR.....3</p> <p>OTHER _____ 7</p> <p>(SPECIFY)</p> <p>DON'T KNOW.....8</p>			

712B	CHECK 712, CIRCLE THE APPROPRIATE WORK CODE.	AGRICULTURE.....01 ANIMAL HUSBANDRY.....02 AGRICULTURE AND A.HUSBANDRY...03 FORESTRY.....04 FISHERY.....05 MINING.....06 MANUFACTURE INDUSTRY (FOOD)...07 MANUFACTURE INDUSTRY (TEXTILE).08 MANUFACTURE INDUSTRY (METAL)...09 MANUFACTURE INDUSTRY (OTHER)...10 ELECTRICITY/GAS/WATER.....11 CONSTRUCTION.....12 WHOLESALE/RETAIL TRADE.....13 HOTEL AND RESTAURANT.....14 COMMUNICATION/TRANSPORTATION...15 HEALTH.....16 EDUCATION/CULTURE.....17 PUBLIC ADMINISTRATION.....18 TOURISM.....19 BANKING/INSURANCE.....20 OTHER SOCIAL SERVICES.....21 PERSONAL SERVICES.....22 OTHER _____ 96 (SPECIFY)	
------	--	--	--

712C	CHECK 712 , CIRCLE THE APPROPRIATE CODE FOR THE POSITION AT WORK	EMPLOYER EMPLOYING TEN OR MORE THAN TEN PERSONS.....01 SMALL EMPLOYER EMPLOYING LESS THAN TEN PERSONS.....02 SALARIED (GOVERNMENT OFFICIAL).03 WAGED (PUBLIC SECTOR).....04 WAGED (PRIVATE SECTOR).....05 FOR HIS OWN.....06 UNPAID FAMILY LABOURER.....07 OTHER _____ 96 (SPECIFY)	
------	--	---	--

713	CHECK 712B :		
	AGRICULTURE (01) OR ARRICULTURE AND ANIMAL HUSBANDRY (03) <input type="checkbox"/>	OTHER WORK CODES <input type="checkbox"/>	→716

714	Whom does the land you work on belong to ?	HER LAND.....01 FAMILY LAND.....02 RENTED LAND.....03 SOMEONE ELSE'S LAND.....04 STATE LAND.....05 OTHER _____ 96 (SPECIFY)	
-----	--	---	--

716	Do you usually work throughout the year, or do you work seasonally, or only once in a while ?	THROUGHOUT THE YEAR.....1 → 718 SEASONALLY/PART OF THE YEAR.....2 ONCE IN A WHILE.....3 → 719
717	During the last 12 months, how many months did you work ?	NUMBER OF MONTHS..... <input type="text"/>
718	During the last 12 months, how many days a week did you usually work (in the months that you worked) ?	NUMBER OF DAYS..... <input type="text"/> → 720
719	During the last 12 months, approximately, how many days did you work ?	NUMBER OF DAYS..... <input type="text"/>
720	Do/did you earn cash for your work ? PROBE : Do/did you make money for your work ?	YES.....1 NO.....2 → 723
721A	How much do you usually earn for this work ? PROBE : Is this by the day, by the week, or by the month ?	PER HOUR.....1 <input type="text"/> PER DAY.....2 <input type="text"/> PER WEEK.....3 <input type="text"/> PER MONTH.....4 <input type="text"/> PER YEAR.....5 <input type="text"/> OTHER _____ 999996 → 722 (SPECIFY)
721B	CIRCLE THE APPROPRIATE CODE FOR THE GIVEN QUANTITY IN 721A .	MILLION.....1 THOUSAND.....2
721C	(In the times that you work) Generally what amount of the family expenses is met by your earnings? The whole, more than the half, the half, less than the half, or does not your earning have any contribution to the family expenses ?	THE WHOLE.....1 MORE THAN THE HALF.....2 THE HALF.....3 LESS THAN THE HALF.....4 NO CONTRIBUTION.....5 OTHER _____ 7 (SPECIFY) DON'T KNOW.....8

722	Who mainly decides how the money you earn will be used : you or someone else ?	HERSELF DECIDES.....1 SOMEONE ELSE DECIDES.....4 JOINTLY WITH SOMEONE ELSE.....5
-----	--	--

723	Do you usually work at home or away from home ?	HOME.....1 AWAY.....2
-----	---	--------------------------

740	CURRENTLY NOT WORKING (709=2 AND 710=2) <input type="checkbox"/>	CURRENTLY WORKING (709=1 OR 710=1) OR WORKED DURING THE LAST 12 MONTHS (711=1) <input type="checkbox"/>
-----	--	--

741	You told currently you are not working. What is the main reason that you are not working ?	HOLIDAY/ ON VACATION.....01 LOOKS AFTER THE CHILD(S).....02 HOUSEWORK.....03 SICK/HANDICAPPED.....04 CAN'T FIND/LOOKING FOR JOB.....05 ELDERS DON'T WANT.....06 NO NEED FOR WORKING.....07 DOES NOT WANT WORKING ANYMORE..08 NO TALENT/EDUCATION.....09 DISCHARGED.....10 OTHER _____ 96 (SPECIFY)
-----	---	---

741B	CHECK 711:	WORKED DURING THE LAST 12 MONTHS <input type="checkbox"/>	NOT WORKED DURING THE LAST 12 MONTHS <input type="checkbox"/>
------	------------	--	--

742	How long have/had you been working in your current/last job ?	YEAR..... <input type="text"/>
-----	---	--------------------------------

757	What is the main source of income providing your and your family's subsistence?	HER EARNINGS.....03 HER FAMILY'S INCOME.....05 ALIMONY.....06 HER PENSION.....07 PARENTS' PENSION.....08 OTHER _____ 96 (SPECIFY)
-----	---	---

758	<p>Are you covered by any health insurance ?</p> <p>IP YES : According to which schedule ?</p>	<p>NO.....0</p> <p>SSK.....1</p> <p>EMEKLİ SANDIĞI.....2</p> <p>BAÇ-KUR.....3</p> <p>PRIVATE INSURANCE.....4</p> <p>GREEN CARD.....5</p> <p>OTHER _____ 7</p> <p>(SPECIFY)</p> <p>DON'T KNOW.....8</p>																												
766	<p>Women can sometimes make things that may annoy or make angry her husband. Now I will talk about some situations. Can you please tell me if the husband can have the right for beating his wife in those situations?</p> <p>If she burns the food ?</p> <p>If she neglects the care of children ?</p> <p>If she argues with her husband ?</p> <p>If she talks with other men ?</p> <p>If she spends the money needlessly ?</p> <p>If she refuses to have sexual intercourse ?</p>	<table border="1"> <thead> <tr> <th></th> <th>YES</th> <th>NO</th> <th>DK</th> </tr> </thead> <tbody> <tr> <td>BURN FOOD.....1</td> <td>2</td> <td>8</td> <td></td> </tr> <tr> <td>NEGLECT CHILD CARE....1</td> <td>2</td> <td>8</td> <td></td> </tr> <tr> <td>ARGUE WITH HUSBAND....1</td> <td>2</td> <td>8</td> <td></td> </tr> <tr> <td>TALK OTHER MEN.....1</td> <td>2</td> <td>8</td> <td></td> </tr> <tr> <td>SPEND NEEDLESSLY.....1</td> <td>2</td> <td>8</td> <td></td> </tr> <tr> <td>SEXUAL INTERCOURSE....1</td> <td>2</td> <td>8</td> <td></td> </tr> </tbody> </table>		YES	NO	DK	BURN FOOD.....1	2	8		NEGLECT CHILD CARE....1	2	8		ARGUE WITH HUSBAND....1	2	8		TALK OTHER MEN.....1	2	8		SPEND NEEDLESSLY.....1	2	8		SEXUAL INTERCOURSE....1	2	8	
	YES	NO	DK																											
BURN FOOD.....1	2	8																												
NEGLECT CHILD CARE....1	2	8																												
ARGUE WITH HUSBAND....1	2	8																												
TALK OTHER MEN.....1	2	8																												
SPEND NEEDLESSLY.....1	2	8																												
SEXUAL INTERCOURSE....1	2	8																												
767	<p>Now I will read you a few sentences. I would like to learn what you think about the ideas in these sentences. Do you think they are right or wrong ?</p> <p>The important decisions in the family should be made by the male family members.</p> <p>Men are usually wiser than women.</p> <p>A woman should not argue with her husband even if does not share the same views with him.</p> <p>It is better always better for the male child to have education than the female child.</p>	<table border="1"> <thead> <tr> <th></th> <th>AGREES</th> <th>DISAGREES</th> <th>HAS NO IDEA</th> </tr> </thead> <tbody> <tr> <td>The important decisions in the family should be made by the male family members.</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Men are usually wiser than women.</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>A woman should not argue with her husband even if does not share the same views with him.</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>It is better always better for the male child to have education than the female child.</td> <td>1</td> <td>2</td> <td>8</td> </tr> </tbody> </table>		AGREES	DISAGREES	HAS NO IDEA	The important decisions in the family should be made by the male family members.	1	2	8	Men are usually wiser than women.	1	2	8	A woman should not argue with her husband even if does not share the same views with him.	1	2	8	It is better always better for the male child to have education than the female child.	1	2	8								
	AGREES	DISAGREES	HAS NO IDEA																											
The important decisions in the family should be made by the male family members.	1	2	8																											
Men are usually wiser than women.	1	2	8																											
A woman should not argue with her husband even if does not share the same views with him.	1	2	8																											
It is better always better for the male child to have education than the female child.	1	2	8																											

SECTION 8. SEXUALLY TRANSMITTED DISEASES AND AIDS

801A	<p>Now I would like you talk you about sexually transmitted diseases.</p> <p>Have you ever heard of sexually transmitted diseases ?</p>	<p>YES.....1</p> <p>NO.....2 → 801L</p>
------	---	---

801B	<p>Which diseases have you heard ?</p> <p>RECORD ALL MENTIONED.</p>	<p>SYPHILIS.....A</p> <p>BEL SOĞUKLUĞU.....B</p> <p>AIDS.....C</p> <p>WART/SORE IN GENITAL ORGANS.....D</p> <p>FUNGUS.....E</p> <p>OTHER _____ U</p> <p>(SPECIFY)</p> <p>OTHER _____ V</p> <p>(SPECIFY)</p> <p>DOESN'T KNOW THE NAMES.....X</p>
------	---	---

801K	<p>CHECK 801B :</p> <p>NOT MENTIONED AIDS <input type="checkbox"/> MENTIONED AIDS <input type="checkbox"/> → 802A</p>	
------	--	--

801L	<p>Have you ever heard of an illness called AIDS ?</p>	<p>YES.....1</p> <p>NO.....2 → 812</p>
------	--	--

802A	<p>From which sources of information have you learned most about AIDS ?</p> <p>Any other sources ?</p> <p>RECORD ALL MENTIONED.</p>	<p>RADIO.....A</p> <p>TV.....B</p> <p>NEWSPAPERS/MAGAZINES.....C</p> <p>PAMPHLETS/POSTERS.....D</p> <p>HEALTH WORKERS.....E</p> <p>MOSQUES.....F</p> <p>SCHOOLS/TEACHERS.....G</p> <p>FRIENDS/RELATIVES.....I</p> <p>WORK PLACE.....J</p> <p>OTHER _____ U</p> <p>(SPECIFY)</p>
------	---	---

802B	<p>Through what ways a person is transmitted AIDS ?</p> <p>Any other ways ?</p> <p>RECORD ALL MENTIONED.</p>	<p>SEXUAL RELATION.....A</p> <p>SEXUAL RELATION WITH MORE THAN ONE PARTNER.....B</p> <p>SEXUAL RELATION WITH A PROSTITUTE.....C</p> <p>NOT USING CONDOM.....D</p> <p>HOMOSEXUAL RELATION.....E</p> <p>BLOOD TRANSFUSION.....F</p> <p>INJECTION.....G</p> <p>KISSING.....H</p> <p>MOSQUITO BITE.....I</p> <p>OTHER _____ U (SPECIFY)</p> <p>OTHER _____ V (SPECIFY)</p> <p>DON'T KNOW.....X</p>
803	<p>Is there anything a person can do to avoid getting AIDS or the virus that causes AIDS ?</p>	<p>YES.....1</p> <p>NO.....2</p> <p>DON'T KNOW.....8 → 807</p>
804	<p>What can a person do ?</p> <p>Any other ways ?</p> <p>RECORD ALL MENTIONED.</p>	<p>SAFE SEX.....A</p> <p>ABSTAIN FROM SEX.....B</p> <p>USE CONDOMS.....C</p> <p>HAVE ONLY ONE SEX PARTNER.....D</p> <p>AVOID SEX WITH PROSTITUTES.....E</p> <p>AVOID SEX WITH HOMOSEXUALS.....F</p> <p>AVOID BLOOD TRANSFUSIONS.....G</p> <p>AVOID INJECTIONS.....H</p> <p>AVOID KISSING.....I</p> <p>AVOID MOSQUITO BITES.....J</p> <p>SEEK PROTECTION FROM TRADITIONAL HEALER.....K</p> <p>OTHER _____ U (SPECIFY)</p> <p>OTHER _____ V (SPECIFY)</p> <p>DON'T KNOW.....X</p>
807	<p>Is it possible for a healthy-looking person to have the AIDS virus ?</p>	<p>YES.....1</p> <p>NO.....2</p> <p>DON'T KNOW.....8</p>
808A	<p>Do you think that persons with AIDS almost never die from the disease, sometimes die, or almost always die from the disease ?</p>	<p>ALMOST NEVER.....1</p> <p>SOMETIMES.....2</p> <p>ALMOST ALWAYS.....3</p> <p>DON'T KNOW.....8</p>

808B	Is there a medical treatment for AIDS ?	YES.....1 NO.....2 DON'T KNOW.....8	
808C	Is AIDS transmitted from mother to child ?	YES.....1 NO.....2 DON'T KNOW.....8	
812	RECORD THE TIME.	HOUR..... <input type="text"/> <input type="text"/> MINUTES..... <input type="text"/> <input type="text"/>	
813	PRESENCE OF OTHERS DURING THE INTERVIEW : CIRCLE ALL APPROPRIATE ALTERNATIVES.	NO ONE.....A CHILDREN UNDER 10.....B HER MOTHER.....E OTHER MEN.....F OTHER WOMEN.....G	
814	WAS THE INTERVIEW INTERRUPTED ? IF YES, FOR HOW LONG, APPROXIMATELY ? (IN MINUTES)	NO.....000 YES.....1 <input type="text"/> <input type="text"/>	
815	WHAT IS THE RELIABILITY OF THE RESPONSES, IN YOUR OPINION ?	POOR.....1 FAIR.....2 GOOD.....3 VERY GOOD.....4	
816	WHAT LANGUAGE WAS USED DURING THE INTERVIEW ?	TURKISH.....1 → 921 KURDISH.....2 ARABIC.....3 OTHER _____ 7 (SPECIFY)	
817	WAS AN INTERPRETER USED DURING THE INTERVIEW ?	YES.....1 NO.....2	

SECTION 9. HEIGHT AND WEIGHT

921	<p>■ WRITE NAME OF WOMAN TO Q.923</p> <p>■ WRITE THE HEIGHT AND WEIGHT OF WOMAN IN 926 AND 928</p>																
	<table border="1"> <tr> <td data-bbox="950 478 998 598">1</td> <td data-bbox="998 478 1174 598">WOMAN</td> </tr> </table>	1	WOMAN														
1	WOMAN																
923	<table border="1"> <tr> <td data-bbox="609 598 950 703">NAME</td> <td data-bbox="950 598 1174 703">(NAME)</td> </tr> <tr> <td colspan="2" data-bbox="609 661 1174 703">_____</td> </tr> </table>	NAME	(NAME)	_____													
NAME	(NAME)																

926	<table border="1"> <tr> <td data-bbox="609 703 950 808">HEIGHT (in centimeters)</td> <td data-bbox="950 703 1174 808"> <table border="1"> <tr> <td> </td><td> </td><td> </td><td> </td><td> </td> </tr> </table> </td> </tr> </table>	HEIGHT (in centimeters)	<table border="1"> <tr> <td> </td><td> </td><td> </td><td> </td><td> </td> </tr> </table>														
HEIGHT (in centimeters)	<table border="1"> <tr> <td> </td><td> </td><td> </td><td> </td><td> </td> </tr> </table>																
928	<table border="1"> <tr> <td data-bbox="609 808 950 913">WEIGHT (in kilograms)</td> <td data-bbox="950 808 1174 913"> <table border="1"> <tr> <td> </td><td> </td><td> </td><td> </td><td> </td> </tr> </table> </td> </tr> </table>	WEIGHT (in kilograms)	<table border="1"> <tr> <td> </td><td> </td><td> </td><td> </td><td> </td> </tr> </table>														
WEIGHT (in kilograms)	<table border="1"> <tr> <td> </td><td> </td><td> </td><td> </td><td> </td> </tr> </table>																
929	<table border="1"> <tr> <td data-bbox="609 913 950 1144">DATE WEIGHTED AND MEASURED</td> <td data-bbox="950 913 1174 1144"> <table border="1"> <tr> <td data-bbox="950 913 1079 997">DAY.....</td> <td data-bbox="1079 913 1174 997"> <table border="1"> <tr><td> </td><td> </td></tr> </table> </td> </tr> <tr> <td data-bbox="950 997 1079 1081">MONTH.....</td> <td data-bbox="1079 997 1174 1081"> <table border="1"> <tr><td> </td><td> </td></tr> </table> </td> </tr> <tr> <td data-bbox="950 1081 1079 1144">YEAR..</td> <td data-bbox="1079 1081 1174 1144"> <table border="1"> <tr> <td>1</td><td>9</td><td> </td><td> </td> </tr> </table> </td> </tr> </table> </td> </tr> </table>	DATE WEIGHTED AND MEASURED	<table border="1"> <tr> <td data-bbox="950 913 1079 997">DAY.....</td> <td data-bbox="1079 913 1174 997"> <table border="1"> <tr><td> </td><td> </td></tr> </table> </td> </tr> <tr> <td data-bbox="950 997 1079 1081">MONTH.....</td> <td data-bbox="1079 997 1174 1081"> <table border="1"> <tr><td> </td><td> </td></tr> </table> </td> </tr> <tr> <td data-bbox="950 1081 1079 1144">YEAR..</td> <td data-bbox="1079 1081 1174 1144"> <table border="1"> <tr> <td>1</td><td>9</td><td> </td><td> </td> </tr> </table> </td> </tr> </table>	DAY.....	<table border="1"> <tr><td> </td><td> </td></tr> </table>			MONTH.....	<table border="1"> <tr><td> </td><td> </td></tr> </table>			YEAR..	<table border="1"> <tr> <td>1</td><td>9</td><td> </td><td> </td> </tr> </table>	1	9		
DATE WEIGHTED AND MEASURED	<table border="1"> <tr> <td data-bbox="950 913 1079 997">DAY.....</td> <td data-bbox="1079 913 1174 997"> <table border="1"> <tr><td> </td><td> </td></tr> </table> </td> </tr> <tr> <td data-bbox="950 997 1079 1081">MONTH.....</td> <td data-bbox="1079 997 1174 1081"> <table border="1"> <tr><td> </td><td> </td></tr> </table> </td> </tr> <tr> <td data-bbox="950 1081 1079 1144">YEAR..</td> <td data-bbox="1079 1081 1174 1144"> <table border="1"> <tr> <td>1</td><td>9</td><td> </td><td> </td> </tr> </table> </td> </tr> </table>	DAY.....	<table border="1"> <tr><td> </td><td> </td></tr> </table>			MONTH.....	<table border="1"> <tr><td> </td><td> </td></tr> </table>			YEAR..	<table border="1"> <tr> <td>1</td><td>9</td><td> </td><td> </td> </tr> </table>	1	9				
DAY.....	<table border="1"> <tr><td> </td><td> </td></tr> </table>																
MONTH.....	<table border="1"> <tr><td> </td><td> </td></tr> </table>																
YEAR..	<table border="1"> <tr> <td>1</td><td>9</td><td> </td><td> </td> </tr> </table>	1	9														
1	9																
930	<table border="1"> <tr> <td data-bbox="609 1144 950 1438">RESULT</td> <td data-bbox="950 1144 1174 1438"> <table border="1"> <tr> <td data-bbox="950 1144 1174 1228">MEASURED.....1</td> </tr> <tr> <td data-bbox="950 1228 1174 1291">NOT PRESENT.....3</td> </tr> <tr> <td data-bbox="950 1291 1174 1354">REFUSED.....4</td> </tr> <tr> <td data-bbox="950 1354 1174 1438">OTHER _____ 7 (SPECIFY)</td> </tr> </table> </td> </tr> </table>	RESULT	<table border="1"> <tr> <td data-bbox="950 1144 1174 1228">MEASURED.....1</td> </tr> <tr> <td data-bbox="950 1228 1174 1291">NOT PRESENT.....3</td> </tr> <tr> <td data-bbox="950 1291 1174 1354">REFUSED.....4</td> </tr> <tr> <td data-bbox="950 1354 1174 1438">OTHER _____ 7 (SPECIFY)</td> </tr> </table>	MEASURED.....1	NOT PRESENT.....3	REFUSED.....4	OTHER _____ 7 (SPECIFY)										
RESULT	<table border="1"> <tr> <td data-bbox="950 1144 1174 1228">MEASURED.....1</td> </tr> <tr> <td data-bbox="950 1228 1174 1291">NOT PRESENT.....3</td> </tr> <tr> <td data-bbox="950 1291 1174 1354">REFUSED.....4</td> </tr> <tr> <td data-bbox="950 1354 1174 1438">OTHER _____ 7 (SPECIFY)</td> </tr> </table>	MEASURED.....1	NOT PRESENT.....3	REFUSED.....4	OTHER _____ 7 (SPECIFY)												
MEASURED.....1																	
NOT PRESENT.....3																	
REFUSED.....4																	
OTHER _____ 7 (SPECIFY)																	
931	<table border="1"> <tr> <td data-bbox="609 1438 950 1549">NAME OF MEASURER</td> <td data-bbox="950 1438 1174 1549"> <table border="1"> <tr> <td> </td><td> </td><td> </td><td> </td> </tr> </table> </td> </tr> <tr> <td colspan="2" data-bbox="609 1501 1174 1549">_____</td> </tr> </table>	NAME OF MEASURER	<table border="1"> <tr> <td> </td><td> </td><td> </td><td> </td> </tr> </table>					_____									
NAME OF MEASURER	<table border="1"> <tr> <td> </td><td> </td><td> </td><td> </td> </tr> </table>																

INTERVIEWER'S OBSERVATIONS
(To be filled after completing interview)

COMMENTS ABOUT WOMAN

COMMENTS ON SPECIFIC QUESTIONS

ANY OTHER COMMENTS

SUPERVISOR'S OBSERVATIONS

NAME OF THE SUPERVISOR:
DATE:

EDITOR'S OBSERVATIONS

NAME OF THE EDITOR:
DATE:

CALENDAR

INSTRUCTIONS:
 ONLY ONE CODE SHOULD APPEAR IN ANY BOX.
 FOR COLUMNS 4 AND 5a-5b ALL MONTHS
 SHOULD BE FILLED IN.

COLUMN 1:
 BIRTHS AND PREGNANCIES

N MONTHS OUT OF WEDLOCK

COLUMN 3:
 O NOT MARRIED

COLUMN 4: MOVES AND TYPES OF COMMUNITIES

X CHANGE OF COMMUNITY
 1 PROVINCE CENTRE
 2 DISTRICT CENTRE
 3 SUBDISTRICT/VILLAGE
 4 ABROAD

COLUMN 5a AND 5b: MOVES AND PROVINCES

X CHANGE OF COMMUNITY
 01-80 PROVINCE CODES
 90 ABROAD

		1	2	3	4	5a	5b		
12 DEC	01	N		O				01	DEC
11 NOV	02	N		O				02	NOV
10 OCT	03	N		O				03	OCT
09 SEP	04	N		O				04	SEP
1 08 AUG	05	N		O				05	AUG 1
9 07 JUL	06	N		O				06	JUL 9
9 06 JUN	07	N		O				07	JUN 9
8 05 MAY	08	N		O				08	MAY 8
04 APR	09	N		O				09	APR
03 MAR	10	N		O				10	MAR
02 FEB	11	N		O				11	FEB
01 JAN	12	N		O				12	JAN

12 DEC	13	N		O				13	DEC
11 NOV	14	N		O				14	NOV
10 OCT	15	N		O				15	OCT
09 SEP	16	N		O				16	SEP
1 08 AUG	17	N		O				17	AUG 1
9 07 JUL	18	N		O				18	JUL 9
9 06 JUN	19	N		O				19	JUN 9
7 05 MAY	20	N		O				20	MAY 7
04 APR	21	N		O				21	APR
03 MAR	22	N		O				22	MAR
02 FEB	23	N		O				23	FEB
01 JAN	24	N		O				24	JAN

12 DEC	25	N		O				25	DEC
11 NOV	26	N		O				26	NOV
10 OCT	27	N		O				27	OCT
09 SEP	28	N		O				28	SEP
1 08 AUG	29	N		O				29	AUG 1
9 07 JUL	30	N		O				30	JUL 9
9 06 JUN	31	N		O				31	JUN 9
6 05 MAY	32	N		O				32	MAY 6
04 APR	33	N		O				33	APR
03 MAR	34	N		O				34	MAR
02 FEB	35	N		O				35	FEB
01 JAN	36	N		O				36	JAN

12 DEC	37	N		O				37	DEC
11 NOV	38	N		O				38	NOV
10 OCT	39	N		O				39	OCT
09 SEP	40	N		O				40	SEP
1 08 AUG	41	N		O				41	AUG 1
9 07 JUL	42	N		O				42	JUL 9
9 06 JUN	43	N		O				43	JUN 9
5 05 MAY	44	N		O				44	MAY 5
04 APR	45	N		O				45	APR
03 MAR	46	N		O				46	MAR
02 FEB	47	N		O				47	FEB
01 JAN	48	N		O				48	JAN

12 DEC	49	N		O				49	DEC
11 NOV	50	N		O				50	NOV
10 OCT	51	N		O				51	OCT
09 SEP	52	N		O				52	SEP
1 08 AUG	53	N		O				53	AUG 1
9 07 JUL	54	N		O				54	JUL 9
9 06 JUN	55	N		O				55	JUN 9
4 05 MAY	56	N		O				56	MAY 4
04 APR	57	N		O				57	APR
03 MAR	58	N		O				58	MAR
02 FEB	59	N		O				59	FEB
01 JAN	60	N		O				60	JAN

12 DEC	61	N		O				61	DEC
11 NOV	62	N		O				62	NOV
10 OCT	63	N		O				63	OCT
09 SEP	64	N		O				64	SEP
1 08 AUG	65	N		O				65	AUG 1
9 07 JUL	66	N		O				66	JUL 9
9 06 JUN	67	N		O				67	JUN 9
3 05 MAY	68	N		O				68	MAY 3
04 APR	69	N		O				69	APR
03 MAR	70	N		O				70	MAR
02 FEB	71	N		O				71	FEB
01 JAN	72	N		O				72	JAN

PROVINCE CODES

01 ADANA	21 DIYARBAKIR	41 KOCAELI	61 TRABZON
02 ADIYAMAN	22 EDİRNE	42 KONYA	62 TUNCELI
03 AFYON	23 ELAZIĞ	43 KÖTAHYA	63 ŞANLIURFA
04 AĞRI	24 ERZİNCAN	44 MALATYA	64 UŞAK
05 AMASYA	25 ERZURUM	45 MANISA	65 VAN
06 ANKARA	26 ESKİŞEHİR	46 K.MARAŞ	66 YOZGAT
07 ANTALYA	27 GAZİANTEP	47 MARDİN	67 ZONGULDAK
08 ARTVIN	28 GİRESUN	48 MUĞLA	68 AKSARAY
09 AYDIN	29 GÖMÜŞHANE	49 MUŞ	69 BAYBURT
10 BALIKESİR	30 HAKKARİ	50 NEVŞEHİR	70 KARAMAN
11 BİLECİK	31 HATAY	51 NİĞDE	71 KIRIKKALE
12 BİNGÖL	32 İSPARTA	52 ORDU	72 BATMAN
13 BİTLİS	33 İÇEL	53 RİZE	73 ŞİRNAK
14 BOLU	34 İSTANBUL	54 SAKARYA	74 BARTIN
15 BURDUR	35 İZMİR	55 SAMSUN	75 ARDAHAN
16 BURSA	36 KARS	56 SIIRT	76 İĞDIR
17 ÇANAKKALE	37 KASTAMONU	57 SİNOP	77 YALOVA
18 ÇANKIRI	38 KAYSERİ	58 SİVAS	78 KARABÜK
19 ÇORUM	39 KIRKLARELİ	59 TEKİRDAĞ	79 KİLİS
20 DENİZLİ	40 KİRŞEHİR	60 TOKAT	80 OSMANIYE
90 OTHER COUNTRY			

CONVERSION OF YEARS OF BIRTH FROM
RUMİ CALENDAR TO MILADI CALENDAR
YEARS :

RUMİ YEAR + 584 = MILADI YEAR

HACETTEPE UNIVERSITY INSTITUTE OF POPULATION STUDIES
 1998 TURKISH DEMOGRAPHIC AND HEALTH SURVEY
 HUSBAND QUESTIONNAIRE

IDENTIFICATION			
CLUSTER NO.....	<input type="text"/>	<input type="text"/>	PROVINCE _____
HOUSEHOLD NO.....	<input type="text"/>	<input type="text"/>	DISTRICT _____
REGION.....	<input type="text"/>	<input type="text"/>	SUBDISTRICT _____
URBAN (1) / RURAL (2).....	<input type="text"/>	<input type="text"/>	VILLAGE _____
			QUARTER _____
			STREET _____ NO _____

NAME-SURNAME OF MAN _____	LINE NO OF MAN	<input type="text"/>
	LINE NO OF WOMAN	<input type="text"/>

INTERVIEWER VISITS				
	1	2	3	FINAL VISIT
DATE (DAY-MONTH)	____	____	____	<input type="text"/>
INTERVIEWER'S NAME-SURNAME	_____	_____	_____	<input type="text"/>
RESULT (*)	____	____	____	<input type="text"/>
NEXT DAY-MONTH VISIT HOUR	____	____		TOTAL NO OF VISITS <input type="text"/>

(*) RESULT CODES :

1 COMPLETED	5 PARTLY COMPLETED
2 NOT AT HOME	
3 POSTPONED	7 OTHER _____
4 REFUSED	(SPECIFY)

SUPERVISOR _____	FIELD EDITOR _____	KEYED BY _____
DAY-MONTH <input type="text"/>	DAY-MONTH <input type="text"/>	DAY-MONTH <input type="text"/>

Age	Year of birth	
	Has not had birthday in 1998	Has already had birthday in 1998
	Does not know	
0	1997	--
1	1996	1997
2	1995	1996
3	1994	1995
4	1993	1994
5	1992	1993
6	1991	1992
7	1990	1991
8	1989	1990
9	1988	1989
10	1987	1988
11	1986	1987
12	1985	1986
13	1984	1985
14	1983	1984
15	1982	1983
16	1981	1982
17	1980	1981
18	1979	1980
19	1978	1979
20	1977	1978
21	1976	1977
22	1975	1976
23	1974	1975
24	1973	1974
25	1972	1973
26	1971	1972
27	1970	1971
28	1969	1970
29	1968	1969
30	1967	1968
31	1966	1967
32	1965	1966
33	1964	1965
34	1963	1964

Age	Year of birth	
	Has not had birthday in 1998	Has already had birthday in 1998
	Does not know	
35	1962	1963
36	1961	1962
37	1960	1961
38	1959	1960
39	1958	1959
40	1957	1958
41	1956	1957
42	1955	1956
43	1954	1955
44	1953	1954
45	1952	1953
46	1951	1952
47	1950	1951
48	1949	1950
49	1948	1949
50	1947	1948
51	1946	1947
52	1945	1946
53	1944	1945
54	1943	1944
55	1942	1943
56	1941	1942
57	1940	1941
58	1939	1940
59	1938	1939
60	1937	1938
61	1936	1937
62	1935	1936
63	1934	1935
64	1933	1934
65	1932	1933
66	1931	1932
67	1930	1931
68	1929	1930
69	1928	1929

SECTION 1. RESPONDENT'S BACKGROUND

101	RECORD THE TIME	HOUR - MINUTES..... <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>
102A	First I would like to ask some questions about you and the place you lived. For most of the time until you were 12 years old, where did you live ? _____ (NAME OF THE PLACE) Was it a city centre, district centre, a subdistrict or village ? Or did you live abroad ?	PROVINCE CENTRE.....1 DISTRICT CENTRE.....2 SUBDISTRICT OR VILLAGE.....3 ABROAD.....4 → 103
102B	In which province is this place now ? RECORD THE NAME AND CODE OF THE PROVINCE.	NAME OF PROVINCE <input type="text"/> _____ PROVINCE CODE
103	How long have you been living continuously in (NAME OF PLACE OF INTERVIEW) ?	YEARS..... <input type="text"/> <input type="text"/> ALWAYS.....95 VISITOR.....96 → 105
104A	Where did you live before you moved here ? _____ (NAME OF THE PLACE) Was that a city centre, district centre, a subdistrict or village ? Or did you live abroad ?	PROVINCE CENTRE.....1 DISTRICT CENTRE.....2 SUBDISTRICT OR VILLAGE.....3 ABROAD.....4 → 104C
104B	In which province is this place now ? RECORD THE NAME AND CODE OF THE PROVINCE.	NAME OF PROVINCE <input type="text"/> _____ PROVINCE CODE

104C	<p>What was the main reason for you to move here ?</p>	<p>PERSONAL REASONS MARRIAGE.....01 EDUCATION.....02 LOOKING FOR JOB.....03 CHANGE OF JOB/APPOINTMENT.....04 RETURN TO HOMELAND.....05 SPOUSAL REASONS TO ACCOMPANY HUSBAND.....06 CHANGE OF JOB/APPOINTMENT..07 LOOKING FOR JOB.....08 FAMILIAL REASONS TO JOIN PARENTS.....09 CHANGE OF JOB/APPOINTMENT..10 LOOKING FOR JOB.....11 OTHER _____ 96 (SPECIFY)</p>
105	<p>In what month and year were you born ?</p>	<p>MONTH..... <input type="text"/> <input type="text"/> DON'T KNOW MONTH.....98 YEAR..... <input type="text"/> 1 <input type="text"/> 9 <input type="text"/> <input type="text"/> DON'T KNOW YEAR.....9998</p>
106	<p>How old are you exactly ? What age have you completed ?</p> <p>CHECK ANSWERS TO 105 AND 106 USING AGE-BIRTH YEAR TABLE. IF INCONSISTENT PROBE AND CORRECT AGE INFORMATION MUST BE RECORDED !</p>	<p>AGE IN COMPLETED YEARS..... <input type="text"/> <input type="text"/></p>
107	<p>Have you ever attended school ?</p>	<p>YES.....1 NO.....2 →114</p>
108	<p>What is the highest level you have attended ?</p>	<p>PRIMARY.....1 SECONDARY.....2 HIGH SCHOOL.....3 UNIVERSITY.....4</p>
109A	<p>What is the highest grade you have completed at that level ?</p>	<p>GRADE..... <input type="text"/></p>
109B	<p>Did you graduate from this school ?</p>	<p>YES.....1 NO.....2</p>

113	CHECK 108 :	ATTENDED PRIMARY <input type="checkbox"/>	ATTENDED SECONDARY OR HIGHER <input type="checkbox"/>	→115
-----	-------------	---	---	------

114	Can you read and understand a letter or newspaper easily, with difficulty, or not at all ?	EASILY.....1 WITH DIFFICULTY.....2 NOT AT ALL.....3 →116A
-----	--	---

115	How frequent do you read a newspaper or magazine ?	EVERY DAY/ALMOST EVERY DAY.....1 ONCE-TWICE A WEEK.....2 RARELY/SELDOM.....3 NEVER/ALMOST NEVER.....4
-----	--	--

116A	What is your mother tongue ? RECORD ONLY ONE RESPONSE.	TURKISH.....01 KURDISH AND DIALECTS (KURMANCI, GORANI, ZAZA ETC)...02 ARABIC.....03 GREEK.....04 ARMANIAN.....05 HEBREW(LADINO).....06 CIRCASSIAN.....07 GEORGIAN.....08 LAZ LANGUAGE.....09 PERSIAN.....10 BULGARIAN.....11 RUMANIAN.....12 SERBIAN.....13 ENGLISH.....14 GERMAN.....15 OTHER _____ 96 (SPECIFY)
------	---	--

116B	In addition to your mother tongue, which language(s) can you speak ? RECORD ALL MENTIONED.	TURKISH.....A KURDISH AND DIALECTS (KURMANCI, GORANI, ZAZA ETC)...B ARABIC.....C GREEK.....D ARMANIAN.....E HEBREW(LADINO).....F CIRCASSIAN.....G GEORGIAN.....H LAZ LANGUAGE.....I PERSIAN.....J BULGARIAN.....K RUMANIAN.....L SERBIAN.....M ENGLISH.....N GERMAN.....O OTHER _____ U (SPECIFY) KNOWS NO OTHER LANGUAGE.....Y
------	---	---

116C	What is (was) your mother's and father's mother tongue ? USE THE CODES IN 116A.	MOTHER..... FATHER.....
------	--	--------------------------------

117	What is your religion ? IF THE ANSWER IS "MUSLIM" PROBE FOR RELIGIOUS SECT AND CIRCLE APPROPRIATE CODE.	MUSLIM SUNNI.....01 ALAWI.....02 OTHER _____ 03 (SPECIFY) CHRISTIAN.....06 JEWISH.....07 NO RELIGION.....10 OTHER _____ 96 (SPECIFY)
-----	--	---

118	CHECK Q.4 IN THE HOUSEHOLD QUESTIONNAIRE :		
	THE MAN INTERVIEWED IS NOT A USUAL RESIDENT <input type="checkbox"/>	THE MAN INTERVIEWED IS A USUAL RESIDENT <input type="checkbox"/>	→130

119A	Now I would like to ask about the place in which you usually live. What is the name of the place in which you usually live ? _____ (NAME OF PLACE) Is that a city centre, a district centre, a subdistrict or village, or are you living abroad ?	PROVINCE CENTRE.....1 DISTRICT CENTRE.....2 SUBDISTRICT OR VILLAGE.....3 ABROAD.....4 →120A
------	---	--

119B	In which province is that located ?	NAME OF PROVINCE _____ PROVINCE CODE _____
------	-------------------------------------	---

--	--

120A	Now I would like to ask about the household you usually live. How many persons do usually live in your house ?	NUMBER.....
------	---	-------------

--	--

120B	Does the house you usually live belong to a household member, is it rented from someone else, is it a lodging, or do you just live here without having to pay anything ?	OWNED BY A HOUSEHOLD MEMBER....1 RENTED.....2 LODGING.....3 NO MONEY PAID.....4 OTHER _____ 7 (SPECIFY)
------	--	--

121A	<p>What is the source of drinking water for your household ?</p>	<p>PIPED WATER PIPED WATER IN HOUSE/GARDEN...11 →122A PUBLIC PIPED WATER OUTSIDE HOUSE/GARDEN.....12 WELL WATER WELL IN RESIDENCE/GARDEN.....21 →122A PUBLIC WELL.....22 SURFACE WATER SPRING WATER PIPED INTO HOUSE/GARDEN.....31 →122A SPRING/PUBLIC .32 FOUNTAIN..... RIVER/STREAM/ .33 POND/LAKE/DAM.....41 →122A RAINWATER.....51 TANKER TRUCK.....61 →122A BOTTLED WATER/DEMIJOHN/PET WTR.71 WATER STATION..... 96 OTHER _____ (SPECIFY)</p>
121B	<p>How long does it take you to go there, get water, and come back ?</p>	<p>MINUTES..... <input type="text"/> <input type="text"/> <input type="text"/> ON PREMISES.....996</p>
122A	<p>Is the toilet inside the house or outside ?</p>	<p>NO FACILITY/BUSH/FIELD.....0 →123 INSIDE.....1 OUTSIDE.....2 INSIDE AND OUTSIDE.....3 OTHER _____ 7 (SPECIFY)</p>
122B	<p>What type of facility is used in the toilet ?</p> <p>ASK WHETHER THE TOILET IS CONNECTED TO DRAINAGE SYSTEM</p> <p>IF IT IS CONNECTED : Ask if the toilet is used only by this household or shared with another household ?</p> <p>IF IT IS NOT CONNECTED: What is the facility for excreta disposal ?</p>	<p>CONNECTED TO DRAINAGE SYSTEM USED ONLY BY THIS HOUSEHOLD...11 SHARED WITH ANOTHER HOUSEHOLD.12 PIT OPEN PIT.....21 CLOSED PIT.....22 NO FACILITY.....31 OTHER _____ 96 (SPECIFY)</p>
123	<p>What is the source of heating in winter for your house ?</p>	<p>RADIATOR (CENTRAL HEATING).....1 RADIATOR (PRIVATE).....2 NATURAL GAS STOVE.....3 STOVE (COAL/WOOD).....4 OTHER _____ 7 (SPECIFY)</p>

124	How many rooms in your household are used for sleeping ?	ROOMS USED FOR SLEEPING... <input type="text"/>																																										
125	What is the main material of the floor ?	<p>NATURAL FLOOR</p> <p>EARTH.....11</p> <p>RUDIMENTARY</p> <p>WOOD PLANKS.....21</p> <p>FINISHED FLOOR</p> <p>PARQUET/POLISHED WOOD.....31</p> <p>KARO.....32</p> <p>CEMENT.....34</p> <p>CARPET/WALL TO WALL CARPET..35</p> <p>MARLEY.....36</p> <p>MOSAIC.....37</p> <p>OTHER _____ 96</p> <p>(SPECIFY)</p>																																										
126	I would like to get an estimate of the total income that enters your household each month. Is the total amount of money earned by the members of your household in a month:																																											
126A	More than 100 million ?	<p>YES.....1</p> <p>NO.....2 → 126D</p>																																										
126B	More than 300 million ?	<p>YES.....1</p> <p>NO.....2 → 127</p>																																										
126C	More than 500 million ?	<p>YES.....1</p> <p>NO.....2 → 127</p>																																										
126D	Less than 50 million ?	<p>YES.....1</p> <p>NO.....2</p>																																										
127	Do you have the following in your household ?	<table border="1"> <thead> <tr> <th></th> <th>YES</th> <th>NO</th> </tr> </thead> <tbody> <tr> <td>Refrigerator</td> <td>1</td> <td>2</td> </tr> <tr> <td>Gas or electric oven</td> <td>1</td> <td>2</td> </tr> <tr> <td>Dishwasher</td> <td>1</td> <td>2</td> </tr> <tr> <td>Washing machine</td> <td>1</td> <td>2</td> </tr> <tr> <td>Vacuum cleaner</td> <td>1</td> <td>2</td> </tr> <tr> <td>Television</td> <td>1</td> <td>2</td> </tr> <tr> <td>Video</td> <td>1</td> <td>2</td> </tr> <tr> <td>Camera</td> <td>1</td> <td>2</td> </tr> <tr> <td>Music set with CD player</td> <td>1</td> <td>2</td> </tr> <tr> <td>Telephone</td> <td>1</td> <td>2</td> </tr> <tr> <td>Cellular Telephone</td> <td>1</td> <td>2</td> </tr> <tr> <td>Car (excluding tractors, taxis, etc.)</td> <td>1</td> <td>2</td> </tr> <tr> <td>Computer</td> <td>1</td> <td>2</td> </tr> </tbody> </table>		YES	NO	Refrigerator	1	2	Gas or electric oven	1	2	Dishwasher	1	2	Washing machine	1	2	Vacuum cleaner	1	2	Television	1	2	Video	1	2	Camera	1	2	Music set with CD player	1	2	Telephone	1	2	Cellular Telephone	1	2	Car (excluding tractors, taxis, etc.)	1	2	Computer	1	2
	YES	NO																																										
Refrigerator	1	2																																										
Gas or electric oven	1	2																																										
Dishwasher	1	2																																										
Washing machine	1	2																																										
Vacuum cleaner	1	2																																										
Television	1	2																																										
Video	1	2																																										
Camera	1	2																																										
Music set with CD player	1	2																																										
Telephone	1	2																																										
Cellular Telephone	1	2																																										
Car (excluding tractors, taxis, etc.)	1	2																																										
Computer	1	2																																										

130	<p>Now I would like to ask you questions about working.</p> <p>Are you currently working?</p>	<p>YES.....1 →132</p> <p>NO.....2</p>
-----	---	---------------------------------------

131	<p>Have you worked in any job in the last 12 months ?</p>	<p>YES.....1</p> <p>NO.....2 →141</p>
-----	---	---------------------------------------

132	<p>What type of work are/were you doing ? What kind of job are/were you in ?</p> <p>DO NOT FILL IN THE BOXES.</p>	<table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td style="width: 20px; height: 20px;"></td> <td style="width: 20px; height: 20px;"></td> <td style="width: 20px; height: 20px;"></td> </tr> </table> <p>_____</p> <p>_____</p> <p>_____</p>			

132A	<p>Do/did you pay social security when doing this job ?</p> <p>IF YES: According to which schedule ?</p>	<p>NO.....0</p> <p>SSK.....1</p> <p>EMEKLİ SANDIĞI.....2</p> <p>BAĞ-KUR.....3</p> <p>OTHER _____ 7</p> <p style="text-align: center;">(SPECIFY)</p> <p>DON'T KNOW.....8</p>
------	--	---

132B	<p>CHECK 132, CIRCLE THE APPROPRIATE WORK CODE.</p>	<p>AGRICULTURE.....01</p> <p>ANIMAL HUSBANDRY.....02</p> <p>AGRICULTURE AND A.HUSBANDRY...03</p> <p>FORESTRY.....04</p> <p>FISHERY.....05</p> <p>MINING.....06</p> <p>MANUFACTURE INDUSTRY (FOOD)...07</p> <p>MANUFACTURE INDUSTRY (TEXTILE).08</p> <p>MANUFACTURE INDUSTRY (METAL)...09</p> <p>MANUFACTURE INDUSTRY (OTHER)...10</p> <p>ELECTRICITY/GAS/WATER.....11</p> <p>CONSTRUCTION.....12</p> <p>WHOLESALE/RETAIL TRADE.....13</p> <p>HOTEL AND RESTAURANT.....14</p> <p>COMMUNICATION/TRANSPORTATION...15</p> <p>HEALTH.....16</p> <p>EDUCATION/CULTURE.....17</p> <p>PUBLIC ADMINISTRATION.....18</p> <p>TOURISM.....19</p> <p>BANKING/INSURANCE.....20</p> <p>OTHER SOCIAL SERVICES.....21</p> <p>PERSONAL SERVICES.....22</p> <p>OTHER _____ 96</p> <p style="text-align: center;">(SPECIFY)</p>
------	---	---

132C	CHECK 132 , CIRCLE THE APPROPRIATE CODE FOR THE POSITION AT WORK	EMPLOYER EMPLOYING TEN OR MORE THAN TEN PERSONS.....01 SMALL EMPLOYER EMPLOYING LESS THAN TEN PERSONS.....02 SALARIED (GOVERNMENT OFFICIAL).03 WAGED (PUBLIC SECTOR).....04 WAGED (PRIVATE SECTOR).....05 FOR HIS OWN.....06 UNPAID FAMILY LABOURER.....07 OTHER _____ 96 (SPECIFY)
133	CHECK 132B : AGRICULTURE (01) OR AGRICULTURE AND ANIMAL HUSBANDRY (03)	OTHER WORK CODES <input type="checkbox"/> _____ → 135
134	Whom does the land you work on belong to ?	HIS LAND.....01 FAMILY LAND.....02 RENTED LAND.....03 SOMEONE ELSE'S LAND.....04 STATE LAND.....05 OTHER _____ 96 (SPECIFY)
135	Do you usually work throughout the year, or do you work seasonally, or only once in a while ?	THROUGHOUT THE YEAR.....1 → 137 SEASONALLY/PART OF THE YEAR.NDE.2 ONCE IN A WHILE.....3 → 138
136	During the last 12 months, how many months did you work ?	NUMBER OF MONTHS..... <input type="text"/>
137	During the last 12 months, how many days a week did you usually work (in the months that you worked) ?	NUMBER OF DAYS..... <input type="text"/> → 139
138	During the last 12 months, approximately, how many days did you work ?	NUMBER OF DAYS..... <input type="text"/>
139	Do/did you earn cash for your work ? PROBE : Do/did you make money for your work ?	YES.....1 NO.....2 → 142

140A How much do you usually earn for this work ?

PROBE : Is this by the day, by the week, or by the month ?

PER HOUR.....1							
PER DAY.....2							
PER WEEK.....3							
PER MONTH....4							
PER YEAR.....5							

OTHER _____ 999996 → 142
(SPECIFY)

140B CIRCLE THE APPROPRIATE CODE FOR THE GIVEN QUANTITY (AMOUNT) IN 140A .

MILLION.....1 → 142
THOUSAND.....2

141 You told currently you are not working.
What is the main reason that you are not working ?

RETIRED.....01
INCOME FROM INTEREST, RENT ETC..02
NOT ABLE TO WORK (HANDICAPPED
SICK, TOO OLD ETC.).....03
CAN'T FIND/LOOKING FOR JOB.....04

OTHER _____ 96
(SPECIFY)

142 Are you covered by health insurance ?

IF YES : According to which schedule ?

.....0
SSK.....1
EMEKLİ SANDIĞI.....2
BAĞ-KUR.....3
PRIVATE INSURANCE.....4
GREEN CARD.....5
OTHER _____ 6
(SPECIFY)

SECTION.2 FERTILITY

201	<p>Now I would like to ask you questions about your children. Here I consider your own/self children. Have you evcr had any children born alive ?</p>	<p>YES.....1 NO.....2 → 206</p>									
202	<p>Do you have any sons or daughter who are now living with you ?</p>	<p>YES.....1 NO.....2 → 204</p>									
203	<p>How many sons live with you ? And how many daughters live with you ? IF NONE, RECORD "00".</p>	<p>SONS AT HOME..... <table border="1" data-bbox="1209 682 1274 745"><tr><td></td><td></td></tr><tr><td></td><td></td></tr></table> DAUGHTERS AT HOME..... <table border="1" data-bbox="1209 745 1274 808"><tr><td></td><td></td></tr><tr><td></td><td></td></tr></table></p>									
204	<p>Do you have any sons or daughters who are alive but do not live with you(live elsewhere) ?</p>	<p>YES.....1 NO.....2 → 206</p>									
205	<p>How many sons are alive but do not live with you ? And how many daughters are alive but do not live with you ? IF NONE, RECORD "00".</p>	<p>SONS ELSEWHERE..... <table border="1" data-bbox="1209 1018 1274 1081"><tr><td></td><td></td></tr><tr><td></td><td></td></tr></table> DAUGHTERS ELSEWHERE..... <table border="1" data-bbox="1209 1081 1274 1144"><tr><td></td><td></td></tr><tr><td></td><td></td></tr></table></p>									
206	<p>Have you ever had a boy or a girl born alive but later died ? IF NO. PROBE: Any baby who cried or showed signs of life but only survived a few hours or days ?</p>	<p>YES.....1 NO.....2 → 208</p>									
207	<p>In all, how many boys have died ? And how many girls have died ? IF NONE, PROBE "00".</p>	<p>BOYS DEAD..... <table border="1" data-bbox="1209 1428 1274 1491"><tr><td></td><td></td></tr><tr><td></td><td></td></tr></table> GIRLS DEAD..... <table border="1" data-bbox="1209 1491 1274 1554"><tr><td></td><td></td></tr><tr><td></td><td></td></tr></table></p>									
208	<p>SUM ANSWERS TO 203, 205, AND 207, AND ENTER TOTAL. IF NONE, RECORD "00".</p>	<p>TOTAL..... <table border="1" data-bbox="1209 1648 1274 1711"><tr><td></td><td></td></tr></table></p>									

209	CHECK 208 : Just to make sure that I have this right: you have had in TOTAL ___ births during your life. Is that correct ? YES <input type="checkbox"/> NO <input type="checkbox"/>	PROBE AND CORRECT 201-208 AS NECESSARY.
210	CHECK 208 : HAS CHILD <input type="checkbox"/> HAS NO CHILD <input type="checkbox"/>	→ 301
210A	In what month and year was your last child born ?	MONTH..... YEAR.....
210B	CHECK 210A, LAST CHILD BORN : AFTER JANUARY 1993 <input type="checkbox"/> BEFORE JANUARY 1993 <input type="checkbox"/>	→ 301
211	Before your last child was born, did you want to have child then, did you want to wait until later, or did you not want to have any more children at all ?	THEN.....1 LATER.....2 NOT WANT MORE CHILDREN.....3

SECTION 3. CONTRACEPTION

Now I would like to talk about family planning. There are various methods that a married couple can use to avoid pregnancy.

- CIRCLE CODE '1' IN IN Q. 301 FOR EACH METHOD MENTIONED SPONTANEOUSLY.
- THEN PROCEED DOWN COLUMN 302, READING THE NAME AND DESCRIPTION OF EACH METHOD NOT MENTIONED SPONTANEOUSLY AND ASK WHETHER SHE HAS HEARD THE METHOD. CIRCLE CODE '2' IF METHOD IS RECOGNIZED AND CODE '3' IF NOT RECOGNIZED.
- THEN FOR EACH METHOD WITH CODE '1' OR '2' CIRCLED IN 301 OR 302, ASK 303.
- AFTER ASKING ABOUT ALL METHODS PROCEED TO 304.

	SPON- TANEOUS YES	302 Have you ever heard this method ? PROBED YES NO		303 Have you ever used this method ?
01 PILL Woman can avoid a pregnancy by taking a pill every day.	1	2	3	YES.....1 NO.....2
02 IUD Women can have the so called spiral or IUD placed in them by a doctor or a nurse.	1	2	3	YES.....1 NO.....2
03 INJECTABLES Woman can have an injection by a doctor or a nurse which stops them from becoming pregnant for a certain period of time.	1	2	3	YES.....1 NO.....2
04 NORPLANT Woman can have small rods placed in their arm and this can prevent pregnancy for several years.	1	2	3	YES.....1 NO.....2
05 DIAPHRAGM, FOAM, JELLY Woman can place a sponge, suppository, diaphragm, jelly, or cream inside themselves before intercourse.	1	2	3	YES.....1 NO.....2
06 CONDOM Men can put a rubber sheath on their penis during sexual intercourse.	1	2	3	YES.....1 NO.....2
07 TUBAL LIGATION Women can have an operation of tubal ligation to avoid having any more children.	1	2	3	Has (had) your wife ever have such an operation ? YES.....1 NO.....2
08 MALE STERILIZATION Men can have an operation called vasectomy so that their wives would not get pregnant.	1	2	3	Have you ever had such an operation to avoid having any more children ? YES.....1 NO.....2

301 Which ways or methods have you heard ?	SPON-TANEOUS YES	302 Have you ever heard this method ? PROBED YES NO		303 Have you ever used this method ?
09 RHYTHM Some couples can avoid having sexual intercourse on certain days of the month when the woman is more likely to become pregnant.	1	2	3	YES.....1 NO.....2
10 WITHDRAWAL Some men pull out during sexual intercourse before climax.	1	2	3	YES.....1 NO.....2
11 Have you heard of any other method that women or men can use to avoid pregnancy ?	1	(SPECIFY)		YES.....1 NO.....2
		(SPECIFY)		YES.....1 NO.....2
		(SPECIFY)		YES.....1 NO.....2
304 CHECK 303 : NOT A SINGLE "YES" (NEVER USED) <input type="checkbox"/> AT LEAST ONE "YES" (EVER USED) <input type="checkbox"/> →307				
305 Have you or your wife ever used anything or tried in any way to delay or avoid getting pregnant ?				YES.....1 NO.....2 →309
306 What have you used or done ? CORRECT 303 AND 304 (AND 302 IF NECESSARY).				
307 Are you or your wife currently doing something or using any method to delay or avoid getting pregnant ?				YES.....1 NO.....2 →309
308 Which method are you using ?				PILL.....01 IUD.....02 INJECTABLES.....03 NORPLANT.....04 DIAPHRAGM/FOAM/JELLY.....05 CONDOM.....06 →310 TUBAL LIGATION.....07 MALE STERILIZATION.....08 RHYTHM.....09 WITHDRAWAL.....10 OTHER _____ 96 →310 (SPECIFY)

308A	<p>You said you are currently using withdrawal. Do you use only withdrawal or use it together with any other method ?</p>	<p>YES, TOGETHER WITH OTHER METHOD1 NO, ONLY WITHDRAWAL.....2 → 310</p>
308B	<p>Which method is that ?</p> <p>DO NOT MAKE ANY CORRECTION AT 308 DUE TO THE METHOD MENTIONED HERE. TAKE WITHDRAWAL AS THE CURRENTLY USED METHOD AND SKIP TO 310.</p>	<p>PILL.....01 IUD.....02 INJECTABLES.....03 NORPLANT.....04 DIAPHRAGM/FOAM/JELLY.....05 CONDOM.....06 → 310 TUBAL LIGATION.....07 MALE STERILIZATION.....08 RHYTHM.....09</p> <p>OTHER _____ 96 (SPECIFY)</p>
309	<p>What is the main reason you are not using a method of contraception to avoid pregnancy ?</p>	<p>FERTILITY-RELATED REASONS</p> <p>NOT HAVING SEX.....21 INFREQUENT SEX.....22 WIFE IN MENOPAUSAL/HAD HYSTERECTOMY..23 WIFE SUBFECUND/INFECUND24 WIFE POSTPARTUM/ BREASTFEEDING.....25 WANTS (MORE) CHILDREN.....26 WIFE PREGNANT.....27 HIMSELF INFECUND.....28</p> <p>OPPOSITION TO USE</p> <p>HIMSELF OPPOSED.....31 WIFE OPPOSED.....32 OTHERS OPPOSED.....33 RELIGIOUS REASONS.....34</p> <p>LACK OF KNOWLEDGE</p> <p>KNOWS NO METHOD.....41 KNOWS NO SOURCE.....42</p> <p>METHOD-RELATED REASONS</p> <p>HEALTH CONCERNS.....51 SIDE EFFECTS.....52 LACK OF ACCESS/TOO FAR.....53 COST TOO MUCH.....54 INCONVENIENT TO USE.....55 INTERFERES WITH BODY'S NORMAL PROCESSES.....56</p> <p>LEFT THE CHOICE TO WOMAN.....61</p> <p>OTHER _____ 96 (SPECIFY)</p> <p>DON'T KNOW.....98</p>

310	CHECK 308 :	USING WITHDRAWAL <input type="checkbox"/>	USING OTHER METHOD <input type="checkbox"/>	→ 401
-----	-------------	--	--	-------

311	Now let's talk a little more about withdrawal. You told you are using withdrawal. Who prefers to use it, you, your wife or you and your wife together ?	HIMSELF.....1 WIFE.....2 TOGETHER.....3
-----	--	---

312	Do you think withdrawal is a convenient method for preventing pregnancy ? Does it sufficiently prevent pregnancy ?	YES.....1 NO.....2 DON'T KNOW.....8
-----	--	---

313	Do you think using withdrawal is easy or difficult ?	EASY.....1 DIFFICULT.....2 DON'T KNOW.....8
-----	--	---

314	What is the main reason that you are not using any other method but withdrawal ? DO NOT FILL IN THE BOXES.	<table style="width: 100%; border-collapse: collapse;"> <tr> <td style="border-bottom: 1px solid black; width: 80%;"></td> <td style="width: 20%; border: 1px solid black; text-align: center;"> </td> </tr> <tr> <td style="border-bottom: 1px solid black;"></td> <td style="border: 1px solid black; text-align: center;"> </td> </tr> <tr> <td style="border-bottom: 1px solid black;"></td> <td style="border: 1px solid black; text-align: center;"> </td> </tr> </table>						

SECTION 4. MARRIAGE

401	How many times did you marry ? Once or more than once ?	ONCE.....1 MORE THAN ONCE.....2
402A	Some men can be married with more than one woman at the same time. Is it the case for you ?	YES.....1 NO.....2 → 403
402B	Currently, how many wives do you have ?	NUMBER OF WIVES..... <input type="text"/>
403	WRITE THE LINE NO. OF THE WIFE FROM THE HOUSEHOLD QUESTIONNAIRE. IF THE WIFE DOESN'T LIVE AT HOME, WRITE 00. THE NUMBER IN THE BOX SHOULD BE EQUAL TO THE NUMBER OF WOMEN IN 402B.	<input type="text"/> <input type="text"/> <input type="text"/>
408	CHECK 401 : MARRIED ONCE <input type="checkbox"/> ↓ In what month and year did you marry (started living with) your wife ? MARRIED MORE THAN ONCE <input type="checkbox"/> ↓ Now, let's talk about your first wife. In what month and year did you marry (started living with) your wife ?	MONTH..... <input type="text"/> DON'T KNOW MONTH.....98 YEAR..... <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> → 409A DON'T KNOW YEAR.....9998
409	How old were you when you started living with your (first) wife ?	AGE..... <input type="text"/>
409A	How old was your (first) wife when you started living with her ?	AGE..... <input type="text"/>
411	CHECK 301 AND 302 : KNOWS CONDOM <input type="checkbox"/> ↓ Did you use condom in your last sexual intercourse ? DOES NOT KNOW CONDOM <input type="checkbox"/> ↓ Some men use rubber sheath during the sexual intercourse. Did you use condom in your last sexual intercourse ?	YES.....1 NO.....2

412	Do you know where you can obtain condom ?	YES.....1 NO.....2	→415		
413	Which place is that ? <hr/> (NAME OF PLACE)	PUBLIC SECTOR GOVERNMENT/SAMPLE HOSPITAL.....A MATERNITY HOUSE.....B MOTHER-CHILD HEALTH AND FAMILY PLANNING (MCHPP) CENTRE.....C HEALTH CENTRE.....D HEALTH HOUSE.....E SSK HOSPITAL/DISPENSARY.....F OTHER PUBLIC SECTOR _____ ..G (SPECIFY) PRIVATE SECTOR PRIVATE HOSPITAL.....H PRIVATE POLYCLINIC.....I PRIVATE DOCTOR.....J PRIVATE MIDWIFE/NURSE.....K PHARMACY.....L OTHER PRIVATE SECTOR _____ ..M (SPECIFY) UNIVERSITY HOSPITAL.....N COMMUNITY VOLUNTEERS/ ASSOCIATION/FOUNDATION.....O MARKET/SHOP.....P FRIEND/RELATIVE/NEIGHBOUR.....R OTHER _____ T (SPECIFY)			
415	How old were you when you had your first sexual intercourse ?	AGE..... <table border="1" data-bbox="1128 1354 1193 1417" style="display: inline-table; vertical-align: middle;"> <tr> <td style="width: 20px; height: 20px;"></td> <td style="width: 20px; height: 20px;"></td> </tr> </table>			96

SECTION 5. FERTILITY PREFERENCES

503	Is your wife (or one of your wives) currently pregnant ?	YES.....1 NO.....2 NOT SURE.....8
-----	--	---

→505

504	At the time your wife became pregnant, did you want this pregnancy then, did you want to wait until later, or did you not want to have any more children at all ?	THEN.....1 LATER.....2 NOT WANT MORE CHILDREN.....3
-----	---	---

505	CHECK 503 : WIFE CURRENTLY NOT PREGNANT OR NOT SURE <input type="checkbox"/> WIFE PREGNANT <input type="checkbox"/> Now I have some questions about the future. Would you like to have (a/another) child or would you prefer not to have any (more) children ?	HAVE (A/ANOTHER) CHILD.....1 NO MORE/NONE.....2 WIFE CAN NOT GET PREGNANT.....3 HIMSELF CAN NOT HAVE CHILD.....4 UNDECIDED/DON'T KNOW.....8
-----	---	---

→507

506	CHECK 503 : WIFE NOT PREGNANT OR NOT SURE <input type="checkbox"/> WIFE PREGNANT <input type="checkbox"/> How long would you like to wait from now before the birth of (a/another) child ?	MONTHS.....1 YEARS.....2 SOON/NOW.....993 WIFE CAN NOT GET PREGNANT.....994 OTHER.....996 (SPECIFY) DON'T KNOW.....998
-----	---	--

507	CHECK 308 : USING ANY METHOD ? NOT CURRENTLY USING <input type="checkbox"/> CURRENTLY USING <input type="checkbox"/>	→512
-----	--	------

508	Do you think you will use a method to delay or avoid pregnancy within the next 12 months ?	YES.....1 NO.....2 DON'T KNOW.....8
-----	--	---

→510

509	Do you think you will use a method to delay or avoid pregnancy at any time in the future ?	YES.....1 NO.....2 DON'T KNOW.....8
-----	--	---

→511

510 Which method would you prefer to use ?

- PILL.....01
- IUD.....02
- INJECTABLES.....03
- NORPLANT.....04
- DIAPHRAGM/FOAM/JELLY.....05
- CONDOM.....06
- TUBAL LIGATION.....07
- MALE STERILIZATION.....08 →512
- RHYTHM.....09
- WITHDRAWAL.....10
- OTHER _____ 96
- (SPECIFY)
- NOT SURE.....98

511 What is the main reason that you think you will never use a method ?

- FERTILITY RELATED REASONS
- INFREQUENT SEX.....22
- WIFE IN MENOPAUSAL/HAD HYSTERECTOMY..23
- WIFE SUBFECUND/INFECUND24
- WIFE POSTPARTUM/ BREASTFEEDING.....25
- WANTS (MORE) CHILDREN.....26
- WIFE PREGNANT.....27
- HISSELF INFECUND.....28
- OPPOSITION TO USE
- HIMSELF OPPOSED.....31
- WIFE OPPOSED.....32
- OTHERS OPPOSED.....33
- RELIGIOUS REASONS.....34
- LACK OF KNOWLEDGE
- KNOWS NO METHOD.....41
- KNOWS NO SOURCE.....42
- METHOD-RELATED REASONS
- HEALTH CONCERNS.....51
- SIDE EFFECTS.....52
- LACK OF ACCESS/TOO FAR.....53
- COST TOO MUCH.....54
- INCONVENIENT TO USE.....55
- INTERFERES WITH BODY'S NORMAL PROCESSES.....56
- LEFT THE CHOICE TO WOMAN.....61
- OTHER _____ 96
- (SPECIFY)
- DON'T KNOW.....98

512 CHECK 202 AND 204 :

HAS LIVING CHILDREN

↓
If you could go back to the time you did not have any children and could choose exactly the number of children to have in your whole life, how many would that be ?

NO LIVING CHILDREN

↓
If you could choose exactly the number of children to have in your whole life, how many would that be ?

NUMBER.....

OTHER _____ 96 →514
(SPECIFY)

513	How many of these children would you like to be boys, how many would you like to be girls and for how many? would it not matter ?	<table border="0"> <tr> <td colspan="2"></td> <td style="text-align: right;">BOYS</td> </tr> <tr> <td>NUMBER.....</td> <td></td> <td><input type="text"/></td> </tr> <tr> <td>OTHER _____</td> <td></td> <td style="text-align: right;">96</td> </tr> <tr> <td></td> <td></td> <td style="text-align: center;">(SPECIFY)</td> </tr> <tr> <td colspan="2"></td> <td style="text-align: right;">GIRLS</td> </tr> <tr> <td>NUMBER.....</td> <td></td> <td><input type="text"/></td> </tr> <tr> <td>OTHER _____</td> <td></td> <td style="text-align: right;">96</td> </tr> <tr> <td></td> <td></td> <td style="text-align: center;">(SPECIFY)</td> </tr> <tr> <td colspan="2"></td> <td style="text-align: right;">EITHER</td> </tr> <tr> <td>NUMBER.....</td> <td></td> <td><input type="text"/></td> </tr> <tr> <td>OTHER _____</td> <td></td> <td style="text-align: right;">96</td> </tr> <tr> <td></td> <td></td> <td style="text-align: center;">(SPECIFY)</td> </tr> </table>			BOYS	NUMBER.....		<input type="text"/>	OTHER _____		96			(SPECIFY)			GIRLS	NUMBER.....		<input type="text"/>	OTHER _____		96			(SPECIFY)			EITHER	NUMBER.....		<input type="text"/>	OTHER _____		96			(SPECIFY)
		BOYS																																				
NUMBER.....		<input type="text"/>																																				
OTHER _____		96																																				
		(SPECIFY)																																				
		GIRLS																																				
NUMBER.....		<input type="text"/>																																				
OTHER _____		96																																				
		(SPECIFY)																																				
		EITHER																																				
NUMBER.....		<input type="text"/>																																				
OTHER _____		96																																				
		(SPECIFY)																																				
514	Do you approve of couples using a method to avoid getting pregnant ?	<table border="0"> <tr> <td>APPROVE.....</td> <td style="text-align: right;">1</td> </tr> <tr> <td>DISAPPROVE.....</td> <td style="text-align: right;">2</td> </tr> <tr> <td>NO OPINION.....</td> <td style="text-align: right;">3</td> </tr> </table>	APPROVE.....	1	DISAPPROVE.....	2	NO OPINION.....	3																														
APPROVE.....	1																																					
DISAPPROVE.....	2																																					
NO OPINION.....	3																																					
515	Is it acceptable or not acceptable to you for information on family planning to be provided: On the radio ? On the television ? In newspapers/magazines ? In secondary schools ? In high schools ?	<table border="0"> <thead> <tr> <th></th> <th style="text-align: center;">ACCEPT- ABLE</th> <th style="text-align: center;">NOT ACCEPT- ABLE</th> <th style="text-align: center;">DO NOT KNOW</th> </tr> </thead> <tbody> <tr> <td>RADIO.....</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> <td style="text-align: center;">8</td> </tr> <tr> <td>TELEVISION.....</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> <td style="text-align: center;">8</td> </tr> <tr> <td>NEWSP./MAGZ.....</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> <td style="text-align: center;">8</td> </tr> <tr> <td>SECONDARY.....</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> <td style="text-align: center;">8</td> </tr> <tr> <td>HIGH.....</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> <td style="text-align: center;">8</td> </tr> </tbody> </table>		ACCEPT- ABLE	NOT ACCEPT- ABLE	DO NOT KNOW	RADIO.....	1	2	8	TELEVISION.....	1	2	8	NEWSP./MAGZ.....	1	2	8	SECONDARY.....	1	2	8	HIGH.....	1	2	8												
	ACCEPT- ABLE	NOT ACCEPT- ABLE	DO NOT KNOW																																			
RADIO.....	1	2	8																																			
TELEVISION.....	1	2	8																																			
NEWSP./MAGZ.....	1	2	8																																			
SECONDARY.....	1	2	8																																			
HIGH.....	1	2	8																																			
516	In the last few months have you heard about family On the radio ? On the television ? In newspaper or magazine? From a poster ? From brochures or leaflets ? From billboards or cloth posters ? Any other place ?	<table border="0"> <thead> <tr> <th></th> <th style="text-align: center;">YES</th> <th style="text-align: center;">NO</th> </tr> </thead> <tbody> <tr> <td>RADIO.....</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> </tr> <tr> <td>TELEVISION.....</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> </tr> <tr> <td>NEWSPAPER OR MAGAZINE.....</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> </tr> <tr> <td>POSTER.....</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> </tr> <tr> <td>BROCHURES/LEAFLETS.....</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> </tr> <tr> <td>BILLBOARDS/CLOTH POSTERS.....</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> </tr> <tr> <td>ANOTHER PLACE.....</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> </tr> </tbody> </table>		YES	NO	RADIO.....	1	2	TELEVISION.....	1	2	NEWSPAPER OR MAGAZINE.....	1	2	POSTER.....	1	2	BROCHURES/LEAFLETS.....	1	2	BILLBOARDS/CLOTH POSTERS.....	1	2	ANOTHER PLACE.....	1	2												
	YES	NO																																				
RADIO.....	1	2																																				
TELEVISION.....	1	2																																				
NEWSPAPER OR MAGAZINE.....	1	2																																				
POSTER.....	1	2																																				
BROCHURES/LEAFLETS.....	1	2																																				
BILLBOARDS/CLOTH POSTERS.....	1	2																																				
ANOTHER PLACE.....	1	2																																				
518	In the last few months have you discussed the practice of family planning with friends, neighbours, or relatives ?	<table border="0"> <tr> <td>YES.....</td> <td style="text-align: right;">1</td> </tr> <tr> <td>NO.....</td> <td style="text-align: right;">2</td> </tr> </table>	YES.....	1	NO.....	2																																
YES.....	1																																					
NO.....	2																																					

→520

519	<p>With whom ?</p> <p>Anyone else ?</p> <p>RECORD ALL MENTIONED.</p>	<p>WIFE/PARTNER.....A</p> <p>MOTHER.....B</p> <p>FATHER.....C</p> <p>SISTER(S).....D</p> <p>BROTHER(S).....E</p> <p>DAUGHTER.....F</p> <p>SON.....G</p> <p>MOTHER-IN-LAW.....H</p> <p>FRIEND/RELATIVE.....I</p> <p>NEIGHBOUR.....J</p> <p>OTHER _____ U</p> <p>(SPECIFY)</p>
520	<p>Now I want to ask you about your wife's views on family planning.</p> <p>Do you think that your wife approves of couples using a method to avoid pregnancy ?</p>	<p>APPROVES.....1</p> <p>DISAPPROVES.....2</p> <p>DON'T KNOW.....8</p>
523	<p>Do you think your wife wants the same number of children that you want, or does she want more or fewer than you want ?</p>	<p>SAME NUMBER.....1</p> <p>MORE CHILDREN.....2</p> <p>FEWER CHILDREN.....3</p> <p>DON'T KNOW.....8</p>
524	<p>Do you think that the use of family planning is against religion ?</p>	<p>YES.....1</p> <p>SOME METHODS ARE.....2</p> <p>NO.....3</p> <p>NOT RELIGIOUS/NO RELIGION.....4</p> <p>DON'T KNOW.....8</p> <p style="text-align: right;">→801A</p>
525	<p>Which method(s) do you think (are) is against religion ?</p> <p>RECORD ALL MENTIONED.</p>	<p>PILL.....A</p> <p>IUD.....B</p> <p>INJECTABLES.....C</p> <p>NORPLANT.....D</p> <p>DIAPHRAGM/FOAM/JELLY.....E</p> <p>CONDOM.....F</p> <p>TUBAL LIGATION.....G</p> <p>MALE STERILIZATION.....H</p> <p>RHYTHM.....I</p> <p>WITHDRAWAL.....J</p> <p>SEXUAL ABSTINENCE.....K</p> <p>OTHER METHOD _____ L</p> <p>(SPECIFY)</p> <p>INDUCED ABORTION.....M</p>

SECTION 8. SEXUALLY TRANSMITTED DISEASES AND AIDS

801A	<p>Now I would like you talk you about sexually transmitted diseases.</p> <p>Have you ever heard of sexually transmitted diseases ?</p>	<p>YES.....1</p> <p>NO.....2 → 801L</p>
------	---	---

801B	<p>Which diseases have you heard ?</p> <p>RECORD ALL MENTIONED.</p>	<p>SYPHILIS.....A</p> <p>GONORE.....B</p> <p>AIDS.....C</p> <p>WART/SORE IN GENITAL ORGANS.....D</p> <p>FUNGUS.....E</p> <p>OTHER _____ U</p> <p>(SPECIFY)</p> <p>OTHER _____ V</p> <p>(SPECIFY)</p> <p>DOESN'T KNOW THE NAMES.....X</p>
------	---	--

801K	<p>CHECK 801B :</p> <p>NOT MENTIONED AIDS <input type="checkbox"/> MENTIONED AIDS <input type="checkbox"/> → 802A</p>		
------	--	--	--

801L	<p>Have you ever heard of an illness called AIDS ?</p>	<p>YES.....1</p> <p>NO.....2 → 901</p>
------	--	--

802A	<p>From which sources of information have you learned most about AIDS ?</p> <p>Any other sources ?</p> <p>RECORD ALL MENTIONED.</p>	<p>RADIO.....A</p> <p>TV.....B</p> <p>NEWSPAPERS/MAGAZINES.....C</p> <p>PAMPHLETS/POSTERS.....D</p> <p>HEALTH WORKERS.....E</p> <p>MOSQUES.....F</p> <p>SCHOOLS/TEACHERS.....G</p> <p>FRIENDS/RELATIVES.....I</p> <p>WORK PLACE.....J</p> <p>OTHER _____ U</p> <p>(SPECIFY)</p>
------	---	---

802B	<p>Through what ways a person is transmitted AIDS ?</p> <p>Any other ways ?</p> <p>RECORD ALL MENTIONED.</p>	<p>SEXUAL RELATION.....A</p> <p>SEXUAL RELATION WITH MORE THAN ONE PARTNER.....B</p> <p>SEXUAL RELATION WITH A PROSTITUTE.....C</p> <p>NOT USING CONDOM.....D</p> <p>HOMOSEXUAL RELATION.....E</p> <p>BLOOD TRANSFUSION.....F</p> <p>INJECTION.....G</p> <p>KISSING.....H</p> <p>MOSQUITO BITE.....I</p> <p>OTHER _____ U (SPECIFY)</p> <p>OTHER _____ V (SPECIFY)</p> <p>DON'T KNOW.....X</p>
803	<p>Is there anything a person can do to avoid getting AIDS or the virus that causes AIDS ?</p>	<p>YES.....1</p> <p>NO.....2</p> <p>DON'T KNOW.....8 → 807</p>
804	<p>What can a person do ?</p> <p>Any other ways ?</p> <p>RECORD ALL MENTIONED.</p>	<p>SAFE SEX.....A</p> <p>ABSTAIN FROM SEX.....D</p> <p>USE CONDOMS.....C</p> <p>HAVE ONLY ONE SEX PARTNER.....D</p> <p>AVOID SEX WITH PROSTITUTES.....E</p> <p>AVOID SEX WITH HOMOSEXUALS.....F</p> <p>AVOID BLOOD TRANSFUSIONS.....G</p> <p>AVOID INJECTIONS.....H</p> <p>AVOID KISSING.....I</p> <p>AVOID MOSQUITO BITES.....J</p> <p>SEEK PROTECTION FROM TRADITIONAL HEALER.....K</p> <p>OTHER _____ U (SPECIFY)</p> <p>OTHER _____ V (SPECIFY)</p> <p>DON'T KNOW.....X</p>
807	<p>Is it possible for a healthy-looking person to have the AIDS virus ?</p>	<p>YES.....1</p> <p>NO.....2</p> <p>DON'T KNOW.....8</p>
808A	<p>Do you think that persons with AIDS almost never die from the disease, sometimes die, or almost always die from the disease ?</p>	<p>ALMOST NEVER.....1</p> <p>SOMETIMES.....2</p> <p>ALMOST ALWAYS.....3</p> <p>DON'T KNOW.....8</p>

808B	Is there a medical treatment for AIDS ?	YES.....1 NO.....2 DON'T KNOW.....8
808C	Is AIDS transmitted from mother to child ?	YES.....1 NO.....2 DON'T KNOW.....8
808D	Do you know anyone with HIV virus or anyone died from AIDS ?	YES.....1 NO.....2 DON'T KNOW.....8

SECTION 9. ATTITUDE

904	<p>Woman can sometimes make things that may annoy or make angry her husband. Now I will talk about some situations. Can you please tell me if the husband can have the right for beating his wife in those situations?</p>	<p>YES NO DK</p>
	<p>If she burns the food ? If she neglects the care of children ? If she argues with her husband ? If she talks with other men ? If she spends the money needlessly ? If she refuses to have sexual intercourse ?</p>	<p>BURN FOOD.....1 2 8 NEGLECT CHILD CARE....1 2 8 ARGUE WITH HUSBAND....1 2 8 TALK OTHER MEN.....1 2 8 SPEND NEEDLESSLY.....1 2 8 SEXUAL INTERCOURSE....1 2 8</p>
905	<p>Now I will read you a few sentences. I would like to learn what you think about the ideas in these sentences. Do you agree or disagree with the following:</p>	<p>AGREES DISAGREES HAS NO IDEA</p>
	<p>The important decisions in the family should be made by the male family members.</p>	<p>1 2 8</p>
	<p>Men are usually wiser than women.</p>	<p>1 2 8</p>
	<p>A woman should not argue with her husband even if she does not share the same view with him.</p>	<p>1 2 8</p>
	<p>It is better for the male child to be educated than the female child.</p>	<p>1 2 8</p>

912	RECORD THE TIME.	HOUR..... MINUTES.....	<table border="1" style="width: 20px; height: 20px; margin-left: auto; margin-right: auto;"> <tr><td style="width: 10px; height: 10px;"></td><td style="width: 10px; height: 10px;"></td></tr> <tr><td style="width: 10px; height: 10px;"></td><td style="width: 10px; height: 10px;"></td></tr> </table>				
913	PRESENCE OF OTHERS DURING INTERVIEW : CIRCLE ALL APPROPRIATE ALTERNATIVES.	NO ONE.....A CHILDREN UNDER 10.....B WIFE.....C MOTHER IN LAW.....D RESPONDENT'S MOTHER.....E OTHER MEN.....P OTHER WOMEN.....G					
914	WAS THE INTERVIEW INTERRUPTED ? IF YES, FOR HOW LONG , APPROXIMATELY ? (IN MINUTES)	NO.....000 YES.....1	<table border="1" style="width: 20px; height: 20px; margin-left: auto; margin-right: auto;"> <tr><td style="width: 10px; height: 10px;"></td><td style="width: 10px; height: 10px;"></td></tr> </table>				
915	WHAT LANGUAGE WAS USED DURING THE INTERVIEW ?	TURKISH.....1 KURDISH.....2 ARABIC.....3 OTHER.....4 (SPECIFY)	→917				
916	WAS AN INTERPRETER USED DURING THE INTERVIEW ?	YES.....1 NO.....2					
917	WHAT IS THE RELIABILITY OF THE RESPONSES, IN YOUR OPINION ?	POOR.....1 FAIR.....2 GOOD.....3 VERY GOOD.....4					

NAME OF THE EDITOR:
DATE:

EDITOR'S OBSERVATIONS

NAME OF THE SUPERVISOR:
DATE:

SUPERVISOR'S OBSERVATIONS

ANY OTHER COMMENTS

COMMENTS ON SPECIFIC QUESTIONS

COMMENTS ABOUT HUSBAND

INTERVIEWER'S OBSERVATIONS
(To be filled after completing interview)

PROVINCE CODES

01 ADANA	21 DIYARBAKIR	41 KOCAELI	61 TRABZON
02 ADIYAMAN	22 EDİRNE	42 KONYA	62 TUNCELI
03 AFYON	23 ELAZIĞ	43 KÜTAHYA	63 ŞANLIURFA
04 AĞRI	24 ERZINCAN	44 MALATYA	64 UŞAK
05 AMASYA	25 ERZURUM	45 MANISA	65 VAN
06 ANKARA	26 ESKİŞEHİR	46 K.MARAŞ	66 YOZGAT
07 ANTALYA	27 GAZİANTEP	47 MARDİN	67 ZONGULDAK
08 ARTVIN	28 GİRESUN	48 MUĞLA	68 AKSARAY
09 AYDIN	29 GÜMÜŞHANE	49 MUŞ	69 BAYBURT
10 BALIKESİR	30 HAKKARİ	50 NEVŞEHİR	70 KARAMAN
11 BİLECİK	31 HATAY	51 NİĞDE	71 KIRIKKALE
12 BİNGÖL	32 İSPARTA	52 ORDU	72 BATMAN
13 BİTLİS	33 İÇEL	53 RİZE	73 ŞIRNAK
14 BOLU	34 İSTANBUL	54 SAKARYA	74 BARTIN
15 BURDUR	35 İZMİR	55 SAMSUN	75 ARDAHAN
16 BURSA	36 KARS	56 SİRT	76 İGDIR
17 ÇANAKKALE	37 KASTAMONU	57 SİNOP	77 YALOVA
18 ÇANKIRI	38 KAYSERİ	58 SIVAS	78 KARABÜK
19 ÇORUM	39 KIRKLARELİ	59 TEKİRDAĞ	79 KİLİS
20 DENİZLİ	40 KIRŞEHİR	60 TOKAT	80 OSMANIYE

90 OTHER COUNTRY

CONVERSION OF YEARS OF BIRTH FROM
RUMİ CALENDAR TO MILADI CALENDAR
YEARS :

RUMİ YEAR + 584 = MILADI YEAR